

IMO 2013 Solution Notes

COMPILED BY EVAN CHEN

April 15, 2019

This is an compilation of solutions for the 2013 IMO. Some of the solutions are my own work, but many are from the official solutions provided by the organizers (for which they hold any copyrights), and others were found on the Art of Problem Solving forums.

Corrections and comments are welcome!

Contents

0 Problems	2
1 IMO 2013/1, proposed by Japan	3
2 IMO 2013/2, proposed by Ivan Guo (AUS)	4
3 IMO 2013/3, proposed by Alexander A. Polyansky (RUS)	5
4 IMO 2013/4, proposed by Warut Suksompong and Potcharapol Suteparuk (THA)	7
5 IMO 2013/5, proposed by Bulgaria	9
6 IMO 2013/6, proposed by Russia	10

§0 Problems

1. Let k and n be positive integers. Prove that there exist positive integers m_1, \dots, m_k such that

$$1 + \frac{2^k - 1}{n} = \left(1 + \frac{1}{m_1}\right) \left(1 + \frac{1}{m_2}\right) \dots \left(1 + \frac{1}{m_k}\right).$$

2. A configuration of 4027 points in the plane is called *Colombian* if it consists of 2013 red points and 2014 blue points, and no three of the points of the configuration are collinear. By drawing some lines, the plane is divided into several regions. An arrangement of lines is *good* for a Colombian configuration if the following two conditions are satisfied:

- (i) No line passes through any point of the configuration.
- (ii) No region contains points of both colors.

Find the least value of k such that for any Colombian configuration of 4027 points, there is a good arrangement of k lines.

3. Let the excircle of triangle ABC opposite the vertex A be tangent to the side BC at the point A_1 . Define the points B_1 on CA and C_1 on AB analogously, using the excircles opposite B and C , respectively. Suppose that the circumcenter of triangle $A_1B_1C_1$ lies on the circumcircle of triangle ABC . Prove that triangle ABC is right-angled.
4. Let ABC be an acute triangle with orthocenter H , and let W be a point on the side \overline{BC} , between B and C . The points M and N are the feet of the altitudes drawn from B and C , respectively. Suppose ω_1 is the circumcircle of triangle BWN and X is a point such that \overline{WX} is a diameter of ω_1 . Similarly, ω_2 is the circumcircle of triangle CWM and Y is a point such that \overline{WY} is a diameter of ω_2 . Show that the points X, Y , and H are collinear.
5. Suppose a function $f : \mathbb{Q}_{>0} \rightarrow \mathbb{R}$ satisfies:
- (i) If $x, y \in \mathbb{Q}_{>0}$, then $f(x)f(y) \geq f(xy)$.
 - (ii) If $x, y \in \mathbb{Q}_{>0}$, then $f(x+y) \geq f(x) + f(y)$.
 - (iii) There exists a rational number $a > 1$ with $f(a) = a$.

Prove that f is the identity function.

6. Let $n \geq 3$ be an integer, and consider a circle with $n + 1$ equally spaced points marked on it. Consider all labellings of these points with the numbers $0, 1, \dots, n$ such that each label is used exactly once; two such labellings are considered to be the same if one can be obtained from the other by a rotation of the circle. A labelling is called *beautiful* if, for any four labels $a < b < c < d$ with $a + d = b + c$, the chord joining the points labelled a and d does not intersect the chord joining the points labelled b and c . Let M be the number of beautiful labellings, and let N be the number of ordered pairs (x, y) of positive integers such that $x + y \leq n$ and $\gcd(x, y) = 1$. Prove that $M = N + 1$.

§1 IMO 2013/1, proposed by Japan

Let k and n be positive integers. Prove that there exist positive integers m_1, \dots, m_k such that

$$1 + \frac{2^k - 1}{n} = \left(1 + \frac{1}{m_1}\right) \left(1 + \frac{1}{m_2}\right) \dots \left(1 + \frac{1}{m_k}\right).$$

By induction on $k \geq 1$. When $k = 1$ there is nothing to prove.

For the inductive step, if n is even, write

$$\frac{n + (2^k - 1)}{n} = \left(1 + \frac{1}{n + (2^k - 2)}\right) \cdot \frac{\frac{n}{2} + (2^{k-1} - 1)}{\frac{n}{2}}$$

and use inductive hypothesis on the second term. On the other hand if n is odd then write

$$\frac{n + (2^k - 1)}{n} = \left(1 + \frac{1}{n}\right) \cdot \frac{\frac{n+1}{2} + (2^{k-1} - 1)}{\frac{n+1}{2}}$$

and use inductive hypothesis on the second term.

§2 IMO 2013/2, proposed by Ivan Guo (AUS)

A configuration of 4027 points in the plane is called *Colombian* if it consists of 2013 red points and 2014 blue points, and no three of the points of the configuration are collinear. By drawing some lines, the plane is divided into several regions. An arrangement of lines is *good* for a Colombian configuration if the following two conditions are satisfied:

- (i) No line passes through any point of the configuration.
- (ii) No region contains points of both colors.

Find the least value of k such that for any Colombian configuration of 4027 points, there is a good arrangement of k lines.

The answer is $k \geq 2013$.

To see that $k = 2013$ is necessary, consider a regular 4026-gon and alternatively color the points red and blue, then place the last blue point anywhere in general position (it doesn't matter). Each side of the 4026 is a red-blue line segment which needs to be cut by one of the k lines, and each line can cut at most two of the segments.

Now, we prove that $k = 2013$ lines is sufficient. Consider the convex hull of all the points.

- If the convex hull has any red points, cut that red point off from everyone else by a single line. Then, for each of the remaining 2012 red points, break them into 1006 pairs arbitrarily, and for each pair $\{A, B\}$ draw two lines parallel to AB and very close to them.
- If the convex hull has two consecutive blue points, cut that red point off from everyone else by a single line. Then repeat the above construction for the remaining 2012 blue points.

The end.

§3 IMO 2013/3, proposed by Alexander A. Polyansky (RUS)

Let the excircle of triangle ABC opposite the vertex A be tangent to the side BC at the point A_1 . Define the points B_1 on CA and C_1 on AB analogously, using the excircles opposite B and C , respectively. Suppose that the circumcenter of triangle $A_1B_1C_1$ lies on the circumcircle of triangle ABC . Prove that triangle ABC is right-angled.


We ignore for now the given condition and prove the following important lemma.

Lemma

Let (AB_1C_1) meet (ABC) again at X . From $BC_1 = B_1C$ follows $XC_1 = XB_1$, and X is the midpoint of major arc \widehat{BC} .

Proof. This follows from the fact that we have a spiral similarity $\triangle XBC_1 \sim \triangle XCB_1$ which must actually be a spiral congruence since $BC_1 = B_1C$. \square

We define the arc midpoints Y and Z similarly, which lie on the perpendicular bisectors of $\overline{A_1C_1}$, $\overline{A_1B_1}$.



We now turn to the problem condition which asserts the circumcenter W of $\triangle A_1B_1C_1$ lies on (ABC) .

Claim — We may assume WLOG that $W = X$.

Proof. This is just configuration issues since we already knew that the arc midpoints both lie on (ABC) and the relevant perpendicular bisectors.

Suppose (WLOG) that $\angle B_1A_1C_1 > 90^\circ$ (since W lies (ABC) and hence outside $\triangle ABC$, hence outside $\triangle A_1B_1C_1$). Then A and X lie on the same side of line $\overline{B_1C_1}$, and since W is supposed to lie both on (ABC) and the perpendicular bisector of $\overline{B_1C_1}$ it follows $W = X$. \square

Consequently, \overline{XY} and \overline{XZ} are exactly the perpendicular bisectors of $\overline{A_1C_1}$, $\overline{A_1B_1}$. The rest is angle chase, the fastest one is

$$\begin{aligned}\angle A &= \angle C_1XB_1 = \angle C_1XA_1 + \angle A_1XB_1 = 2\angle YXA_1 + 2\angle A_1XZ \\ &= 2\angle YXZ = 180^\circ - \angle A\end{aligned}$$

which solves the problem.


Remark. Angle chasing is also possible even without the points Y and Z , though it takes much longer. Introduce the Bevan point V and use the fact that VA_1B_1C is cyclic (with diameter \overline{VC}) and similarly VA_1C_1B is cyclic; a calculation then gives $\angle CVB = 180^\circ - \frac{1}{2}\angle A$. Thus V lies on the circle with diameter $\overline{I_bI_c}$.

§4 IMO 2013/4, proposed by Warut Suksompong and Potcharapol Suteparuk (THA)

Let ABC be an acute triangle with orthocenter H , and let W be a point on the side \overline{BC} , between B and C . The points M and N are the feet of the altitudes drawn from B and C , respectively. Suppose ω_1 is the circumcircle of triangle BWN and X is a point such that \overline{WX} is a diameter of ω_1 . Similarly, ω_2 is the circumcircle of triangle CWM and Y is a point such that \overline{WY} is a diameter of ω_2 . Show that the points X, Y , and H are collinear.

We present two solutions, an elementary one and then an advanced one by moving points.

First solution, classical Let P be the second intersection of ω_1 and ω_2 ; this is the Miquel point, so P also lies on the circumcircle of AMN , which is the circle with diameter \overline{AH} .



We now contend:

Claim — Points P, H, X collinear. (Similarly, points P, H, Y are collinear.)

Proof using power of a point. By radical axis on $BNMC$, ω_1, ω_2 , it follows that A, P, W are collinear. We know that $\angle APH = 90^\circ$, and also $\angle XPW = 90^\circ$ by construction. Thus P, H, X are collinear. □

Proof using angle chasing. This is essentially Reim's theorem:

$$\angle NPH = \angle NAH = \angle BAH = \angle ABX = \angle NBX = \angle NPX$$

as desired. Alternatively, one may prove A, P, W are collinear by $\angle NPA = \angle NMA = \angle NMC = \angle BNC = \angle BNW = \angle NPW$. □

Second solution, by moving points Fix $\triangle ABC$ and vary W . Let ∞ be the point at infinity perpendicular to \overline{BC} for brevity.

By spiral similarity, the point X moves linearly on $\overline{B\infty}$ as W varies linearly on \overline{BC} . Similarly, so does Y . So in other words, the map

$$X \mapsto W \mapsto Y$$

is linear. However, the map

$$X \mapsto Y' \stackrel{\text{def}}{=} \overline{XH} \cap \overline{C\infty}$$

is linear too.

To show that these maps are the same, it suffices to check it thus at two points.

- When $W = B$, the circle (BNW) degenerates to the circle through B tangent to \overline{BC} , and $X = \overline{CN} \cap \overline{B\infty}$. We have $Y = Y' = C$.
- When $W = C$, the result is analogous.
- Although we don't need to do so, it's also easy to check the result if W is the foot from A since then $XHWB$ and $YHWC$ are rectangles.

§5 IMO 2013/5, proposed by Bulgaria

Suppose a function $f : \mathbb{Q}_{>0} \rightarrow \mathbb{R}$ satisfies:

- (i) If $x, y \in \mathbb{Q}_{>0}$, then $f(x)f(y) \geq f(xy)$.
- (ii) If $x, y \in \mathbb{Q}_{>0}$, then $f(x+y) \geq f(x) + f(y)$.
- (iii) There exists a rational number $a > 1$ with $f(a) = a$.

Prove that f is the identity function.

First, we dispense of negative situations by proving:

Claim — For any integer $n > 0$, we have $f(n) \geq n$.

Proof. Note by induction on (ii) we have $f(nx) \geq nf(x)$. Taking $(x, y) = (a, 1)$ in (i) gives $f(1) \geq 1$, and hence $f(n) \geq n$. \square

Claim — The f takes only positive values, and hence by (ii) is strictly increasing.

Proof, suggested by Gopal Goel. Let $p, q > 0$ be integers. Then $f(q)f(p/q) \geq f(p)$, and since both $\min(f(p), f(q)) > 0$ it follows $f(p/q) > 0$. \square

Claim — For any $x > 1$ we have $f(x) \geq x$.

Proof. Note that

$$f(x)^N \geq f(x^N) \geq f(\lfloor x^N \rfloor) \geq \lfloor x^N \rfloor > x^N - 1$$

for any integer N . Since m can be arbitrarily large, we conclude $f(x) \geq x$ for $x > 1$. \square

On the other hand, f has arbitrarily large fixed points (namely powers of a) so from (ii) we're essentially done. First, for $x > 1$ pick a large m and note

$$a^m = f(a^m) \geq f(a^m - x) + f(x) \geq (a^m - x) + x = a^m.$$

Finally, for $x \leq 1$ use

$$nf(x) = f(n)f(x) \geq f(nx) \geq nf(x)$$


for large n .

Note that $a > 1$ is essential; if $b \geq 1$ then $f(x) = bx^2$ works with unique fixed point $1/b \leq 1$.

§6 IMO 2013/6, proposed by Russia

Let $n \geq 3$ be an integer, and consider a circle with $n + 1$ equally spaced points marked on it. Consider all labellings of these points with the numbers $0, 1, \dots, n$ such that each label is used exactly once; two such labellings are considered to be the same if one can be obtained from the other by a rotation of the circle. A labelling is called *beautiful* if, for any four labels $a < b < c < d$ with $a + d = b + c$, the chord joining the points labelled a and d does not intersect the chord joining the points labelled b and c . Let M be the number of beautiful labellings, and let N be the number of ordered pairs (x, y) of positive integers such that $x + y \leq n$ and $\gcd(x, y) = 1$. Prove that $M = N + 1$.

First, here are half of the beautiful labellings up to reflection for $n = 6$, just for concreteness.



Abbreviate “beautiful labelling of points around a circle” to ring. Moreover, throughout the solution we will allow degenerate chords that join a point to itself; this has no effect on the problem statement.

The idea is to proceed by induction in the following way. A ring of $[0, n]$ is called *linear* if it is an arithmetic progression modulo $n + 1$. For example, the first two rings in the diagram and the last one are linear for $n = 6$, while the other three are not.

Of course we can move from any ring on $[0, n]$ to a ring on $[0, n - 1]$ by deleting n . We are going to prove that:

- Each linear ring on $[0, n - 1]$ yields exactly two rings of $[0, n]$, and
- Each nonlinear ring on $[0, n - 1]$ yields exactly one rings of $[0, n]$.

In light of the fact there are obviously $\varphi(n)$ linear rings on $[0, n]$, the conclusion will follow by induction.

We say a set of chords (possibly degenerate) is *pseudo-parallel* if for any three of them, one of them separates the two. (Intuitively, one can perturb the endpoints along the circle in order to make them parallel in Euclidean sense.) The main structure lemma is going to be:

Lemma

In any ring, the chords of sum k are pseudo-parallel.

Proof. By induction on k . You can assume one of the chords is $\{0, k\}$ else we can just shift. Moreover, discard all points greater than k (i.e. assume $n = k$). (Details omitted.) \square

Next, we give another characterization of linear rings.

Lemma

A ring on $[0, n - 1]$ is linear if and only if the point 0 does not lie between two chords of sum n .

Proof. It's obviously true for linear rings. Conversely, assume the property holds for some ring. Note that the chords with sum $n - 1$ are pseudo-parallel and encompass every point, so they are parallel. Similarly, the chords of sum n are parallel and encompass every point other than 0. So the map

$$t \mapsto n - t \mapsto (n - 1) - (n - t) = t + 1 \pmod{n}$$


is rotation as desired. \square

Lemma

Every nonlinear ring on $[0, n - 1]$ induces exactly one ring on $[0, n]$.

Proof. Because the chords of sum n are pseudo-parallel, there is at most one possibility for the location n .

Conversely, we claim that this works. The chords of sum n (and less than n) are OK by construction, so assume for contradiction that there exists $a, b, c \in \{1, \dots, n - 1\}$ such that $a + b = n + c$. Then, we can “reflect” them using the (pseudo-parallel) chords of length n to find that $(n - a) + (n - b) = 0 + (n - c)$, and the chords joining 0 to $n - c$ and $n - a$ to $n - b$ intersect, by definition.



This is a contradiction that the original numbers on $[0, n - 1]$ form a ring. \square

Lemma

Every linear ring on $[0, n - 1]$ induces exactly one ring on $[0, n]$.

Proof. Because the chords of sum n are pseudo-parallel, the point n must lie either directly to the left or right of 0. For the same reason as in the previous proof, both of them work. \square