

# USAMO 2002 Solution Notes

COMPILED BY EVAN CHEN

April 15, 2019

This is an compilation of solutions for the 2002 USAMO. Some of the solutions are my own work, but many are from the official solutions provided by the organizers (for which they hold any copyrights), and others were found on the Art of Problem Solving forums.

Corrections and comments are welcome!

## Contents

<a href="#">0 Problems</a>	2
<a href="#">1 USAMO 2002/1</a>	3
<a href="#">2 USAMO 2002/2</a>	4
<a href="#">3 USAMO 2002/3</a>	5
<a href="#">4 USAMO 2002/4</a>	6
<a href="#">5 USAMO 2002/5</a>	7
<a href="#">6 USAMO 2002/6</a>	8

## §0 Problems

- Let  $S$  be a set with 2002 elements, and let  $N$  be an integer with  $0 \leq N \leq 2^{2002}$ . Prove that it is possible to color every subset of  $S$  either black or white so that the following conditions hold:
  - the union of any two white subsets is white;
  - the union of any two black subsets is black;
  - there are exactly  $N$  white subsets.
- Let  $ABC$  be a triangle such that

$$\left(\cot \frac{A}{2}\right)^2 + \left(2 \cot \frac{B}{2}\right)^2 + \left(3 \cot \frac{C}{2}\right)^2 = \left(\frac{6s}{7r}\right)^2,$$

where  $s$  and  $r$  denote its semiperimeter and its inradius, respectively. Prove that triangle  $ABC$  is similar to a triangle  $T$  whose side lengths are all positive integers with no common divisors and determine these integers.

- Prove that any monic polynomial (a polynomial with leading coefficient 1) of degree  $n$  with real coefficients is the average of two monic polynomials of degree  $n$  with  $n$  real roots.
- Determine all functions  $f : \mathbb{R} \rightarrow \mathbb{R}$  such that

$$f(x^2 - y^2) = xf(x) - yf(y)$$

for all pairs of real numbers  $x$  and  $y$ .

- Let  $a, b$  be integers greater than 2. Prove that there exists a positive integer  $k$  and a finite sequence  $n_1, n_2, \dots, n_k$  of positive integers such that  $n_1 = a$ ,  $n_k = b$ , and  $n_i n_{i+1}$  is divisible by  $n_i + n_{i+1}$  for each  $i$  ( $1 \leq i < k$ ).
- I have an  $n \times n$  sheet of stamps, from which I've been asked to tear out blocks of three adjacent stamps in a single row or column. (I can only tear along the perforations separating adjacent stamps, and each block must come out of the sheet in one piece.) Let  $b(n)$  be the smallest number of blocks I can tear out and make it impossible to tear out any more blocks. Prove that there are real constants  $c$  and  $d$  such that

$$\frac{1}{7}n^2 - cn \leq b(n) \leq \frac{1}{5}n^2 + dn$$

for all  $n > 0$ .

**§1 USAMO 2002/1**

Let  $S$  be a set with 2002 elements, and let  $N$  be an integer with  $0 \leq N \leq 2^{2002}$ . Prove that it is possible to color every subset of  $S$  either black or white so that the following conditions hold:

- (a) the union of any two white subsets is white;
- (b) the union of any two black subsets is black;
- (c) there are exactly  $N$  white subsets.

---

Trivial, do it for any  $n$  other than 2002, noting white and black are dual. In this way we can "induct" – (either color  $N$  white or  $2^{2002} - N$  white kind of thing).

## §2 USAMO 2002/2

Let  $ABC$  be a triangle such that

$$\left(\cot \frac{A}{2}\right)^2 + \left(2 \cot \frac{B}{2}\right)^2 + \left(3 \cot \frac{C}{2}\right)^2 = \left(\frac{6s}{7r}\right)^2,$$

where  $s$  and  $r$  denote its semiperimeter and its inradius, respectively. Prove that triangle  $ABC$  is similar to a triangle  $T$  whose side lengths are all positive integers with no common divisors and determine these integers.

Let  $x = s - a$ ,  $y = s - b$ ,  $z = s - c$  in the usual fashion, then the equation reads

$$x^2 + 4y^2 + 9z^2 = \left(\frac{6}{7}(x + y + z)\right)^2.$$

However, by Cauchy-Schwarz, we have

$$\left(1 + \frac{1}{4} + \frac{1}{9}\right)(x^2 + 4y^2 + 9z^2) \geq (x + y + z)^2$$

with equality if and only if  $1 : \frac{1}{2} : \frac{1}{3} = x : 2y : 3z$ , id est  $x : y : z = 1 : \frac{1}{4} : \frac{1}{9} = 36 : 9 : 4$ . This is equivalent to  $y + z : z + x : x + y = 13 : 40 : 45$ .

**Remark.** You can tell this is not a geometry problem because you eliminate the cotangents right away to get an algebra problem... and then you realize the problem claims that one equation can determine three variables up to scaling, at which point you realize it has to be an inequality (otherwise degrees of freedom don't work). So of course, Cauchy-Schwarz...

### §3 USAMO 2002/3

Prove that any monic polynomial (a polynomial with leading coefficient 1) of degree  $n$  with real coefficients is the average of two monic polynomials of degree  $n$  with  $n$  real roots.

First,

#### Lemma

If  $p$  is a monic polynomial of degree  $n$ , and  $p(1)p(2) < 0$ ,  $p(2)p(3) < 0$ ,  $\dots$ ,  $p(n-1)p(n) < 0$  then  $p$  has  $n$  real roots.

*Proof.* The intermediate value theorem already guarantees the existence of  $n-1$  real roots.

The last root is obtained by considering cases on  $n \pmod{2}$ . If  $n$  is even, then  $p(1)$  and  $p(n)$  have opposite sign, while we must have either

$$\lim_{x \rightarrow -\infty} p(x) = \lim_{x \rightarrow \infty} p(x) = \pm\infty$$

so we get one more root. The  $n$  odd case is similar, with  $p(1)$  and  $p(n)$  now having the same sign, but  $\lim_{x \rightarrow -\infty} p(x) = -\lim_{x \rightarrow \infty} p(x)$  instead.  $\square$

Let  $f(n)$  be the monic polynomial and let  $M > 1000 \max_{t=1, \dots, n} |f(t)| + 1000$ . Then we may select reals  $a_1, \dots, a_n$  and  $b_1, \dots, b_n$  such that for each  $k = 1, \dots, n$ , we have

$$\begin{aligned} a_k + b_k &= 2f(k) \\ (-1)^k a_k &> M \\ (-1)^{k+1} b_k &> M. \end{aligned}$$

We may interpolate monic polynomials  $g$  and  $h$  through the  $a_k$  and  $b_k$  (if the  $a_k, b_k$  are selected “generically” from each other). Then one can easily check  $f = \frac{1}{2}(g+h)$  works.

**Remark.** This is like Cape Town all over again...

## §4 USAMO 2002/4

Determine all functions  $f : \mathbb{R} \rightarrow \mathbb{R}$  such that

$$f(x^2 - y^2) = xf(x) - yf(y)$$

for all pairs of real numbers  $x$  and  $y$ .

---

The answer is  $f(x) = cx$ ,  $c \in \mathbb{R}$  (these obviously work).

First, by putting  $x = 0$  and  $y = 0$  respectively we have

$$f(x^2) = xf(x) \quad \text{and} \quad f(-y^2) = -yf(y).$$

From this we deduce that  $f$  is odd, in particular  $f(0) = 0$ . Then, we can rewrite the given as  $f(x^2 - y^2) + f(y^2) = f(x^2)$ . Combined with the fact that  $f$  is odd, we deduce that  $f$  is additive (i.e.  $f(a + b) = f(a) + f(b)$ ).

**Remark** (Philosophy). At this point we have  $f(x^2) \equiv xf(x)$  and  $f$  additive, and everything we have including the given equation is a direct corollary of these two. So it makes sense to only focus on these two conditions.

Then

$$\begin{aligned} f((x+1)^2) &= (x+1)f(x+1) \\ \implies f(x^2) + 2f(x) + f(1) &= (x+1)f(x) + (x+1)f(1) \end{aligned}$$

which readily gives  $f(x) = f(1)x$ .

**§5 USAMO 2002/5**

Let  $a, b$  be integers greater than 2. Prove that there exists a positive integer  $k$  and a finite sequence  $n_1, n_2, \dots, n_k$  of positive integers such that  $n_1 = a$ ,  $n_k = b$ , and  $n_i n_{i+1}$  is divisible by  $n_i + n_{i+1}$  for each  $i$  ( $1 \leq i < k$ ).

---

Consider a graph  $G$  on the vertex set  $\{3, 4, \dots\}$  and with edges between  $v, w$  if  $v + w \mid vw$ ; the problem is equivalent to showing that  $G$  is connected.

First, note that  $n$  is connected to  $n(n-1)$ ,  $n(n-1)(n-2)$ , etc. up to  $n!$ . But for  $n > 2$ ,  $n!$  is connected to  $(n+1)!$  too:

- $n! \rightarrow (n+1)!$  if  $n$  is even
- $n! \rightarrow 2n! \rightarrow (n+1)!$  if  $n$  is odd.

This concludes the problem.

## §6 USAMO 2002/6

I have an  $n \times n$  sheet of stamps, from which I've been asked to tear out blocks of three adjacent stamps in a single row or column. (I can only tear along the perforations separating adjacent stamps, and each block must come out of the sheet in one piece.) Let  $b(n)$  be the smallest number of blocks I can tear out and make it impossible to tear out any more blocks. Prove that there are real constants  $c$  and  $d$  such that

$$\frac{1}{7}n^2 - cn \leq b(n) \leq \frac{1}{5}n^2 + dn$$

for all  $n > 0$ .

For the lower bound: there are  $2n(n - 2)$  places one could put a block. Note that each block eliminates at most 14 such places.

For the upper bound, the construction of  $\frac{1}{5}$  is easy to build. Here is an illustration of one possible construction for  $n = 9$  which generalizes readily, using only vertical blocks.

$$\begin{bmatrix} A & & E & & I & L & & P \\ A & & E & G & & L & & P & R \\ A & C & & G & & L & N & & R \\ & C & & G & J & & N & & R \\ & C & F & & J & & N & Q & \\ B & & F & & J & M & & Q & \\ B & & F & H & & M & & Q & S \\ B & D & & H & & M & O & & S \\ & D & & H & K & & O & & S \end{bmatrix}$$

Actually, for the lower bound, one may improve  $1/7$  to  $1/6$ . Count the number  $A$  of pairs of adjacent squares one of which is torn out and the other which is not:

- For every deleted block, there are eight neighboring squares, at least two on each long edge which have been deleted too. Hence  $N \leq 6b(n)$ .
- For every block still alive and not on the border, there are four neighboring squares, and clearly at least two are deleted. Hence  $N \geq 2((n - 2)^2 - 3b(n))$ .

Collating these solves the problem.