# **IMO 1998 Solution Notes**

EVAN CHEN《陳誼廷》

15 April 2024

This is a compilation of solutions for the 1998 IMO. The ideas of the solution are a mix of my own work, the solutions provided by the competition organizers, and solutions found by the community. However, all the writing is maintained by me.

These notes will tend to be a bit more advanced and terse than the "official" solutions from the organizers. In particular, if a theorem or technique is not known to beginners but is still considered "standard", then I often prefer to use this theory anyways, rather than try to work around or conceal it. For example, in geometry problems I typically use directed angles without further comment, rather than awkwardly work around configuration issues. Similarly, sentences like "let  $\mathbb{R}$  denote the set of real numbers" are typically omitted entirely.

Corrections and comments are welcome!

# Contents

0	Problems	2
1	Solutions to Day 1	3
	1.1 IMO 1998/1	3
	1.2 IMO 1998/2	5
	1.3 IMO 1998/3	6
2	Solutions to Day 2	7
	2.1 IMO 1998/4	7
	2.2 IMO 1998/5	8
	2.3 IMO 1998/6	9

## §0 Problems

- 1. A convex quadrilateral ABCD has perpendicular diagonals. The perpendicular bisectors of the sides AB and CD meet at a unique point P inside ABCD. Prove that the quadrilateral ABCD is cyclic if and only if triangles ABP and CDP have equal areas.
- 2. In a competition, there are a contestants and b judges, where  $b \ge 3$  is an odd integer. Each judge rates each contestant as either "pass" or "fail". Suppose k is a number such that for any two judges, their ratings coincide for at most k contestants. Prove that

$$\frac{k}{a} \ge \frac{b-1}{2b}$$

**3.** For any positive integer n, let  $\tau(n)$  denote the number of its positive divisors (including 1 and itself). Determine all positive integers m for which there exists a positive integer n such that

$$\frac{\tau(n^2)}{\tau(n)} = m.$$

- 4. Determine all pairs (x, y) of positive integers such that  $x^2y + x + y$  is divisible by  $xy^2 + y + 7$ .
- 5. Let I be the incenter of triangle ABC. Let the incircle of ABC touch the sides BC, CA, and AB at K, L, and M, respectively. The line through B parallel to MK meets the lines LM and LK at R and S, respectively. Prove that angle RIS is acute.
- **6.** Classify all functions  $f \colon \mathbb{N} \to \mathbb{N}$  satisfying the identity

$$f(n^2 f(m)) = m f(n)^2.$$

# §1 Solutions to Day 1

# §1.1 IMO 1998/1

Available online at https://aops.com/community/p124387.

#### **Problem statement**

A convex quadrilateral ABCD has perpendicular diagonals. The perpendicular bisectors of the sides AB and CD meet at a unique point P inside ABCD. Prove that the quadrilateral ABCD is cyclic if and only if triangles ABP and CDP have equal areas.

If ABCD is cyclic, then P is the circumcenter, and  $\angle APB + \angle PCD = 180^{\circ}$ . The hard part is the converse.



Let M and N be the midpoints of  $\overline{AB}$  and  $\overline{CD}$ .

**Claim** — Unconditionally, we have  $\angle NEM = \angle MPN$ .

*Proof.* Note that  $\overline{EN}$  is the median of right triangle  $\triangle ECD$ , and similarly for  $\overline{EM}$ . Hence  $\angle NED = \angle EDN = \angle BDC$ , while  $\angle AEM = \angle ACB$ . Since  $\angle DEA = 90^{\circ}$ , by looking at quadrilateral XDEA where  $X = \overline{CD} \cap \overline{AB}$ , we derive that  $\angle NED + \angle AEM + \angle DXA = 90^{\circ}$ , so

$$\measuredangle NEM = \measuredangle NED + \measuredangle AEM + 90^{\circ} = -\measuredangle DXA = -\measuredangle NXM = -\measuredangle NPM$$

as needed.

However, the area condition in the problem tells us

$$\frac{EN}{EM} = \frac{CN}{CM} = \frac{PM}{PN}.$$

Finally, we have  $\angle MEN > 90^{\circ}$  from the configuration. These properties uniquely determine the point E: it is the reflection of P across the midpoint of MN.

So EMPN is a parallelogram, and thus  $\overline{ME} \perp \overline{CD}$ . This implies  $\measuredangle BAE = \measuredangle CEM = \measuredangle EDC$  giving ABCD cyclic.

## §1.2 IMO 1998/2

Available online at https://aops.com/community/p124458.

#### **Problem statement**

In a competition, there are a contestants and b judges, where  $b \ge 3$  is an odd integer. Each judge rates each contestant as either "pass" or "fail". Suppose k is a number such that for any two judges, their ratings coincide for at most k contestants. Prove that

$$\frac{k}{a} \ge \frac{b-1}{2b}.$$

This is a "routine" problem with global ideas. We count pairs of coinciding ratings, i.e. the number N of tuples

$$(\{J_1, J_2\}, C)$$

of two distinct judges and a contestant for which the judges gave the same rating.

On the one hand, if we count by the judges, we have

$$N \le \binom{b}{2}k$$

by he problem condition.

On the other hand, if b = 2m + 1, then each contestant C contributes at least  $\binom{m}{2} + \binom{m+1}{2} = m^2$  to N, and so

$$N \geq a \cdot \left(\frac{b-1}{2}\right)^2$$

Putting together the two estimates for N yields the conclusion.

## §1.3 IMO 1998/3

Available online at https://aops.com/community/p124439.

#### **Problem statement**

For any positive integer n, let  $\tau(n)$  denote the number of its positive divisors (including 1 and itself). Determine all positive integers m for which there exists a positive integer n such that

$$\frac{\tau(n^2)}{\tau(n)} = m$$

The answer is odd integers m only. If we write  $n = p_1^{e_1} \dots p_k^{e_k}$  we get

$$\prod \frac{2e_i+1}{e_i+1} = m.$$

It's clear now that m must be odd, since every fraction has odd numerator.

We now endeavor to construct odd numbers. The proof is by induction, in which we are curating sets of fractions of the form  $\frac{2e+1}{e+1}$  that multiply to a given target. The base cases are easy to verify by hand. Generally, assume  $p = 2^t k - 1$  is odd, where

k is odd. Then we can write

$$\frac{2^{2t}k - 2^{t}(k+1) + 1}{2^{2t-1}k - 2^{t-1}(k+1) + 1} \cdot \frac{2^{2t-1}k - 2^{t-1}(k+1) + 1}{2^{2t-2}k - 2^{t-2}(k+1) + 1} \cdots \frac{2^{t+1}k - 2(k+1) + 1}{2^{t}k - 2^{0}(k+1) + 1}$$

Note that  $2^{2t}k - 2^t(k+1) + 1 = (2^tk - 1)(2^t - 1)$ , and  $2^tk - k = k(2^t - 1)$ , so the above fraction simplifies to

$$\frac{2^t k - 1}{k}$$

meaning we just need to multiply by k, which we can do using induction hypothesis.

# §2 Solutions to Day 2

## §2.1 IMO 1998/4

Available online at https://aops.com/community/p124428.

#### **Problem statement**

Determine all pairs (x, y) of positive integers such that  $x^2y + x + y$  is divisible by  $xy^2 + y + 7$ .

The answer is  $(7k^2, 7k)$  for all  $k \ge 1$ , as well as (11, 1) and (49, 1).

We are given  $xy^2 + y + 7 \mid x^2y + x + y$ . Multiplying the right-hand side by y gives

 $xy^2 + y + 7 \mid x^2y^2 + xy + y^2$ 

Then subtracting x times the left-hand side gives

$$xy^2 + y + 7 \mid y^2 - 7x.$$

We consider cases based on the sign of  $y^2 = 7x$ .

- If  $y^2 > 7x$ , then  $0 < y^2 7x < xy^2 + y + 7$ , contradiction.
- If  $y^2 = 7x$ , let y = 7k, so  $x = 7k^2$ . Plugging this back in to the original equation reads

 $343k^4 + 7k + 7 \mid 343k^5 + 7k^2 + 7k$ 

which is always valid, hence these are all solutions.

• If  $y^2 < 7x$ , then  $|y^2 - 7x| \le 7x$ , so  $y \in \{1, 2\}$ . When y = 1 we get

$$x+8 \mid x^2+x+1 \iff x+8 \mid 64-8+1=57.$$

This has solutions x = 11 and x = 49. When y = 2

$$4x + 9 \mid 2x^{2} + x + 2$$
  

$$\implies 4x + 9 \mid 16x^{2} + 8x + 16$$
  

$$\implies 4x + 9 \mid 81 - 18 + 16 = 79$$

which never occurs.

## §2.2 IMO 1998/5

Available online at https://aops.com/community/p121417.

#### **Problem statement**

Let I be the incenter of triangle ABC. Let the incircle of ABC touch the sides BC, CA, and AB at K, L, and M, respectively. The line through B parallel to MK meets the lines LM and LK at R and S, respectively. Prove that angle RIS is acute.

Observe that  $\triangle MKL$  is acute with circumcenter I. We now present two proofs.

¶ First simple proof (grobber). The problem is equivalent to showing  $BI^2 > BR \cdot BS$ . But from

$$\triangle BRK \sim \triangle MKL \sim \triangle BLS$$

we conclude

$$BR = t \cdot \frac{MK}{ML}, \qquad BS = t \cdot \frac{ML}{MK}$$

where t = BK = BL is the length of the tangent from B. Hence  $BR \cdot BS = t^2$ . Since BI > t is clear, we are done.

¶ Second projective proof. Let N be the midpoint of  $\overline{KL}$ , and let ray MN meet the incircle again at P.

Note that line  $\overline{RBS}$  is the polar of N. By Brokard's theorem, lines MK and PL should thus meet the polar of N, so we conclude  $R = \overline{MK} \cap \overline{PL}$ . Analogously,  $S = \overline{ML} \cap \overline{PK}$ .

Again by Brokard's theorem,  $\triangle NRS$  is self-polar, so N is the orthocenter of  $\triangle RIS$ . Since N lies between I and B we are done.

## §2.3 IMO 1998/6

Available online at https://aops.com/community/p124426.

#### **Problem statement**

Classify all functions  $f: \mathbb{N} \to \mathbb{N}$  satisfying the identity

$$f(n^2 f(m)) = m f(n)^2.$$

Let  $\mathcal{P}$  be the set of primes, and let  $g: \mathcal{P} \to \mathcal{P}$  be any involution on them. Extend g to a completely multiplicative function on  $\mathbb{N}$ . Then f(n) = dg(n) is a solution for any  $d \in \mathbb{N}$  which is fixed by g.

It's straightforward to check these all work, since  $g \colon \mathbb{N} \to \mathbb{N}$  is an involution on them. So we prove these are the only functions.

Let d = f(1).

**Claim** — We have 
$$df(n) = f(dn)$$
 and  $d \cdot f(ab) = f(a)f(b)$ .

*Proof.* Let P(m, n) denote the assertion in the problem statement. Off the bat,

- P(1,1) implies  $f(d) = d^2$ .
- P(n, 1) implies  $f(f(n)) = d^2n$ . In particular, f is injective.
- P(1, n) implies  $f(dn^2) = f(n)^2$ .

Then

$$f(a)^{2}f(b)^{2} = f(da^{2})f(b)^{2}$$
by third bullet  
$$= f(b^{2}f(f(da^{2})))$$
by problem statement  
$$= f(b^{2} \cdot d^{2} \cdot da^{2})$$
by second bullet  
$$= f(dab)^{2}$$
by third bullet  
$$\implies f(a)f(b) = f(dab).$$

This implies the first claim by taking (a,b) = (1,n). Then df(a) = f(da), and so we actually have f(a)f(b) = df(ab).

**Claim** — All values of f are divisible by d.

*Proof.* We have

$$f(n^2) = \frac{1}{d}f(n)^2$$
  
$$f(n^3) = \frac{f(n^2)f(n)}{d} = \frac{f(n)^3}{d^2}$$
  
$$f(n^4) = \frac{f(n^3)f(n)}{d} = \frac{f(n)^4}{d^3}$$

and so on, which implies the result.

Then, define g(n) = f(n)/d. We conclude that g is completely multiplicative, with g(1) = 1. However,  $f(f(n)) = d^2n$  also implies g(g(n)) = n, i.e. g is an involution. Moreover, since  $f(d) = d^2$ , g(d) = d.

All that remains is to check that g must map primes to primes to finish the description in the problem. This is immediate; since g is multiplicative and g(1) = 1, if g(g(p)) = pthen g(p) can have at most one prime factor, hence g(p) is itself prime.

**Remark.** The IMO problem actually asked for the least value of f(1998). But for instruction purposes, it is probably better to just find all f. Since  $1998 = 2 \cdot 3^3 \cdot 37$ , this answer is  $2^3 \cdot 3 \cdot 5 = 120$ , anyways.