

# IMO 2000 Solution Notes

COMPILED BY EVAN CHEN

January 22, 2020

This is an compilation of solutions for the 2000 IMO. Some of the solutions are my own work, but many are from the official solutions provided by the organizers (for which they hold any copyrights), and others were found on the Art of Problem Solving forums.

Corrections and comments are welcome!

## Contents

<a href="#">0 Problems</a>	2
<a href="#">1 IMO 2000/1</a>	3
<a href="#">2 IMO 2000/2</a>	4
<a href="#">3 IMO 2000/3</a>	5
<a href="#">4 IMO 2000/4</a>	7
<a href="#">5 IMO 2000/5</a>	9
<a href="#">6 IMO 2000/6</a>	10

## §0 Problems

1. Two circles  $G_1$  and  $G_2$  intersect at two points  $M$  and  $N$ . Let  $AB$  be the line tangent to these circles at  $A$  and  $B$ , respectively, so that  $M$  lies closer to  $AB$  than  $N$ . Let  $CD$  be the line parallel to  $AB$  and passing through the point  $M$ , with  $C$  on  $G_1$  and  $D$  on  $G_2$ . Lines  $AC$  and  $BD$  meet at  $E$ ; lines  $AN$  and  $CD$  meet at  $P$ ; lines  $BN$  and  $CD$  meet at  $Q$ . Show that  $EP = EQ$ .

2. Let  $a, b, c$  be positive real numbers with  $abc = 1$ . Show that

$$\left(a - 1 + \frac{1}{b}\right) \left(b - 1 + \frac{1}{c}\right) \left(c - 1 + \frac{1}{a}\right) \leq 1.$$

3. Let  $n \geq 2$  be a positive integer and  $\lambda$  a positive real number. Initially there are  $n$  fleas on a horizontal line, not all at the same point. We define a move as choosing two fleas at some points  $A$  and  $B$ , with  $A$  to the left of  $B$ , and letting the flea from  $A$  jump over the flea from  $B$  to the point  $C$  so that  $\frac{BC}{AB} = \lambda$ .

Determine all values of  $\lambda$  such that, for any point  $M$  on the line and for any initial position of the  $n$  fleas, there exists a sequence of moves that will take them all to the position right of  $M$ .

4. A magician has one hundred cards numbered 1 to 100. He puts them into three boxes, a red one, a white one and a blue one, so that each box contains at least one card. A member of the audience draws two cards from two different boxes and announces the sum of numbers on those cards. Given this information, the magician locates the box from which no card has been drawn.

How many ways are there to put the cards in the three boxes so that the trick works?


5. Does there exist a positive integer  $n$  such that  $n$  has exactly 2000 distinct prime divisors and  $n$  divides  $2^n + 1$ ?

6. Let  $\overline{AH_1}$ ,  $\overline{BH_2}$ , and  $\overline{CH_3}$  be the altitudes of an acute triangle  $ABC$ . The incircle  $\omega$  of triangle  $ABC$  touches the sides  $BC$ ,  $CA$  and  $AB$  at  $T_1$ ,  $T_2$  and  $T_3$ , respectively. Consider the symmetric images of the lines  $H_1H_2$ ,  $H_2H_3$ , and  $H_3H_1$  with respect to the lines  $T_1T_2$ ,  $T_2T_3$ , and  $T_3T_1$ . Prove that these images form a triangle whose vertices lie on  $\omega$ .

### §1 IMO 2000/1

Two circles  $G_1$  and  $G_2$  intersect at two points  $M$  and  $N$ . Let  $AB$  be the line tangent to these circles at  $A$  and  $B$ , respectively, so that  $M$  lies closer to  $AB$  than  $N$ . Let  $CD$  be the line parallel to  $AB$  and passing through the point  $M$ , with  $C$  on  $G_1$  and  $D$  on  $G_2$ . Lines  $AC$  and  $BD$  meet at  $E$ ; lines  $AN$  and  $CD$  meet at  $P$ ; lines  $BN$  and  $CD$  meet at  $Q$ . Show that  $EP = EQ$ .

First, we have  $\angle EAB = \angle ACM = \angle BAM$  and similarly  $\angle EBA = \angle BDM = \angle ABM$ . Consequently,  $\overline{AB}$  bisects  $\angle EAM$  and  $\angle EBM$ , and hence  $\triangle EAB \cong \triangle MAB$ .



Now it is well-known that  $\overline{MN}$  bisects  $\overline{AB}$  and since  $\overline{AB} \parallel \overline{PQ}$  we deduce that  $M$  is the midpoint of  $\overline{PQ}$ . As  $\overline{AB}$  is the perpendicular bisector of  $\overline{EM}$ , it follows that  $EP = EQ$  as well.

**§2 IMO 2000/2**

Let  $a, b, c$  be positive real numbers with  $abc = 1$ . Show that

$$\left(a - 1 + \frac{1}{b}\right) \left(b - 1 + \frac{1}{c}\right) \left(c - 1 + \frac{1}{a}\right) \leq 1.$$

---

Let  $a = x/y, b = y/z, c = z/x$  for  $x, y, z > 0$ . Then the inequality rewrites as

$$(-x + y + z)(x - y + z)(x + y - z) \leq xyz$$

which when expanded is equivalent to Schur's inequality. Alternatively, if one wants to avoid appealing to Schur, then the following argument works:

- At most one term on the left-hand side is negative; if that occurs we are done from  $xyz > 0 > (-x + y + z)(x - y + z)(x + y - z)$ .
- If all terms in the left-hand side are nonnegative, let us denote  $m = -x + y + z \geq 0$ ,  $n = x - y + z \geq 0$ ,  $p = x + y - z \geq 0$ . Then it becomes

$$mnp \leq \frac{(m+n)(n+p)(p+m)}{8}$$

which follows by AM-GM.

### §3 IMO 2000/3

Let  $n \geq 2$  be a positive integer and  $\lambda$  a positive real number. Initially there are  $n$  fleas on a horizontal line, not all at the same point. We define a move as choosing two fleas at some points  $A$  and  $B$ , with  $A$  to the left of  $B$ , and letting the flea from  $A$  jump over the flea from  $B$  to the point  $C$  so that  $\frac{BC}{AB} = \lambda$ .

Determine all values of  $\lambda$  such that, for any point  $M$  on the line and for any initial position of the  $n$  fleas, there exists a sequence of moves that will take them all to the position right of  $M$ .

The answer is  $\lambda \geq \frac{1}{n-1}$ .

We change the problem by replacing the fleas by **bowling balls**  $B_1, B_2, \dots, B_n$  in that order. Bowling balls aren't exactly great at jumping, so each move can now be described as follows:

- Select two indices  $i < j$ . Then ball  $B_i$  moves to  $B_{i+1}$ 's location,  $B_{i+1}$  moves to  $B_{i+2}$ 's location, and so on; until  $B_{j-1}$  moves to  $B_j$ 's location,
- Finally,  $B_j$  moves some distance forward; the distance is at most  $\lambda \cdot |B_j B_i|$  and  $B_j$  may not pass  $B_{j+1}$ .

**Remark.** Equivalently, you can phrase the proof without bowling balls as follows: if  $a_1 < \dots < a_n$  are the positions of the fleas, the quantity

$$L = a_n - \lambda(a_1 + \dots + a_{n-1})$$

is a monovariant which never increases; i.e.  $L$  is bounded above. Since  $L < (1 - (n-1)\lambda)a_n$ , it follows  $\lambda < \frac{1}{n-1}$  is necessary.

**Claim —** If  $\lambda < \frac{1}{n-1}$  the bowling balls have bounded movement.

*Proof.* Let  $a_i \geq 0$  denote the initial distance between  $B_i$  and  $B_{i+1}$ , and let  $\Delta_i$  denote the distance travelled by ball  $i$ . Of course we have  $\Delta_1 \leq a_1 + \Delta_2$ ,  $\Delta_2 \leq a_2 + \Delta_3$ ,  $\dots$ ,  $\Delta_{n-1} \leq a_{n-1} + \Delta_n$  by the relative ordering of the bowling balls. Finally, distance covered by  $B_n$  is always  $\lambda$  times distance travelled by other bowling balls, so

$$\begin{aligned} \Delta_n &\leq \lambda \sum_{i=1}^{n-1} \Delta_i \leq \lambda \sum_{i=1}^{n-1} ((a_i + a_{i+1} + \dots + a_{n-1}) + \Delta_n) \\ &= (n-1)\lambda \cdot \Delta_n + \sum_{i=1}^{n-1} \lambda a_i \end{aligned}$$

and since  $(n-1)\lambda > 1$ , this gives an upper bound. □

**Claim —** When  $\lambda \geq \frac{1}{n-1}$ , it suffices to always jump the leftmost flea over the rightmost flea.

*Proof.* If we let  $x_i$  denote the distance travelled by  $B_1$  in the  $i$ th step, then  $x_i = a_i$  for  $1 \leq i \leq n-1$  and  $x_i = \lambda(x_{i-1} + x_{i-2} + \dots + x_{i-(n-1)})$ .

In particular, if  $\lambda \geq \frac{1}{n-1}$  then each  $x_i$  is at least the average of the previous  $n-1$  terms. So if the  $a_i$  are not all zero, then  $\{x_n, \dots, x_{2n-2}\}$  are all positive and thereafter

$x_i \geq \min\{x_n, \dots, x_{2n-2}\} > 0$  for every  $i \geq 2n - 1$ . So the partial sums of  $x_i$  are unbounded, as desired.  $\square$

**Remark.** Other inductive constructions are possible. Here is the idea of one of them, although the details are more complicated.

We claim in general that given  $n - 1$  fleas at 0 and one flea at 1, we can get all the fleas arbitrarily close to  $\frac{1}{1-(n-1)\lambda}$  (or as far as we want if  $\lambda > \frac{1}{n-1}$ ). The proof is induction by  $n \geq 2$ ; for  $n = 2$  we get a geometric series. For  $n \geq 3$ , we leave one flea at zero and move the remainder close to  $\frac{1}{1-(n-2)\lambda}$ , then jump the last flea to  $\frac{1+\lambda}{1-(n-2)\lambda}$ .

Now we're in the same situation, except we shifted  $\frac{1}{1-(n-2)\lambda}$  right and have then scaled everything by  $r = \frac{\lambda}{1-(n-2)\lambda}$ . If we repeat this process again and check the geometric series, we see the fleas converge to

$$\frac{1}{1-(n-2)\lambda} (1 + r + r^2 + r^3 + \dots) = \frac{1}{1-(n-2)\lambda} \cdot \frac{1}{1-r} = \frac{1}{1-(n-1)\lambda}.$$

## §4 IMO 2000/4

A magician has one hundred cards numbered 1 to 100. He puts them into three boxes, a red one, a white one and a blue one, so that each box contains at least one card. A member of the audience draws two cards from two different boxes and announces the sum of numbers on those cards. Given this information, the magician locates the box from which no card has been drawn.

How many ways are there to put the cards in the three boxes so that the trick works?

There are  $2 \cdot 3! = 12$  ways, which amount to:

- Partitioning the cards modulo 3, or
- Placing 1 alone in a box, 100 alone in a second box, and all remaining cards in the third box.

These are easily checked to work so we prove they are the only ones.

Let  $A, B, C$  be the sets of cards in the three boxes. Then  $A + B, B + C, C + A$  should be disjoint, and contained in  $\{3, 4, \dots, 199\}$ . On the other hand, we have the following famous fact.

### Lemma

Let  $X$  and  $Y$  be finite nonempty sets of real numbers. We have  $|X+Y| \geq |X|+|Y|-1$ , with equality if and only if  $X$  and  $Y$  are arithmetic progressions with the same common difference, or one of  $X$  and  $Y$  is a singleton set.

Putting these two together gives the estimates

$$197 \geq |A + B| + |B + C| + |C + A| \geq 2(|A| + |B| + |C|) - 3 = 197.$$

So all the inequalities must be sharp. Consequently we conclude that:

**Claim** — Either the sets  $A, B, C$  are disjoint arithmetic progressions with the same common difference  $d = \min_{x \neq y \text{ in same set}} |x - y|$ , or two of the sets are two singleton. Moreover,  $\{3, 4, \dots, 199\} = (A + B) \sqcup (B + C) \sqcup (C + A)$ .

From here it is not hard to deduce the layouts above are the only ones, but there are some details. First, we make the preliminary observation that  $3 = 1 + 2$ ,  $4 = 1 + 3$ ,  $198 = 98 + 100$ ,  $199 = 99 + 100$  and these numbers can't be decomposed in other ways; thus from the remark about the disjoint union:

**Claim (Convenient corollary)** — The pairs  $(1, 2)$ ,  $(1, 3)$ ,  $(98, 100)$ ,  $(99, 100)$  are all in different sets.

We now consider the four cases.

- If two of the boxes are singletons, it follows from the corollary that we should have  $A = \{1\}$ ,  $B = \{100\}$  and  $C = \{2, \dots, 99\}$ , up to permutation.
- Otherwise  $A, B, C$  are disjoint arithmetic progressions with the same common difference  $d$ . As two of  $\{1, 2, 3, 4\}$  are in the same box (by pigeonhole), we must have  $d \leq 3$ .

- If  $d = 3$ , then no two elements of different residues modulo 3 can be in the same box, so we must be in the first construction claimed earlier.
- If  $d = 2$ , then the convenient corollary tells us we may assume WLOG that  $1 \in A$  and  $2 \in B$ , hence  $3 \in C$  (since  $3 \notin A$  by convenient corollary, and  $3 \notin B$  because common difference 2). Thus we must have  $A = \{1\}$ ,  $B = \{2, 4, \dots, 100\}$  and  $C = \{3, 5, \dots, 99\}$  which does not work since  $1 + 4 = 2 + 3$ . Therefore there are no solutions in this case.
- If  $d = 1$ , then by convenient corollary the numbers 1 and 2 are in different sets, as are 99 and 100. So we must have  $A = \{1\}$ ,  $B = \{2, \dots, 99\}$ ,  $C = \{100\}$  which we have already seen is valid.

**Remark.** A more direct solution is to replace 100 by  $n \geq 3$  and then proceed by induction on  $n$ . Indeed, any working configuration for  $\{1, \dots, n\}$  immediately gives a working configuration for  $n - 1$ .



## §5 IMO 2000/5

Does there exist a positive integer  $n$  such that  $n$  has exactly 2000 distinct prime divisors and  $n$  divides  $2^n + 1$ ?

---

Answer: Yes.

We say that  $n$  is *Korean* if  $n \mid 2^n + 1$ . First, observe that  $n = 9$  is Korean. Now, the problem is solved upon the following claim:

**Claim** — If  $n > 3$  is Korean, there exists a prime  $p$  not dividing  $n$  such that  $np$  is Korean too.

*Proof.* I claim that one can take any primitive prime divisor  $p$  of  $2^{2^n} - 1$ , which exists by Zsigmondy theorem. Obviously  $p \neq 2$ . Then:

- Since  $p \nmid 2^{\varphi(n)} - 1$  it follows then that  $p \nmid n$ .
- Moreover,  $p \mid 2^n + 1$  since  $p \nmid 2^n - 1$ .

Hence  $np \mid 2^n + 1 \mid 2^{np} + 1$  by Chinese Theorem, since  $\gcd(n, p) = 1$ . □

## §6 IMO 2000/6

Let  $\overline{AH_1}$ ,  $\overline{BH_2}$ , and  $\overline{CH_3}$  be the altitudes of an acute triangle  $ABC$ . The incircle  $\omega$  of triangle  $ABC$  touches the sides  $BC$ ,  $CA$  and  $AB$  at  $T_1$ ,  $T_2$  and  $T_3$ , respectively. Consider the symmetric images of the lines  $H_1H_2$ ,  $H_2H_3$ , and  $H_3H_1$  with respect to the lines  $T_1T_2$ ,  $T_2T_3$ , and  $T_3T_1$ . Prove that these images form a triangle whose vertices lie on  $\omega$ .

We use complex numbers with  $\omega$  the unit circle. Let  $T_1 = a$ ,  $T_2 = b$ ,  $T_3 = c$ . The main content of the problem is to show that the triangle in question has vertices  $ab/c$ ,  $bc/a$ ,  $ca/b$  (which is evident from a good diagram).


Since  $A = \frac{2bc}{b+c}$ , we have

$$H_1 = \frac{1}{2} \left( \frac{2bc}{b+c} + a + a - a^2 \cdot \frac{2}{b+c} \right) = \frac{ab + bc + ca - a^2}{b+c}.$$

The reflection of  $H_1$  over  $\overline{T_1T_2}$  is

$$\begin{aligned} H_1^C &= a + b - ab\overline{H_1} = a + b - b \cdot \frac{ac + ab + a^2 - bc}{a(b+c)} \\ &= \frac{a(a+b)(b+c) - b(a^2 + ab + ac - bc)}{a(b+c)} = \frac{c(a^2 + b^2)}{a(b+c)}. \end{aligned}$$

Now, we claim that  $H_1^C$  lies on the chord joining  $\frac{ca}{b}$  and  $\frac{cb}{a}$ ; by symmetry so will  $H_2^C$  and this will imply the problem (it means that the desired triangle has vertices  $ab/c$ ,  $bc/a$ ,  $ca/b$ ). A cartoon of this is shown below.



To see this, it suffices to compute

$$\begin{aligned} H_1^C + \left(\frac{ca}{b}\right) \left(\frac{cb}{a}\right) \overline{H_1^C} &= \frac{c(a^2 + b^2)}{a(b+c)} + c^2 \frac{\frac{1}{c} \cdot \frac{a^2+b^2}{a^2b^2}}{\frac{1}{a} \left(\frac{b+c}{bc}\right)} \\ &= \frac{c(a^2 + b^2)}{a(b+c)} + \frac{c(a^2 + b^2)}{abc^{-1}(b+c)} \\ &= \frac{c(a^2 + b^2)}{a(b+c)} \left(\frac{b+c}{b}\right) \\ &= \frac{c(a^2 + b^2)}{ab} = \frac{ca}{b} + \frac{cb}{a} \end{aligned}$$

as desired.