

IMO 2006 Solution Notes

Evan Chen 《陳誼廷》

23 July 2025

This is a compilation of solutions for the 2006 IMO. The ideas of the solution are a mix of my own work, the solutions provided by the competition organizers, and solutions found by the community. However, all the writing is maintained by me.

These notes will tend to be a bit more advanced and terse than the “official” solutions from the organizers. In particular, if a theorem or technique is not known to beginners but is still considered “standard”, then I often prefer to use this theory anyways, rather than try to work around or conceal it. For example, in geometry problems I typically use directed angles without further comment, rather than awkwardly work around configuration issues. Similarly, sentences like “let \mathbb{R} denote the set of real numbers” are typically omitted entirely.

Corrections and comments are welcome!

Contents

0 Problems	2
1 Solutions to Day 1	3
1.1 IMO 2006/1, proposed by Hojoo Lee (KOR)	3
1.2 IMO 2006/2, proposed by Dušan Djukić (SRB)	4
1.3 IMO 2006/3, proposed by Finbarr Holland (IRL)	5
2 Solutions to Day 2	6
2.1 IMO 2006/4, proposed by Zuming Feng (USA)	6
2.2 IMO 2006/5, proposed by Dan Schwarz (ROU)	7
2.3 IMO 2006/6, proposed by Dušan Djukić (SRB)	8

§0 Problems

1. Let ABC be a triangle with incenter I . A point P in the interior of the triangle satisfies

$$\angle PBA + \angle PCA = \angle PBC + \angle PCB.$$

Show that $AP \geq AI$ and that equality holds if and only if $P = I$.

2. Let P be a regular 2006-gon. A diagonal is called *good* if its endpoints divide the boundary of P into two parts, each composed of an odd number of sides of P . The sides of P are also called *good*. Suppose P has been dissected into triangles by 2003 diagonals, no two of which have a common point in the interior of P . Find the maximum number of isosceles triangles having two good sides that could appear in such a configuration.

3. Determine the least real number M such that the inequality

$$|ab(a^2 - b^2) + bc(b^2 - c^2) + ca(c^2 - a^2)| \leq M(a^2 + b^2 + c^2)^2$$

holds for all real numbers a , b and c .

4. Determine all pairs (x, y) of integers such that

$$1 + 2^x + 2^{2x+1} = y^2.$$

5. Let $P(x)$ be a polynomial of degree $n > 1$ with integer coefficients and let k be a positive integer. Consider the polynomial

$$Q(x) = P(P(\dots P(P(x)) \dots))$$

where P occurs k times. Prove that there are at most n integers t such that $Q(t) = t$.

6. Assign to each side b of a convex polygon P the maximum area of a triangle that has b as a side and is contained in P . Show that the sum of the areas assigned to the sides of P is at least twice the area of P .

§1 Solutions to Day 1

§1.1 IMO 2006/1, proposed by Hojoo Lee (KOR)

Available online at <https://aops.com/community/p571966>.

Problem statement

Let ABC be a triangle with incenter I . A point P in the interior of the triangle satisfies

$$\angle PBA + \angle PCA = \angle PBC + \angle PCB.$$

Show that $AP \geq AI$ and that equality holds if and only if $P = I$.

The condition rewrites as

$$\angle PBC + \angle PCB = (\angle B - \angle PBC) + (\angle C - \angle PCB) \implies \angle PBC + \angle PCB = \frac{\angle B + \angle C}{2}$$

which means that

$$\angle BPC = 180^\circ - \frac{\angle B + \angle C}{2} = 90^\circ + \frac{\angle A}{2} = \angle BIC.$$

Since P and I are both inside $\triangle ABC$ that implies P lies on the circumcircle of $\triangle BIC$.

It's well-known (by "Fact 5") that the circumcenter of $\triangle BIC$ is the arc midpoint M of \widehat{BC} . Therefore

$$AI + IM = AM \leq AP + PM \implies AI \leq AP$$

with equality holding iff A, P, M are collinear, or $P = I$.

§1.2 IMO 2006/2, proposed by Dušan Djukić (SRB)

Available online at <https://aops.com/community/p571973>.

Problem statement

Let P be a regular 2006-gon. A diagonal is called *good* if its endpoints divide the boundary of P into two parts, each composed of an odd number of sides of P . The sides of P are also called *good*. Suppose P has been dissected into triangles by 2003 diagonals, no two of which have a common point in the interior of P . Find the maximum number of isosceles triangles having two good sides that could appear in such a configuration.

Call a triangle with the desired property *special*. We prove the maximum number of special triangles is 1003, achieved by paring up the sides of the polygon.

We present two solutions for the upper bound. Both of them rely first on two geometric notes:

- In a special triangle, the good sides are congruent (and not congruent to the third side).
- No two isosceles triangles share a good side.

¶ **Solution using bijections.** Call a good diagonal **special** if it's part of a special triangle; special diagonals come in pairs. Consider the minor arc cut out by a special diagonal d , which has an odd number of sides. Since special diagonals come in pairs, one can associate to d a side of the polygon not covered by any special diagonals from d . Hence there are at most 2006 special diagonals, so at most 1003 special triangles.

¶ **Solution using graph theory.** Consider the tree T formed by the 2004 triangles in the dissection, with obvious adjacency. Let F be the forest obtained by deleting any edge corresponding to a good diagonal. Then the resulting graph F has only degrees 1 and 3, with special triangles only occurring at degree 1 vertices.

If there are k good diagonals drawn, then this forest consists of $k + 1$ trees. A tree with n_i vertices ($0 \leq i \leq k$) consequently has $\frac{n_i + 2}{2}$ leaves. However by the earlier remark at least k leaves don't give special triangles (one on each side of a special diagonal); so the number of leaves that do give good triangles is at most

$$-k + \sum_i \frac{n_i + 2}{2} = -k + \frac{2004 + 2(k + 1)}{2} = 1003.$$

§1.3 IMO 2006/3, proposed by Finbarr Holland (IRL)

Available online at <https://aops.com/community/p571945>.

Problem statement

Determine the least real number M such that the inequality

$$|ab(a^2 - b^2) + bc(b^2 - c^2) + ca(c^2 - a^2)| \leq M(a^2 + b^2 + c^2)^2$$

holds for all real numbers a, b and c .

It's the same as

$$|(a-b)(b-c)(c-a)(a+b+c)| \leq M(a^2 + b^2 + c^2)^2.$$

Let $x = a - b$, $y = b - c$, $z = c - a$, $s = a + b + c$. Then we want to have

$$|xyzs| \leq \frac{M}{9}(x^2 + y^2 + z^2 + s^2)^2.$$

Here $x + y + z = 0$.

Now if x and y have the same sign, we can replace them with the average (this increases the LHS and decreases RHS). So we can have $x = y$, $z = -2x$. Now WLOG $x > 0$ to get

$$2x^3 \cdot s \leq \frac{M}{9}(6x^2 + s^2)^2.$$

After this routine calculation gives $M = \frac{9}{32}\sqrt{2}$ works and is optimal (by $6x^2 + s^2 = 2x^2 + 2x^2 + 2x^2 + s^2$ and AM-GM).

§2 Solutions to Day 2

§2.1 IMO 2006/4, proposed by Zuming Feng (USA)

Available online at <https://aops.com/community/p572815>.

Problem statement

Determine all pairs (x, y) of integers such that

$$1 + 2^x + 2^{2x+1} = y^2.$$

Answers: $(0, \pm 2)$, $(4, \pm 23)$, which work.

Assume $x \geq 4$.

$$2^x (1 + 2^{x+1}) = 2^x + 2^{2x+1} = y^2 - 1 = (y-1)(y+1).$$

So either:

- $y = 2^{x-1}m + 1$ for some odd m , so

$$1 + 2^{x+1} = m(2^{x-2}m + 1) \implies 2^x = \frac{4(1-m)}{m^2-8}.$$

- $y = 2^{x-1}m - 1$ for some odd m , so

$$1 + 2^{x+1} = m(2^{x-2}m - 1) \implies 2^x = \frac{4(1+m)}{m^2-8}.$$

In particular we need $4|1 \pm m| \geq 2^4|m^2 - 8|$, which is enough to imply $m < 5$. From here easily recover $x = 4$, $m = 3$ as the last solution (in the second case).

§2.2 IMO 2006/5, proposed by Dan Schwarz (ROU)

Available online at <https://aops.com/community/p572821>.

Problem statement

Let $P(x)$ be a polynomial of degree $n > 1$ with integer coefficients and let k be a positive integer. Consider the polynomial

$$Q(x) = P(P(\dots P(P(x)) \dots))$$

where P occurs k times. Prove that there are at most n integers t such that $Q(t) = t$.

First, we prove that:

Claim (Putnam 2000 et al) — If a number is periodic under P then in fact it's fixed by $P \circ P$.

Proof. Let x_1, x_2, \dots, x_n be a minimal orbit. Then

$$x_i - x_{i+1} \mid P(x_i) - P(x_{i+1}) = x_{i+1} - x_{i+2}$$

and so on cyclically.

If any of the quantities are zero we are done. Else, we must eventually have $x_i - x_{i+1} = -(x_{i+1} - x_{i+2})$, so $x_i = x_{i+2}$ and we get 2-periodicity. \square

The tricky part is to study the 2-orbits. Suppose there exists a fixed pair $u \neq v$ with $P(u) = v$, $P(v) = u$. (If no such pair exists, we are already done.) Let (a, b) be any other pair with $P(a) = b$, $P(b) = a$, possibly even $a = b$, but $\{a, b\} \cap \{u, v\} = \emptyset$. Then we should have

$$u - a \mid P(u) - P(a) = v - b \mid P(v) - P(b) = u - a$$

and so $u - a$ and $v - b$ divide each other (and are nonzero). Similarly, $u - b$ and $v - a$ divide each other.

Hence $u - a = \pm(v - b)$ and $u - b = \pm(v - a)$. We consider all four cases:

- If $u - a = v - b$ and $u - b = v - a$ then $u - v = b - a = a - b$, contradiction.
- If $u - a = -(v - b)$ and $u - b = -(v - a)$ then $u + v = u - v = a + b$.
- If $u - a = -(v - b)$ and $u - b = v - a$, we get $a + b = u + v$ from the first one (discarding the second).
- If $u - a = v - b$ and $u - b = -(v - a)$, we get $a + b = u + v$ from the second one (discarding the first one).

Thus in all possible situations we have

$$a + b = c := u + v$$

a fixed constant.

Therefore, any pair (a, b) with $P(a) = b$ and $P(b) = a$ actually satisfies $P(a) = c - a$. And since $\deg P > 1$, this means there are at most n roots to $a + P(a) = c$, as needed.

§2.3 IMO 2006/6, proposed by Dušan Djukić (SRB)

Available online at <https://aops.com/community/p572824>.

Problem statement

Assign to each side b of a convex polygon P the maximum area of a triangle that has b as a side and is contained in P . Show that the sum of the areas assigned to the sides of P is at least twice the area of P .

We say a polygon is *almost convex* if all its angles are at most 180° .


Note that given any convex or almost convex polygon, we can take any side b and add another vertex on it, and the sum of the labels doesn't change (since the label of a side is the length of the side times the distance of the farthest point).

Lemma

Let N be an even integer. Then any almost convex N -gon with area S should have an inscribed triangle with area at least $2S/N$.

The main work is the proof of the lemma.

Label the polygon $P_0P_1 \dots P_{N-1}$. Consider the $N/2$ major diagonals of the almost convex N -gon, $P_0P_{N/2}$, $P_1P_{N/2+1}$, et cetera. A *butterfly* refers to a self-intersecting quadrilateral $P_iP_{i+1}P_{i+1+N/2}P_{i+N/2}$. An example of a butterfly is shown below for $N = 8$.



Claim — Every point X in the polygon is contained in the wingspan of some butterfly.

Proof. Consider a windmill-like process which

- starts from some oriented red line $P_0P_{N/2}$, oriented to face $P_0P_{N/2}$

- rotates through $P_0P_{N/2} \cap P_1P_{N/2+1}$ to get line $P_1P_{N/2+1}$,
- rotates through $P_1P_{N/2+1} \cap P_2P_{N/2+2}$ to get line $P_2P_{N/2+2}$,
- ...et cetera, until returning to line $P_{N/2}P_0$, but in the reverse orientation.

At the end of the process, every point in the plane has switched sides with our moving line. The moment that X crosses the moving red line, we get it contained in a butterfly, as needed. \square

Claim — If $ABDC = P_iP_{i+1}P_{i+1+N/2}P_{i+N/2}$ is a butterfly, one of the triangles ABC , BCD , CDA , DAB has area at least that of the butterfly.

Proof. Let the diagonals of the butterfly meet at O , and let $a = AO$, $b = BO$, $c = CO$, $d = DO$. If we assume WLOG $d = \min(a, b, c, d)$ then it follows $[ABC] = [AOB] + [BOC] \geq [AOB] + [COD]$, as needed. \square

Now, since the $N/2$ butterflies cover an area of S , it follows that one of the butterflies has area at least $S/(N/2) = 2S/N$, and so that butterfly gives a triangle with area at least $2S/N$, completing the proof of the lemma.

¶ **Main proof.** Let a_1, \dots, a_n be the numbers assigned to the sides. Assume for contradiction $a_1 + \dots + a_n < 2S$. We pick even integers m_1, m_2, \dots, m_n such that

$$\begin{aligned} \frac{a_1}{S} &< \frac{2m_1}{m_1 + \dots + m_n} \\ \frac{a_2}{S} &< \frac{2m_2}{m_1 + \dots + m_n} \\ &\vdots \\ \frac{a_n}{S} &< \frac{2m_n}{m_1 + \dots + m_n}. \end{aligned}$$

which is possible by rational approximation, since the right-hand sides sum to 2 and the left-hand sides sum to strictly less than 2.

Now we break every side of P into m_i equal parts to get an almost convex N -gon, where $N = m_1 + \dots + m_n$.

The main lemma then gives us a triangle Δ of the almost convex N -gon which has area at least $\frac{2S}{N}$. If Δ used the i th side then it then follows the label a_i on that side should be at least $m_i \cdot \frac{2S}{N}$, contradiction.