

Team Selection Test for the 55th International Mathematical Olympiad

United States of America

Day I

Thursday, December 12, 2013

Time limit: 4.5 hours. Each problem is worth 7 points.

IMO TST 1. Let ABC be an acute triangle, and let X be a variable interior point on the minor arc BC of its circumcircle. Let P and Q be the feet of the perpendiculars from X to lines CA and CB , respectively. Let R be the intersection of line PQ and the perpendicular from B to AC . Let ℓ be the line through P parallel to XR . Prove that as X varies along minor arc BC , the line ℓ always passes through a fixed point.

IMO TST 2. Let a_1, a_2, a_3, \dots be a sequence of integers, with the property that every consecutive group of a_i 's averages to a perfect square. More precisely, for all positive integers n and k , the quantity

$$\frac{a_n + a_{n+1} + \cdots + a_{n+k-1}}{k}$$

is always the square of an integer. Prove that the sequence must be constant (all a_i are equal to the same perfect square).

IMO TST 3. Let n be an even positive integer, and let G be an n -vertex (simple) graph with exactly $\frac{n^2}{4}$ edges. An unordered pair of distinct vertices $\{x, y\}$ is said to be *amicable* if they have a common neighbor (there is a vertex z such that xz and yz are both edges). Prove that G has at least $2\binom{n/2}{2}$ pairs of vertices which are amicable.

Team Selection Test for the 55th International Mathematical Olympiad

United States of America

Day II

Thursday, January 23, 2014

Time limit: 4.5 hours. Each problem is worth 7 points.

IMO TST 4. Let n be a positive even integer, and let c_1, c_2, \dots, c_{n-1} be real numbers satisfying

$$\sum_{i=1}^{n-1} |c_i - 1| < 1.$$

Prove that

$$2x^n - c_{n-1}x^{n-1} + c_{n-2}x^{n-2} - \dots - c_1x^1 + 2$$

has no real roots.

IMO TST 5. Let $ABCD$ be a cyclic quadrilateral, and let $E, F, G,$ and H be the midpoints of $AB, BC, CD,$ and DA respectively. Let W, X, Y and Z be the orthocenters of triangles AHE, BEF, CFG and $DGH,$ respectively. Prove that the quadrilaterals $ABCD$ and $WXYZ$ have the same area.

IMO TST 6. For a prime $p,$ a subset S of residues modulo p is called a *sum-free multiplicative subgroup* of \mathbb{F}_p if

- there is a nonzero residue α modulo p such that $S = \{1, \alpha^1, \alpha^2, \dots\}$ (all considered mod p), and
- there are no $a, b, c \in S$ (not necessarily distinct) such that $a + b \equiv c \pmod{p}$.

Prove that for every integer $N,$ there is a prime p and a sum-free multiplicative subgroup S of \mathbb{F}_p such that $|S| \geq N.$