

IMO 2022 Solution Notes

COMPILED BY EVAN CHEN

27 January 2023

This is an compilation of solutions for the 2022 IMO. Some of the solutions are my own work, but many are from the official solutions provided by the organizers (for which they hold any copyrights), and others were found by users on the Art of Problem Solving forums.

These notes will tend to be a bit more advanced and terse than the “official” solutions from the organizers. In particular, if a theorem or technique is not known to beginners but is still considered “standard”, then I often prefer to use this theory anyways, rather than try to work around or conceal it. For example, in geometry problems I typically use directed angles without further comment, rather than awkwardly work around configuration issues. Similarly, sentences like “let \mathbb{R} denote the set of real numbers” are typically omitted entirely.

Corrections and comments are welcome!

Contents

0 Problems	2
1 IMO 2022/1, proposed by France	3
2 IMO 2022/2, proposed by Netherlands	4
3 IMO 2022/3, proposed by Ankan Bhattacharya (USA)	5
4 IMO 2022/4, proposed by Patrik Bak (SVK)	6
5 IMO 2022/5, proposed by Belgium	8
6 IMO 2022/6, proposed by Nikola Petrović (SRB)	9

§0 Problems

1. The Bank of Oslo issues two types of coin: aluminum (denoted A) and bronze (denoted B). Marianne has n aluminum coins and n bronze coins arranged in a row in some arbitrary initial order. A chain is any subsequence of consecutive coins of the same type. Given a fixed positive integer $k \leq 2n$, Gilberty repeatedly performs the following operation: he identifies the longest chain containing the k^{th} coin from the left and moves all coins in that chain to the left end of the row. For example, if $n = 4$ and $k = 4$, the process starting from the ordering $AABBBABA$ would be $AABBBABA \rightarrow BBBAAABA \rightarrow AAABBBBA \rightarrow BBBBAAAA \rightarrow \dots$.

Find all pairs (n, k) with $1 \leq k \leq 2n$ such that for every initial ordering, at some moment during the process, the leftmost n coins will all be of the same type.

2. Find all functions $f: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that for each $x \in \mathbb{R}^+$, there is exactly one $y \in \mathbb{R}^+$ satisfying

$$xf(y) + yf(x) \leq 2.$$

3. Let k be a positive integer and let S be a finite set of odd prime numbers. Prove that there is at most one way (up to rotation and reflection) to place the elements of S around the circle such that the product of any two neighbors is of the form $x^2 + x + k$ for some positive integer x .
4. Let $ABCDE$ be a convex pentagon such that $BC = DE$. Assume that there is a point T inside $ABCDE$ with $TB = TD, TC = TE$ and $\angle ABT = \angle TEA$. Let line AB intersect lines CD and CT at points P and Q , respectively. Assume that the points P, B, A, Q occur on their line in that order. Let line AE intersect CD and DT at points R and S , respectively. Assume that the points R, E, A, S occur on their line in that order. Prove that the points P, S, Q, R lie on a circle.
5. Find all triples (a, b, p) of positive integers with p prime and

$$a^p = b! + p.$$

6. Let n be a positive integer. A *Nordic square* is an $n \times n$ board containing all the integers from 1 to n^2 so that each cell contains exactly one number. An *uphill path* is a sequence of one or more cells such that:
- the first cell in the sequence is a *valley*, meaning the number written is less than all its orthogonal neighbors,
 - each subsequent cell in the sequence is orthogonally adjacent to the previous cell, and
 - the numbers written in the cells in the sequence are in increasing order.

Find, as a function of n , the smallest possible total number of uphill paths in a Nordic square.

§1 IMO 2022/1, proposed by France

The Bank of Oslo issues two types of coin: aluminum (denoted A) and bronze (denoted B). Marianne has n aluminum coins and n bronze coins arranged in a row in some arbitrary initial order. A chain is any subsequence of consecutive coins of the same type. Given a fixed positive integer $k \leq 2n$, Gilberty repeatedly performs the following operation: he identifies the longest chain containing the k^{th} coin from the left and moves all coins in that chain to the left end of the row. For example, if $n = 4$ and $k = 4$, the process starting from the ordering $AABBBABA$ would be $AABBBABA \rightarrow BBBA AABA \rightarrow AAABBBBA \rightarrow BBBBAAAA \rightarrow \dots$.

Find all pairs (n, k) with $1 \leq k \leq 2n$ such that for every initial ordering, at some moment during the process, the leftmost n coins will all be of the same type.

Answer: $n \leq k \leq \lceil \frac{3}{2}n \rceil$.

Call a maximal chain a *block*. Then the line can be described as a sequence of blocks: it's one of:

$$\begin{array}{c} \underbrace{A \dots A}_{e_1} \underbrace{B \dots B}_{e_2} \underbrace{A \dots A}_{e_3} \dots \underbrace{A \dots A}_{e_m} \text{ for odd } m \\ \underbrace{A \dots A}_{e_1} \underbrace{B \dots B}_{e_2} \underbrace{A \dots A}_{e_3} \dots \underbrace{B \dots B}_{e_m} \text{ for even } m \end{array}$$

or the same thing with the roles of A and B flipped.

The main claim is the following:

Claim — The number m of blocks will never increase after an operation. Moreover, it stays the same if and only if

- $k \leq e_1$; or
- m is even and $e_m \geq 2n + 1 - k$.

Proof. This is obvious, just run the operation and see! □

The problem asks for which values of k we always reach $m = 2$ eventually; we already know that it's non-increasing. We consider a few cases:

- If $k < n$, then any configuration with $e_1 = n - 1$ will never change.
- If $k > \lceil 3n/2 \rceil$, then take $m = 4$ and $e_1 = e_2 = \lfloor n/2 \rfloor$ and $e_3 = e_4 = \lceil n/2 \rceil$. This configuration retains $m = 4$ always: the blocks simply rotate.
- Conversely, suppose $k \geq n$ has the property that $m > 2$ stays fixed. If after the first three operations m hasn't changed, then we must have $m \geq 4$ even, and $e_m, e_{m-1}, e_{m-2} \geq 2n + 1 - k$. Now,

$$n \geq e_m + e_{m-2} \geq 2(2n + 1 - k) \implies k \geq \frac{3}{2}n + 1$$

so this completes the proof.

§2 IMO 2022/2, proposed by Netherlands

Find all functions $f: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that for each $x \in \mathbb{R}^+$, there is exactly one $y \in \mathbb{R}^+$ satisfying

$$xf(y) + yf(x) \leq 2.$$

The answer is $f(x) \equiv 1/x$ which obviously works (here $y = x$).

For the converse, assume we have f such that each $x \in \mathbb{R}^+$ has a *friend* y with $xf(y) + yf(x) \leq 2$. By symmetry y is also the friend of x .

Claim — In fact every number is its own friend.

Proof. Assume for contradiction $a \neq b$ are friends. Then we know that $af(a) + af(a) > 2 \implies f(a) > \frac{1}{a}$. Analogously, $f(b) > \frac{1}{b}$. However, we then get

$$2 \geq af(b) + bf(a) > \frac{a}{b} + \frac{b}{a} \stackrel{\text{AMGM}}{\geq} 2$$

which is impossible. □

The problem condition now simplifies to saying

$$f(x) \leq \frac{1}{x} \text{ for all } x, \quad xf(y) + yf(x) > 2 \text{ for all } x \neq y.$$

In particular, for any $x > 0$ and $\varepsilon > 0$ we have

$$\begin{aligned} 2 &< xf(x + \varepsilon) + (x + \varepsilon)f(x) \leq \frac{x}{x + \varepsilon} + (x + \varepsilon)f(x) \\ \implies f(x) &> \frac{x + 2\varepsilon}{(x + \varepsilon)^2} = \frac{1}{x + \frac{\varepsilon^2}{x + 2\varepsilon}}. \end{aligned}$$

Since this holds for all $\varepsilon > 0$ this forces $f(x) \geq \frac{1}{x}$ as well. We're done.

§3 IMO 2022/3, proposed by Ankan Bhattacharya (USA)

Let k be a positive integer and let S be a finite set of odd prime numbers. Prove that there is at most one way (up to rotation and reflection) to place the elements of S around the circle such that the product of any two neighbors is of the form $x^2 + x + k$ for some positive integer x .

We replace “positive integer x ” with “nonnegative integer x ”, and say numbers of the form $x^2 + x + k$ are *good*. We could also replace “nonnegative integer x ” with “integer x ” owing to the obvious map $x \mapsto 1 - x$.

Claim — If p is an odd prime, there are at most two odd primes q and r less than p for which $pq = x^2 + x + k$ and $pr = y^2 + y + k$ are good.

Moreover, if the above occurs and $x, y \geq 0$, then $x + y + 1 = p$ and $xy \equiv k \pmod{p}$.

Proof. The equation $T^2 + T + k \equiv 0 \pmod{p}$ has at most two solutions modulo p , i.e. at most two solutions in the interval $[0, p - 1]$. Because $0 \leq x, y < p$ from $p > \max(q, r)$ and $k > 0$, the first half follows.

For the second half, Vieta also gives $x + y \equiv 1 \pmod{p}$ and $xy \equiv k \pmod{p}$, and we know $0 < x + y < 2p$. \square

Claim — If two such primes do exist as above, then qr is also good (!).

Proof. Let $pq = x^2 + x + k$ and $pr = y^2 + y + k$ for $x, y \geq 0$ as before. Fix $\alpha \in \mathbb{C}$ such that $\alpha^2 + \alpha + k = 0$; then for any $n \in \mathbb{Z}$, we have

$$n^2 + n + k = \text{Norm}(n - \alpha).$$

Hence

$$pq \cdot pr = \text{Norm}\left((x - \alpha)(y - \alpha)\right) = \text{Norm}\left((xy - k) - (x + y + 1)\alpha\right)$$

But $\text{Norm}(p) = p^2$, so combining with the second half of the previous claim gives

$$qr = \text{Norm}\left(\frac{1}{p}(xy - k) - \alpha\right)$$

as needed. \square

These two claims imply the claim directly by induction on $|S|$, since one can now delete the largest element of S .

Remark. To show that the condition is not vacuous, the author gave a ring of 385 primes for $k = 41$; see <https://aops.com/community/p26068963>.


§4 IMO 2022/4, proposed by Patrik Bak (SVK)

Let $ABCDE$ be a convex pentagon such that $BC = DE$. Assume that there is a point T inside $ABCDE$ with $TB = TD, TC = TE$ and $\angle ABT = \angle TEA$. Let line AB intersect lines CD and CT at points P and Q , respectively. Assume that the points P, B, A, Q occur on their line in that order. Let line AE intersect CD and DT at points R and S , respectively. Assume that the points R, E, A, S occur on their line in that order. Prove that the points P, S, Q, R lie on a circle.

The conditions imply

$$\triangle BTC \cong \triangle DTE, \quad \text{and} \quad \triangle BTY \sim \triangle ETX.$$

Define $K = \overline{CT} \cap \overline{AE}$, $L = \overline{DT} \cap \overline{AB}$, $X = \overline{BT} \cap \overline{AE}$, $Y = \overline{ET} \cap \overline{BY}$.



Claim (Main claim) — We have

$$\triangle BTQ \sim \triangle ETS, \quad \text{and} \quad BY : YL : LQ = EX : XK : KS.$$

In other words, $TBYLQ \sim TEXKS$.

Proof. We know $\triangle BTY \sim \triangle ETX$. Also, $\angle BTL = \angle BTD = \angle CTE = \angle KTE$ and $\angle BTQ = \angle BTC = \angle DTE = \angle STE$. \square

It follows from the claim that:

- $TL/TQ = TK/TS$, ergo $TL \cdot TS = TK \cdot TQ$, so $KLSQ$ is cyclic; and
- $TC/TK = TE/TK = TB/TL = TD/TL$, so $\overline{KL} \parallel \overline{PCDR}$.

With these two bullets, we're done by Reim theorem.

§5 IMO 2022/5, proposed by Belgium

Find all triples (a, b, p) of positive integers with p prime and

$$a^p = b! + p.$$

The answer is $(2, 2, 2)$ and $(3, 4, 3)$ only, which work.
In what follows we assume $a \geq 2$.

Claim — We have $b \leq 2p - 2$, and hence $a < p^2$.

Proof. For the first half, assume first for contradiction that $b \geq 2p$. Then $b! + p \equiv p \pmod{p^2}$, so $\nu_p(b! + p) = 1$, but $\nu_p(a^p) = 1$ never occurs.

We can also rule out $b = 2p - 1$ since that would give

$$(2p - 1)! + p = p[(p - 1)!(p + 1)(p + 2) \dots (2p - 1) + 1]$$

By Wilson theorem the inner bracket is $(-1)^2 + 1 \equiv 2 \pmod{p}$ exactly, contradiction for $p > 2$. And when $p = 2$, $3! + 2 = 8$ is not a perfect square.

The second half follows as $a^p \leq (2p - 2)! + p < p^{2p}$. (Here we used the crude estimate $(2p - 2)! = \prod_{k=1}^{p-1} k \cdot (2p - 1 - k) < (p(p - 1))^{p-1}$.) \square

Claim — We have $a \geq p$, and hence $b \geq p$.

Proof. For the first half, assume for contradiction that $p > a$. Then

$$b! + p = a^p \geq a^{p-1} + p \geq a^a + p > a! + p \implies b > a.$$

Then taking modulo a now gives $0 \equiv 0 + p \pmod{a}$, which is obviously impossible.

The second half follows from $b! = a^p - p \geq p! - p > (p - 1)!$. \square

Claim — We have $a = p$ exactly.

Proof. We know $p \geq b$ hence $p \mid b! + p$, so let $a = pk$ for $k < p$. Then $k \mid b!$ yet $k \nmid a^p - p$, contradiction. \square

Let's get the small p out of the way:

- For $p = 2$, checking $2 \leq b \leq 3$ gives $(a, b) = (2, 2)$ only.
- For $p = 3$, checking $3 \leq b \leq 5$ gives $(a, b) = (3, 4)$ only.

Once $p \geq 5$, if $b! = p^p - p = p(p^{p-1} - 1)$ then applying Zsigmondy gets a prime factor $q \equiv 1 \pmod{p - 1}$ which divides $p^{p-1} - 1$. Yet $q \leq b \leq 2p - 2$ and $q \neq p$, contradiction.

§6 IMO 2022/6, proposed by Nikola Petrović (SRB)

Let n be a positive integer. A *Nordic square* is an $n \times n$ board containing all the integers from 1 to n^2 so that each cell contains exactly one number. An *uphill path* is a sequence of one or more cells such that:

1. the first cell in the sequence is a *valley*, meaning the number written is less than all its orthogonal neighbors,
2. each subsequent cell in the sequence is orthogonally adjacent to the previous cell, and
3. the numbers written in the cells in the sequence are in increasing order.

Find, as a function of n , the smallest possible total number of uphill paths in a Nordic square.

Answer: $2n^2 - 2n + 1$.

Bound The lower bound is the “obvious” one:

- For any pair of adjacent cells, say $a > b$, one can extend it to a downhill path (the reverse of an uphill path) by walking downwards until one reaches a valley. This gives $2n(n-1) = 2n^2 - 2n$ uphill paths of length ≥ 2 .
- There is always at least one uphill path of length 1, namely the single cell $\{1\}$ (or indeed any valley).

Construction For the construction, the idea is to build a tree T on the grid such that no two cells not in T are adjacent.

An example of such a grid is shown below for $n = 15$ with T in yellow and cells not in T in black; it generalizes to any $3 \mid n$, and then to any n by deleting the last $n \bmod 3$ rows and either/both of the leftmost/rightmost column.

1	2	3	32		36	62	63	64	93		97	123	124	125
	4		33	34	35		65		94	95	96		126	
6	5	7		37		67	66	68		98		128	127	129
	8		39	38	40		69		100	99	101		130	
10	9	11		41		71	70	72		102		132	131	133
	12		43	42	44		73		104	103	105		134	
14	13	15		45		75	74	76		106		136	135	137
	16		47	46	48		77		108	107	109		138	
18	17	19		49		79	78	80		110		140	139	141
	20		51	50	52		81		112	111	113		142	
22	21	23		53		83	82	84		114		144	143	145
	24		55	54	56		85		116	115	117		146	
26	25	27		57		87	86	88		118		148	147	149
	28		59	58	60		89		120	119	121		150	
30	29	31		61		91	90	92		122		152	151	153

Place 1 anywhere in T and then place all the small numbers at most $|T|$ adjacent to previously placed numbers (example above). Then place the remaining numbers outside T arbitrarily.

By construction, as 1 is the only valley, any uphill path must start from 1. And by construction, it may only reach a given pair of terminal cells in one way, i.e. the downhill paths we mentioned are the only one. End proof.