

IMO 2025 Solution Notes

Evan Chen 《陳誼廷》

20 July 2025

This is a compilation of solutions for the 2025 IMO. The ideas of the solution are a mix of my own work, the solutions provided by the competition organizers, and solutions found by the community. However, all the writing is maintained by me.

These notes will tend to be a bit more advanced and terse than the “official” solutions from the organizers. In particular, if a theorem or technique is not known to beginners but is still considered “standard”, then I often prefer to use this theory anyways, rather than try to work around or conceal it. For example, in geometry problems I typically use directed angles without further comment, rather than awkwardly work around configuration issues. Similarly, sentences like “let $\mathbb{R}$ denote the set of real numbers” are typically omitted entirely.

Corrections and comments are welcome!

Contents

0 Problems	2
1 Solutions to Day 1	3
1.1 IMO 2025/1, proposed by Linus Tang (USA)	3
1.2 IMO 2025/2, proposed by Tran Quang Hung (VNM)	5
1.3 IMO 2025/3, proposed by Lorenzo Sarria (COL)	8
2 Solutions to Day 2	10
2.1 IMO 2025/4, proposed by Paulius Aleknavičius (LIT)	10
2.2 IMO 2025/5, proposed by Massimiliano Foschi, Leonardo Franchi (ITA)	12
2.3 IMO 2025/6, proposed by Zhao Yu Ma and David Lin Kewei (SGP)	14

§0 Problems

1. A line in the plane is called *sunny* if it is not parallel to any of the x -axis, the y -axis, or the line $x + y = 0$.

Let $n \geq 3$ be a given integer. Determine all nonnegative integers k such that there exist n distinct lines in the plane satisfying both of the following:

- for all positive integers a and b with $a + b \leq n + 1$, the point (a, b) lies on at least one of the lines; and
- exactly k of the n lines are sunny.

2. Let Ω and Γ be circles with centres M and N , respectively, such that the radius of Ω is less than the radius of Γ . Suppose Ω and Γ intersect at two distinct points A and B . Line MN intersects Ω at C and Γ at D , so that C, M, N, D lie on MN in that order. Let P be the circumcenter of triangle ACD . Line AP meets Ω again at $E \neq A$ and meets Γ again at $F \neq A$. Let H be the orthocenter of triangle PMN . Prove that the line through H parallel to AP is tangent to the circumcircle of triangle BEF .

3. A function $f: \mathbb{N} \rightarrow \mathbb{N}$ is said to be *bonza* if

$$f(a) \text{ divides } b^a - f(b)^{f(a)}$$

for all positive integers a and b .

Determine the smallest real constant c such that $f(n) \leq cn$ for all bonza functions f and all positive integers n .

4. An infinite sequence $a_1, a_2, \dots$ consists of positive integers has each of which has at least three proper divisors. Suppose that for each $n \geq 1$, a_{n+1} is the sum of the three largest proper divisors of a_n . Determine all possible values of a_1 .
5. Alice and Bazza are playing the *inekoalaty game*, a two-player game whose rules depend on a positive real number λ which is known to both players. On the n th turn of the game (starting with $n = 1$) the following happens:

- If n is odd, Alice chooses a nonnegative real number x_n such that

$$x_1 + x_2 + \dots + x_n \leq \lambda n.$$

- If n is even, Bazza chooses a nonnegative real number x_n such that

$$x_1^2 + x_2^2 + \dots + x_n^2 \leq n.$$

If a player cannot choose a suitable x_n , the game ends and the other player wins. If the game goes on forever, neither player wins. All chosen numbers are known to both players.

Determine all values of λ for which Alice has a winning strategy and all those for which Bazza has a winning strategy.

6. Consider a 2025×2025 grid of unit squares. Matilda wishes to place on the grid some rectangular tiles, possibly of different sizes, such that each side of every tile lies on a grid line and every unit square is covered by at most one tile.

Determine the minimum number of tiles Matilda needs to place so that each row and each column of the grid has exactly one unit square that is not covered by any tile.

§1 Solutions to Day 1

§1.1 IMO 2025/1, proposed by Linus Tang (USA)

Available online at <https://aops.com/community/p35332003>.

Problem statement

A line in the plane is called *sunny* if it is not parallel to any of the x -axis, the y -axis, or the line $x + y = 0$.

Let $n \geq 3$ be a given integer. Determine all nonnegative integers k such that there exist n distinct lines in the plane satisfying both of the following:

- for all positive integers a and b with $a + b \leq n + 1$, the point (a, b) lies on at least one of the lines; and
- exactly k of the n lines are sunny.

The answer is 0, 1, or 3 sunny lines.

In what follows, we draw the grid as equilateral instead of a right triangle; this has no effect on the problem statement but is more symmetric.


We say a *long line* is one of the three lines at the edge of the grid, i.e. one of the (non-sunny) lines passing through n points. The main claim is the following.

Claim — If $n \geq 4$, any set of n lines must have at least one long line.

Proof. Consider the $3(n-1)$ points on the outer edge of the grid. If there was no long line, each of the n lines passes through at most two such points. So we obtain $2n \geq 3(n-1)$, which forces $n \leq 3$. $\square$

Hence, by induction we may repeatedly delete a long line without changing the number of sunny lines until $n = 3$ (and vice-versa: given a construction for smaller n we can increase n by one and add a long line).

We now classify *all* the ways to cover the $1 + 2 + 3 = 6$ points in an $n = 3$ grid with 3 lines.



- If there is a long line (say, the red one in the figure), the remaining $1 + 2 = 3$ points (circled in blue) are covered with two lines. One of the lines passes through 2 points

and must not be sunny; the other line may or may not be sunny. Hence in this case the possible counts of sunny lines are 0 or 1.

- If there is no long line, each of the three lines passes through at most 2 points. But there are 6 total lines, so in fact each line must pass through *exactly* two points. The only way to do this is depicted in the figure in the right. In this case there are 3 sunny lines.

This proves that 0, 1, 3 are the only possible answers.

Remark. The concept of a sunny line is not that important to the problem. The proof above essentially classifies all the ways to cover the $1 + 2 + \cdots + n$ points with exactly n lines. Namely, one should repeatedly take a long line and decrease n until $n = 3$, and then pick one of the finitely many cases for $n = 3$. The count of sunny lines just happens to be whatever is possible for $n = 3$, since long lines are not sunny.

§1.2 IMO 2025/2, proposed by Tran Quang Hung (VNM)

Available online at <https://aops.com/community/p35332018>.

Problem statement

Let Ω and Γ be circles with centres M and N , respectively, such that the radius of Ω is less than the radius of Γ . Suppose Ω and Γ intersect at two distinct points A and B . Line MN intersects Ω at C and Γ at D , so that C, M, N, D lie on MN in that order. Let P be the circumcenter of triangle ACD . Line AP meets Ω again at $E \neq A$ and meets Γ again at $F \neq A$. Let H be the orthocenter of triangle PMN .

Prove that the line through H parallel to AP is tangent to the circumcircle of triangle BEF .

Throughout the solution, we define


$$\begin{aligned}\alpha &:= \angle DCA = \angle BCD \implies \angle PAD = \angle CAB = 90^\circ - \alpha \\ \beta &:= \angle ADC = \angle CDB \implies \angle CAP = \angle BAD = 90^\circ - \beta.\end{aligned}$$

Ignore the points H, M, N for now and focus on the remaining ones.

Claim — We have $\overline{CE} \parallel \overline{AD}$ and $\overline{DF} \parallel \overline{AC}$.

Proof. $\angle AEC = \angle ABC = \angle CAB = 90^\circ - \alpha$. □

Hence, if we let $A' := \overline{CE} \cap \overline{DF}$, we have a parallelogram $ACA'D$. Note in particular that $\overline{BA'} \parallel \overline{CD}$.



Claim — Point T also lies on $\overline{BA'}$ and is also the arc midpoint of $\widehat{EF}$ on (BEF) .

$$\begin{aligned}\angle FEA' &= \angle AEC = \angle ABC = \angle CAB = 90^\circ - \alpha \\ \angle A'FE &= \angle DFA = \angle DBA = \angle BAD = 90^\circ - \beta \\ \angle EA'F &= \alpha + \beta.\end{aligned}$$
$$\angle EA'T = \angle 90^\circ - \angle A'FE = \beta = \angle A'CD = \angle CA'B.$$

Also, we have $\angle ETF = 2\angle EA'F = 2(\alpha + \beta)$ and

$$\angle EBF = \angle EBA + \angle ABF = \angle ECA + \angle ADF$$

$$= \angle A'CA + \angle ADA' = (\alpha + \beta) + (\alpha + \beta) = 2(\alpha + \beta)$$

which proves that T also lies on $\overline{AB'}$. $\square$

We then bring M and N into the picture as follows:

Claim — Point T lies on both lines ME and NF .

Proof. To show that F, T, N are collinear, note that $\triangle FEA' \sim \triangle FAD$ via a homothety at F . This homothety maps T to N . $\square$

We now deal with point H using two claims.

Claim — We have $\overline{MH} \parallel \overline{AD}$ and $\overline{NH} \parallel \overline{AC}$.

Proof. Note that $\overline{MH} \perp \overline{PN}$, but $\overline{PN}$ is the perpendicular bisector of $\overline{AD}$, so in fact $\overline{MH} \parallel \overline{AD}$. Similarly, $\overline{NH} \parallel \overline{AC}$. $\square$

Claim — Lines $\overline{MH}$ and $\overline{NH}$ bisect $\angle NMT$ and $\angle MNT$. In fact, point H is the incenter of $\triangle TMN$, and $\angle NTH = \angle HTM = 90^\circ - (\alpha + \beta)$.

Proof. Hence, $\angle HMN = \angle A'CD = \angle ADC = \beta$. But $\angle TMN = \angle CME = 2\angle CAE = -2(90^\circ - 2\beta) = 2\beta$. That proves $\overline{MH}$ bisects $\angle NMT$; the other one is similar.

To show that H is an incenter (rather than an excenter) and get the last angle equality, we need to temporarily undirect our angles. Assume WLOG that $\triangle ACD$ is directed counterclockwise. The problem condition that C and D are the farther intersections of line MN mean that $\angle NHM = \angle CAD > 90^\circ$. We are also promised C, M, N, D are collinear in that order. Hence the reflections of line MN over lines MH and NH , which meet at T , should meet at a point for which T lies on the same side as H . In other words, $\triangle MTN$ is oriented counterclockwise and contains H .

Working with undirected $\alpha = \angle DCA$ and $\beta = \angle ADC$ with $\alpha + \beta < 90^\circ$,

$$\angle NTH = \angle HTM = \frac{1}{2}\angle NTM = \frac{1}{2}(180^\circ - 2(\alpha + \beta)) = 90^\circ - (\alpha + \beta).$$

This matches the claim and finishes the result. $\square$

Now

$$\angle NFA = 90^\circ - \angle ADF = 90^\circ - (\alpha + \beta) = \angle NTH$$

so $\overline{HT} \parallel \overline{AP}$. And since $TE = TF$, we have the tangency requested too now, as desired.

Remark. There are many other ways to describe the point T . For example, $AMTN$ is a parallelogram and $MBTN$ is an isosceles trapezoid. In coordination, we joked that it was impossible to write a false conjecture.

§1.3 IMO 2025/3, proposed by Lorenzo Sarria (COL)

Available online at <https://aops.com/community/p35332016>.

Problem statement

A function $f: \mathbb{N} \rightarrow \mathbb{N}$ is said to be *bonza* if

$$f(a) \text{ divides } b^a - f(b)^{f(a)}$$

for all positive integers a and b .

Determine the smallest real constant c such that $f(n) \leq cn$ for all bonza functions f and all positive integers n .

The answer is $c = 4$.

Let $P(a, b)$ denote the given statement $f(a) \mid b^a - f(b)^{f(a)}$.

Claim — We have $f(n) \mid n^n$ for all n .

Proof. Take $P(n, n)$. □

Claim — Unless $f = \text{id}$, we have $f(p) = 1$ for all odd primes p .

Proof. Consider any prime q with $f(q) > 1$. Then $f(q)$ is a power of q , and for each n we get

$$P(q, n) \implies q \mid f(q) \mid n^q - f(n)^{f(q)}.$$

Fermat's little theorem now gives $n^q \equiv n \pmod{q}$ and $f(n)^{f(q)} \equiv f(n) \pmod{q}$ (since $f(q)$ is a power of q), and therefore $q \mid n - f(n)$. Hence, unless f is the identity function, only finitely many q could have $f(q) > 1$.

Now let p be any odd prime, and let q be a large prime such that $q \not\equiv 1 \pmod{p}$ (possible for all $p > 2$, say by Dirichlet). Then

$$P(p, q) \implies f(p) \mid q^p - 1^p.$$

The RHS is $q^p - 1 \equiv q - 1 \not\equiv 0 \pmod{p}$, so $f(p) = 1$. □

Claim — We have $f(n) \mid 2^\infty$ for all n .

Proof. If $p \mid f(n)$ is odd then $P(n, p)$ gives $p \mid f(n) \mid p^n - 1^n$, contradiction. □

(In particular, we now know $f(n) = 1$ for all odd n , though we don't use this.)

Claim — We have $f(n) \leq 2^{\nu_2(n)+2}$ for all n .

Proof. Consider $P(n, 5) \implies f(n) \mid 5^n - 1^n$. It's well-known that $\nu_2(5^n - 1) = \nu_2(n) + 2$ for all n . □

This immediately shows $f(n) \leq 4n$ for all n , hence $c = 4$ in the problem statement works.

For the construction, the simplest one seems to be

$$f(n) = \begin{cases} 1 & n \text{ odd} \\ 16 & n = 4 \\ 2 & n \text{ even, } n \neq 4 \end{cases}$$

which is easily checked to work and has $f(4) = 16$.

Remark. With a little more case analysis we can classify all functions f . The two trivial solutions are $f(n) = n$ and $f(n) = 1$; the others are described by writing $f(n) = 2^{e(n)}$ for any function e satisfying

- $e(n) = 0$ for odd n ;
- $1 \leq e(2) \leq 2$;
- $1 \leq e(n) \leq \nu_2(n) + 2$ for even $n > 2$.

This basically means that there are almost no additional constraints beyond what is suggested by the latter two claims.

§2 Solutions to Day 2

§2.1 IMO 2025/4, proposed by Paulius Aleknavičius (LIT)

Available online at <https://aops.com/community/p35347364>.

Problem statement

An infinite sequence $a_1, a_2, \dots$ consists of positive integers has each of which has at least three proper divisors. Suppose that for each $n \geq 1$, a_{n+1} is the sum of the three largest proper divisors of a_n . Determine all possible values of a_1 .

The answer is $a_1 = 12^e \cdot 6 \cdot \ell$ for any $e, \ell \geq 0$ with $\gcd(\ell, 10) = 1$.

Let $\mathbf{S}$ denote the set of positive integers with at least three divisors. For $x \in \mathbf{S}$, let $\psi(x)$ denote the sum of the three largest ones, so that $\psi(a_n) = a_{n+1}$.

¶ **Proof that all such a_1 work.** Let $x = 12^e \cdot 6 \cdot \ell \in \mathbf{S}$ with $\gcd(\ell, 10) = 1$. As $\frac{1}{2} + \frac{1}{3} + \frac{1}{4} = \frac{13}{12}$, we get

$$\psi(x) = \begin{cases} x & e = 0 \\ \frac{13}{12}x & e > 0 \end{cases}$$

so by induction on the value of e we see that $\psi(x) \in \mathbf{S}$ (the base $e = 0$ coming from ψ fixing x).

¶ **Proof that all a_1 are of this form.** In what follows x is always an element of $\mathbf{S}$, not necessarily an element of the sequence.

Claim — Let $x \in \mathbf{S}$. If $2 \mid \psi(x)$ then $2 \mid x$.

Proof. If x is odd then every divisor of x is odd, so $\psi(x)$ is the sum of three odd numbers. $\square$

Claim — Let $x \in \mathbf{S}$. If $6 \mid \psi(x)$ then $6 \mid x$.

Proof. We consider only x even because of the previous claim. We prove the contrapositive that $3 \nmid x \implies 3 \nmid \psi(x)$.

- If $4 \mid x$, then letting d be the third largest proper divisor of x ,

$$\psi(x) = \frac{x}{2} + \frac{x}{4} + d = \frac{3}{4}x + d \equiv d \not\equiv 0 \pmod{3}.$$

- Otherwise, let $p \mid x$ be the smallest prime dividing x , with $p > 3$. If the third smallest nontrivial divisor of x is $2p$, then

$$\psi(x) = \frac{x}{2} + \frac{x}{p} + \frac{x}{2p} = \frac{3}{2p}x + \frac{x}{2} \equiv \frac{x}{2} \not\equiv 0 \pmod{3}.$$

If the third smallest nontrivial divisor of x is instead an odd prime q , then

$$\psi(x) = \frac{x}{2} + \frac{x}{p} + \frac{x}{q} \equiv 1 + 0 + 0 \equiv 1 \pmod{2}.$$

□

To tie these two claims into the problem, we assert:

Claim — Every a_i must be divisible by 6.

Proof. The idea is to combine the previous two claims (which have no dependence on the sequence) with a size argument.

- For odd $x \in \mathbf{S}$ note that $\psi(x) < \left(\frac{1}{3} + \frac{1}{5} + \frac{1}{7}\right)x < x$ and $\psi(x)$ is still odd. So if any a_i is odd the sequence is strictly decreasing and that's impossible. Hence, we may assume $a_1, a_2, \dots$ are all even.
- If $x \in \mathbf{S}$ is even but $3 \nmid x$ then $\psi(x) < \left(\frac{1}{2} + \frac{1}{4} + \frac{1}{5}\right)x < x$ and $\psi(x)$ is still not a multiple of 3. So if any a_i is not divisible by 3 the sequence is again strictly decreasing. □

On the other hand, if x is a multiple of 6, we have the following formula for $\psi(x)$:

$$\psi(x) = \begin{cases} \frac{13}{12}x & 4 \mid x \\ \frac{31}{30}x & 4 \nmid x \text{ but } 5 \mid x \\ x & 4 \nmid x \text{ and } 5 \nmid x. \end{cases}$$

Looking back on our sequence of a_i (which are all multiples of 6), the center case cannot happen with our a_i , because $\frac{31}{30}x$ is odd when $x \equiv 2 \pmod{4}$. Hence in actuality

$$a_{n+1} = \frac{13}{12}a_n \quad \text{or} \quad a_{n+1} = a_n$$

for every n .

Let T be the smallest index such that $a_T = a_{T+1} = a_{T+2} = \dots$ (it must exist because we cannot multiply by $\frac{13}{12}$ forever). Then we can exactly describe the sequence by

$$a_n = a_1 \cdot \left(\frac{13}{12}\right)^{\min(n, T)-1}.$$

Hence $a_1 = \left(\frac{12}{13}\right)^{T-1} a_T$, and since a_T is a multiple of 6 not divisible by 4 or 5, it follows a_1 has the required form.

§2.2 IMO 2025/5, proposed by Massimiliano Foschi, Leonardo Franchi (ITA)

Available online at <https://aops.com/community/p35341177>.

Problem statement

Alice and Bazza are playing the *inekoalaty game*, a two-player game whose rules depend on a positive real number λ which is known to both players. On the n th turn of the game (starting with $n = 1$) the following happens:

- If n is odd, Alice chooses a nonnegative real number x_n such that

$$x_1 + x_2 + \cdots + x_n \leq \lambda n.$$

- If n is even, Bazza chooses a nonnegative real number x_n such that

$$x_1^2 + x_2^2 + \cdots + x_n^2 \leq n.$$

If a player cannot choose a suitable x_n , the game ends and the other player wins. If the game goes on forever, neither player wins. All chosen numbers are known to both players.

Determine all values of λ for which Alice has a winning strategy and all those for which Bazza has a winning strategy.

The answer is that Alice has a winning strategy for $\lambda > 1/\sqrt{2}$, and Bazza has a winning strategy for $\lambda < 1/\sqrt{2}$. (Neither player can guarantee winning for $\lambda = 1/\sqrt{2}$.)

We divide the proof into two parts.

¶ **Alice's strategy when $\lambda \geq 1/\sqrt{2}$.** Consider the strategy where Alice always plays $x_{2i+1} = 0$ for $i = 0, \dots, k-1$.

In this situation, when $n = 2k+1$ we have

$$\begin{aligned} \sum_{i=1}^{2k} x_i &= 0 + x_2 + 0 + x_4 + \cdots + 0 + x_{2k} \\ &\leq k \cdot \sqrt{\frac{x_2^2 + \cdots + x_{2k}^2}{k}} = \sqrt{2} \cdot k < \lambda \cdot (2k+1) \end{aligned}$$

and so the choices for x_{2k+1} are

$$x_{2k+1} \in [0, \lambda \cdot (2k+1) - \sqrt{2}k]$$

which is nonempty. Hence Alice can't ever lose with this strategy.

But suppose further $\lambda > \frac{1}{\sqrt{2}}$; we show Alice can win. Choose k large enough that

$$\sqrt{2} \cdot k < \lambda \cdot (2k+1) - \sqrt{2k+2}.$$

Then on the $(2k+1)$ st turn, Alice can (after playing 0 on all earlier turns) play a number greater than $\sqrt{2k+2}$ and cause Bob to lose.

¶ **Bob strategy when $\lambda \leq 1/\sqrt{2}$.** Consider the strategy where Bob always plays $x_{2i+2} = \sqrt{2 - x_{2i+1}^2}$ for all $i = 0, \dots, k-1$ (i.e. that is, the largest possible value Bob can play).

To analyze Bob's choices on each of his turns, we first need to estimate x_{2k+1} . We do this by writing

$$\begin{aligned} \lambda \cdot (2k+1) &\geq x_1 + x_2 + \dots + x_{2k+1} \\ &= \left(x_1 + \sqrt{2 - x_1^2}\right) + \left(x_3 + \sqrt{2 - x_3^2}\right) \\ &\quad + \dots + \left(x_{2k-1} + \sqrt{2 - x_{2k-1}^2}\right) + x_{2k+1} \\ &\geq \underbrace{\sqrt{2} + \dots + \sqrt{2}}_{k \text{ times}} + x_{2k+1} = \sqrt{2} \cdot k + x_{2k+1} \end{aligned}$$

where we have used the fact that $t + \sqrt{2 - t^2} \geq 2$ for all $t \geq 0$. This means that

$$x_{2k+1} \leq \lambda \cdot (2k+1) - \sqrt{2}k < \sqrt{2}.$$

And $x_1^2 + x_2^2 + \dots + x_{2k}^2 + x_{2k+1}^2 = (2 + \dots + 2) + x_{2k+1}^2$, Bob can indeed choose $x_{2k+2} = \sqrt{2 - x_{2k+1}^2}$ and always has a move.

But suppose further $\lambda < 1/\sqrt{2}$. Then the above calculation *also* shows that Alice couldn't have made a valid choice for large enough k , since $\lambda \cdot (2k+1) - \sqrt{2}k < 0$ for large k .

Remark. In the strategies above, we saw that Alice prefers to always play 0 and Bob prefers to always play as large as possible. One could consider what happens in the opposite case:

- If Alice tries to always play the largest number possible, her strategy still wins for $\lambda > 1$.
- If Bob tries to always play 0, Alice can win no matter the value for $\lambda > 0$.

§2.3 IMO 2025/6, proposed by Zhao Yu Ma and David Lin Kewei (SGP)

Available online at <https://aops.com/community/p35341197>.

Problem statement

Consider a 2025×2025 grid of unit squares. Matilda wishes to place on the grid some rectangular tiles, possibly of different sizes, such that each side of every tile lies on a grid line and every unit square is covered by at most one tile.

Determine the minimum number of tiles Matilda needs to place so that each row and each column of the grid has exactly one unit square that is not covered by any tile.

The answer is $2112 = 2025 + 2 \cdot 45 - 3$. In general, the answer turns out to be $\lceil n + 2\sqrt{n} - 3 \rceil$, but when n is not a perfect square the solution is more complicated.

¶ **Construction.** To be added.

¶ **Bound.** To be added.

Remark. Ironically, the construction obtaining the answer in the floor pattern at Sunshine Coast airport, the closest airport to the site of the exam. See [this image](#) or [this image](#).