

JMO 2011 Solution Notes

EVAN CHEN 《陳誼廷》

11 December 2023

This is a compilation of solutions for the 2011 JMO. Some of the solutions are my own work, but many are from the official solutions provided by the organizers (for which they hold any copyrights), and others were found by users on the Art of Problem Solving forums.

These notes will tend to be a bit more advanced and terse than the “official” solutions from the organizers. In particular, if a theorem or technique is not known to beginners but is still considered “standard”, then I often prefer to use this theory anyways, rather than try to work around or conceal it. For example, in geometry problems I typically use directed angles without further comment, rather than awkwardly work around configuration issues. Similarly, sentences like “let \mathbb{R} denote the set of real numbers” are typically omitted entirely.

Corrections and comments are welcome!

Contents

0 Problems	2
1 Solutions to Day 1	3
1.1 JMO 2011/1, proposed by Titu Andreescu	3
1.2 JMO 2011/2, proposed by Titu Andreescu	4
1.3 JMO 2011/3, proposed by Zuming Feng	5
2 Solutions to Day 2	7
2.1 JMO 2011/4, proposed by Gabriel Carroll	7
2.2 JMO 2011/5, proposed by Zuming Feng	8
2.3 JMO 2011/6, proposed by Sam Vandervelde	9

§0 Problems

1. Find all positive integers n such that $2^n + 12^n + 2011^n$ is a perfect square.
2. Let a, b, c be positive real numbers such that $a^2 + b^2 + c^2 + (a + b + c)^2 \leq 4$. Prove that

$$\frac{ab+1}{(a+b)^2} + \frac{bc+1}{(b+c)^2} + \frac{ca+1}{(c+a)^2} \geq 3.$$

3. For a point $P = (a, a^2)$ in the coordinate plane, let $\ell(P)$ denote the line passing through P with slope $2a$. Consider the set of triangles with vertices of the form $P_1 = (a_1, a_1^2)$, $P_2 = (a_2, a_2^2)$, $P_3 = (a_3, a_3^2)$, such that the intersection of the lines $\ell(P_1), \ell(P_2), \ell(P_3)$ form an equilateral triangle Δ . Find the locus of the center of Δ as $P_1 P_2 P_3$ ranges over all such triangles.
4. A *word* is defined as any finite string of letters. A word is a *palindrome* if it reads the same backwards and forwards. Let a sequence of words W_0, W_1, W_2, \dots be defined as follows: $W_0 = a, W_1 = b$, and for $n \geq 2$, W_n is the word formed by writing W_{n-2} followed by W_{n-1} . Prove that for any $n \geq 1$, the word formed by writing $W_1, W_2, W_3, \dots, W_n$ in succession is a palindrome.
5. Points A, B, C, D, E lie on a circle ω and point P lies outside the circle. The given points are such that (i) lines PB and PD are tangent to ω , (ii) P, A, C are collinear, and (iii) $\overline{DE} \parallel \overline{AC}$. Prove that \overline{BE} bisects \overline{AC} .
6. Consider the assertion that for each positive integer $n \geq 2$, the remainder upon dividing 2^{2^n} by $2^n - 1$ is a power of 4. Either prove the assertion or find (with proof) a counterexample.

§1 Solutions to Day 1

§1.1 JMO 2011/1, proposed by Titu Andreescu

Available online at <https://aops.com/community/p2254778>.

Problem statement

Find all positive integers n such that $2^n + 12^n + 2011^n$ is a perfect square.

The answer $n = 1$ works, because $2^1 + 12^1 + 2011^1 = 45^2$. We prove it's the only one.

- If $n \geq 2$ is even, then modulo 3 we have $2^n + 12^n + 2011^n \equiv 1 + 0 + 1 \equiv 2 \pmod{3}$ so it is not a square.
- If $n \geq 3$ is odd, then modulo 4 we have $2^n + 12^n + 2011^n \equiv 0 + 0 + 3 \equiv 3 \pmod{4}$ so it is not a square.

This completes the proof.

§1.2 JMO 2011/2, proposed by Titu Andreescu

Available online at <https://aops.com/community/p2254758>.

Problem statement

Let a, b, c be positive real numbers such that $a^2 + b^2 + c^2 + (a + b + c)^2 \leq 4$. Prove that

$$\frac{ab+1}{(a+b)^2} + \frac{bc+1}{(b+c)^2} + \frac{ca+1}{(c+a)^2} \geq 3.$$

The condition becomes $2 \geq a^2 + b^2 + c^2 + ab + bc + ca$. Therefore,

$$\begin{aligned} \sum_{\text{cyc}} \frac{2ab+2}{(a+b)^2} &\geq \sum_{\text{cyc}} \frac{2ab + (a^2 + b^2 + c^2 + ab + bc + ca)}{(a+b)^2} \\ &= \sum_{\text{cyc}} \frac{(a+b)^2 + (c+a)(c+b)}{(a+b)^2} \\ &= 3 + \sum_{\text{cyc}} \frac{(c+a)(c+b)}{(a+b)^2} \\ &\geq 3 + 3 \sqrt[3]{\prod_{\text{cyc}} \frac{(c+a)(c+b)}{(a+b)^2}} = 3 + 3 = 6 \end{aligned}$$

with the last line by AM-GM. This completes the proof.

§1.3 JMO 2011/3, proposed by Zuming Feng

Available online at <https://aops.com/community/p2254823>.

Problem statement

For a point $P = (a, a^2)$ in the coordinate plane, let $\ell(P)$ denote the line passing through P with slope $2a$. Consider the set of triangles with vertices of the form $P_1 = (a_1, a_1^2)$, $P_2 = (a_2, a_2^2)$, $P_3 = (a_3, a_3^2)$, such that the intersection of the lines $\ell(P_1), \ell(P_2), \ell(P_3)$ form an equilateral triangle Δ . Find the locus of the center of Δ as $P_1P_2P_3$ ranges over all such triangles.

The answer is the line $y = -1/4$. I did not find this problem inspiring, so I will not write out most of the boring calculations since most solutions are just going to be “use Cartesian coordinates and grind all the way through”.

The “nice” form of the main claim is as follows (which is certainly overkill for the present task, but is too good to resist including):

Claim (Naoki Sato) — In general, the orthocenter of Δ lies on the directrix $y = -1/4$ of the parabola (even if the triangle Δ is not equilateral).

Proof. By writing out the equation $y = 2a_ix - a_i^2$ for $\ell(P_i)$, we find the vertices of the triangle are located at

$$\left(\frac{a_1 + a_2}{2}, a_1a_2\right); \quad \left(\frac{a_2 + a_3}{2}, a_2a_3\right); \quad \left(\frac{a_3 + a_1}{2}, a_3a_1\right).$$

The coordinates of the orthocenter can be checked explicitly to be

$$H = \left(\frac{a_1 + a_2 + a_3 + 4a_1a_2a_3}{2}, -\frac{1}{4}\right).$$

An advanced synthetic proof of this fact is given at <https://aops.com/community/p2255814>. \square

This claim already shows that every point lies on $y = -1/4$. We now turn to showing that, even when restricted to equilateral triangles, we can achieve every point on $y = -1/4$. In what follows $a = a_1$, $b = a_2$, $c = a_3$ for legibility.

Claim — Lines $\ell(a)$, $\ell(b)$, $\ell(c)$ form an equilateral triangle if and only if

$$\begin{aligned} a + b + c &= -12abc \\ ab + bc + ca &= -\frac{3}{4}. \end{aligned}$$

Moreover, the x -coordinate of the equilateral triangle is $\frac{1}{3}(a + b + c)$.

Proof. The triangle is equilateral if and only if the centroid and orthocenter coincide, i.e.

$$\left(\frac{a + b + c}{3}, \frac{ab + bc + ca}{3}\right) = G = H = \left(\frac{a + b + c + 4abc}{2}, -\frac{1}{4}\right).$$

Setting the x and y coordinates equal, we derive the claimed equations. \square

Let λ be any real number. We are tasked to show that

$$P(X) = X^3 - 3\lambda \cdot X^2 - \frac{3}{4}X + \frac{\lambda}{4}$$

has three real roots (with multiplicity); then taking those roots as (a, b, c) yields a valid equilateral-triangle triple whose x -coordinate is exactly λ , by the previous claim.

To prove that, pick the values

$$P(-\sqrt{3}/2) = -2\lambda$$

$$P(0) = \frac{1}{4}\lambda$$

$$P(\sqrt{3}/2) = -2\lambda.$$

The intermediate value theorem (at least for $\lambda \neq 0$) implies that P should have at least two real roots now, and since P has degree 3, it has all real roots. That's all.

§2 Solutions to Day 2

§2.1 JMO 2011/4, proposed by Gabriel Carroll

Available online at <https://aops.com/community/p2254808>.

Problem statement

A *word* is defined as any finite string of letters. A word is a *palindrome* if it reads the same backwards and forwards. Let a sequence of words W_0, W_1, W_2, \dots be defined as follows: $W_0 = a, W_1 = b$, and for $n \geq 2$, W_n is the word formed by writing W_{n-2} followed by W_{n-1} . Prove that for any $n \geq 1$, the word formed by writing $W_1, W_2, W_3, \dots, W_n$ in succession is a palindrome.

To aid in following the solution, here are the first several words:

$$\begin{aligned} W_0 &= a \\ W_1 &= b \\ W_2 &= ab \\ W_3 &= bab \\ W_4 &= abbab \\ W_5 &= bababbab \\ W_6 &= abbabbababbab \\ W_7 &= bababbababbababbab \end{aligned}$$

We prove that $W_1 W_2 \dots W_n$ is a palindrome by induction on n . The base cases $n = 1, 2, 3, 4$ can be verified by hand.

For the inductive step, we let \overline{X} denote the word X written backwards. Then

$$\begin{aligned} W_1 W_2 \dots W_{n-3} W_{n-2} W_{n-1} W_n &\stackrel{\text{IH}}{=} (\overline{W_{n-1} W_{n-2} W_{n-3}} \dots \overline{W_2 W_1}) W_n \\ &= (\overline{W_{n-1} W_{n-2} W_{n-3}} \dots \overline{W_2 W_1}) W_{n-2} W_{n-1} \\ &= \overline{W_{n-1} W_{n-2}} (\overline{W_{n-3}} \dots \overline{W_2 W_1}) W_{n-2} W_{n-1} \end{aligned}$$

with the first equality being by the induction hypothesis. By induction hypothesis again the inner parenthesized term is also a palindrome, and so this completes the proof.

§2.2 JMO 2011/5, proposed by Zuming Feng


Available online at <https://aops.com/community/p2254813>.

Problem statement

Points A, B, C, D, E lie on a circle ω and point P lies outside the circle. The given points are such that (i) lines PB and PD are tangent to ω , (ii) P, A, C are collinear, and (iii) $\overline{DE} \parallel \overline{AC}$. Prove that \overline{BE} bisects \overline{AC} .

We present two solutions.

¶ **First solution using harmonic bundles.** Let $M = \overline{BE} \cap \overline{AC}$ and let ∞ be the point at infinity along $\overline{DE} \parallel \overline{AC}$.



Note that $ABCD$ is harmonic, so

$$-1 = (AC; BD) \stackrel{E}{=} (AC; M\infty)$$

implying M is the midpoint of \overline{AC} .

¶ **Second solution using complex numbers (Cynthia Du).** Suppose we let b, d, e be free on unit circle, so $p = \frac{2bd}{b+d}$. Then $d/c = a/e$, and $a + c = p + ac\overline{p}$. Consequently,

$$\begin{aligned} ac &= de \\ \frac{1}{2}(a + c) &= \frac{bd}{b+d} + de \cdot \frac{1}{b+d} = \frac{d(b+e)}{b+d}. \\ \frac{a+c}{2ac} &= \frac{(b+e)}{e(b+d)}. \end{aligned}$$

From here it's easy to see

$$\frac{a+c}{2} + \frac{a+c}{2ac} \cdot be = b+e$$

which is what we wanted to prove.

§2.3 JMO 2011/6, proposed by Sam Vandervelde

Available online at <https://aops.com/community/p2254810>.

Problem statement

Consider the assertion that for each positive integer $n \geq 2$, the remainder upon dividing 2^{2^n} by $2^n - 1$ is a power of 4. Either prove the assertion or find (with proof) a counterexample.

We claim $n = 25$ is a counterexample. Since $2^{25} \equiv 2^0 \pmod{2^{25} - 1}$, we have

$$2^{2^{25}} \equiv 2^{2^{25} \bmod 25} \equiv 2^7 \pmod{2^{25} - 1}$$

and the right-hand side is actually the remainder, since $0 < 2^7 < 2^{25}$. But 2^7 is not a power of 4.

Remark. Really, the problem is just equivalent for asking 2^n to have odd remainder when divided by n .