

# JMO 2014 Solution Notes

COMPILED BY EVAN CHEN

March 22, 2020

This is an compilation of solutions for the 2014 JMO. Some of the solutions are my own work, but many are from the official solutions provided by the organizers (for which they hold any copyrights), and others were found on the Art of Problem Solving forums.

Corrections and comments are welcome!

## Contents

<a href="#">0 Problems</a>	2
<a href="#">1 JMO 2014/1, proposed by Titu Andreescu</a>	3
<a href="#">2 JMO 2014/2, proposed by Zuming Feng</a>	4
<a href="#">3 JMO 2014/3, proposed by Titu Andreescu</a>	6
<a href="#">4 JMO 2014/4, proposed by Palmer Mebane</a>	8
<a href="#">5 JMO 2014/5, proposed by Palmer Mebane</a>	9
<a href="#">6 JMO 2014/6, proposed by Titu Andreescu, Cosmin Pohoata</a>	11

## §0 Problems

1. Let  $a, b, c$  be real numbers greater than or equal to 1. Prove that

$$\min \left( \frac{10a^2 - 5a + 1}{b^2 - 5b + 10}, \frac{10b^2 - 5b + 1}{c^2 - 5c + 10}, \frac{10c^2 - 5c + 1}{a^2 - 5a + 10} \right) \leq abc.$$

2. Let  $\triangle ABC$  be a non-equilateral, acute triangle with  $\angle A = 60^\circ$ , and let  $O$  and  $H$  denote the circumcenter and orthocenter of  $\triangle ABC$ , respectively.
- (a) Prove that line  $OH$  intersects both segments  $AB$  and  $AC$  at two points  $P$  and  $Q$ , respectively.
- (b) Denote by  $s$  and  $t$  the respective areas of triangle  $APQ$  and quadrilateral  $BPQC$ . Determine the range of possible values for  $s/t$ .
3. Find all  $f : \mathbb{Z} \rightarrow \mathbb{Z}$  such that

$$xf(2f(y) - x) + y^2f(2x - f(y)) = \frac{f(x)^2}{x} + f(yf(y))$$

for all  $x, y \in \mathbb{Z}$  such that  $x \neq 0$ .

4. Let  $b \geq 2$  be a fixed integer, and let  $s_b(n)$  denote the sum of the base- $b$  digits of  $n$ . Show that there are infinitely many positive integers that cannot be represented in the form  $n + s_b(n)$  where  $n$  is a positive integer.
5. Let  $k$  be a positive integer. Two players  $A$  and  $B$  play a game on an infinite grid of regular hexagons. Initially all the grid cells are empty. Then the players alternately take turns with  $A$  moving first. In her move,  $A$  may choose two adjacent hexagons in the grid which are empty and place a counter in both of them. In his move,  $B$  may choose any counter on the board and remove it. If at any time there are  $k$  consecutive grid cells in a line all of which contain a counter,  $A$  wins. Find the minimum value of  $k$  for which  $A$  cannot win in a finite number of moves, or prove that no such minimum value exists.
6. Let  $ABC$  be a triangle with incenter  $I$ , incircle  $\gamma$  and circumcircle  $\Gamma$ . Let  $M, N, P$  be the midpoints of  $\overline{BC}, \overline{CA}, \overline{AB}$  and let  $E, F$  be the tangency points of  $\gamma$  with  $\overline{CA}$  and  $\overline{AB}$ , respectively. Let  $U, V$  be the intersections of line  $EF$  with line  $MN$  and line  $MP$ , respectively, and let  $X$  be the midpoint of arc  $BAC$  of  $\Gamma$ .
- (a) Prove that  $I$  lies on ray  $CV$ .
- (b) Prove that line  $XI$  bisects  $\overline{UV}$ .

**§1 JMO 2014/1, proposed by Titu Andreescu**

Let  $a, b, c$  be real numbers greater than or equal to 1. Prove that

$$\min \left( \frac{10a^2 - 5a + 1}{b^2 - 5b + 10}, \frac{10b^2 - 5b + 1}{c^2 - 5c + 10}, \frac{10c^2 - 5c + 1}{a^2 - 5a + 10} \right) \leq abc.$$

---

Notice that

$$\frac{10a^2 - 5a + 1}{a^2 - 5a + 10} \leq a^3$$

since it rearranges to  $(a - 1)^5 \geq 0$ . Cyclically multiply to get

$$\prod_{\text{cyc}} \left( \frac{10a^2 - 5a + 1}{b^2 - 5b + 10} \right) \leq (abc)^3$$

and the minimum is at most the geometric mean.

## §2 JMO 2014/2, proposed by Zuming Feng

Let  $\triangle ABC$  be a non-equilateral, acute triangle with  $\angle A = 60^\circ$ , and let  $O$  and  $H$  denote the circumcenter and orthocenter of  $\triangle ABC$ , respectively.

- (a) Prove that line  $OH$  intersects both segments  $AB$  and  $AC$  at two points  $P$  and  $Q$ , respectively.
- (b) Denote by  $s$  and  $t$  the respective areas of triangle  $APQ$  and quadrilateral  $BPQC$ . Determine the range of possible values for  $s/t$ .


We begin with some synthetic work. Let  $I$  denote the incenter, and recall (“fact 5”) that the arc midpoint  $M$  is the center of  $(BIC)$ , which we denote by  $\gamma$ .

Now we have that

$$\angle BOC = \angle BIC = \angle BHC = 120^\circ.$$

Since all three centers lie inside  $ABC$  (as it was acute), and hence on the opposite side of  $\overline{BC}$  as  $M$ , it follows that  $O, I, H$  lie on minor arc  $BC$  of  $\gamma$ .

We note this implies (a) already, as line  $OH$  meets line  $BC$  outside of segment  $BC$ .



**Claim** — Triangle  $APQ$  is equilateral with side length  $\frac{b+c}{3}$ .

*Proof.* Let  $R$  be the circumradius. We have  $R = OM = OA = MH$ , and even  $AH = 2R \cos A = R$ , so  $AOMH$  is a rhombus. Thus  $\overline{OH} \perp \overline{AM}$  and in this way we derive that  $\triangle APQ$  is isosceles, hence equilateral.

Finally, since  $\angle PBH = 30^\circ$ , and  $\angle BPH = 120^\circ$ , it follows that  $\triangle BPH$  is isosceles and  $BP = PH$ . Similarly,  $CQ = QH$ . So  $b+c = AP+BP+AQ+QC = AP+AQ+PQ$  as needed.  $\square$

Finally, we turn to the boring task of extracting the numerical answer. We have

$$\frac{s}{s+t} = \frac{[APQ]}{[ABC]} = \frac{\frac{\sqrt{3}}{4} \left(\frac{b+c}{3}\right)^2}{\frac{\sqrt{3}}{4} bc} = \frac{b^2 + bc + c^2}{9bc} = \frac{1}{9} \left(1 + \frac{b}{c} + \frac{c}{b}\right).$$

So the problem is reduced to analyzing the behavior of  $b/c$ . For this, we imagine fixing  $\Gamma$  the circumcircle of  $ABC$ , as well as the points  $B$  and  $C$ . Then as we vary  $A$  along the “topmost” arc of measure  $120^\circ$ , we find  $b/c$  is monotonic with values  $1/2$  and  $2$  at endpoints, and by continuity all values  $b/c \in (1/2, 2)$  can be achieved.

So

$$\frac{1}{2} < \frac{b}{c} < 2 \implies 4/9 < \frac{s}{s+t} < 1/2 \implies 4/5 < \frac{s}{t} < 1$$

as needed.

### §3 JMO 2014/3, proposed by Titu Andreescu

Find all  $f : \mathbb{Z} \rightarrow \mathbb{Z}$  such that

$$xf(2f(y) - x) + y^2f(2x - f(y)) = \frac{f(x)^2}{x} + f(yf(y))$$

for all  $x, y \in \mathbb{Z}$  such that  $x \neq 0$ .

The answer is  $f(x) \equiv 0$  and  $f(x) \equiv x^2$ . Check that these work.

Now let's prove these are the only solutions. Put  $y = 0$  to obtain

$$xf(2f(0) - x) = \frac{f(x)^2}{x} + f(0).$$

Now we claim  $\boxed{f(0) = 0}$ . If not, select a prime  $p \nmid f(0)$  and put  $x = p \neq 0$ . In the above, we find that  $p \mid f(p)^2$ , so  $p \mid f(p)$  and hence  $p \mid \frac{f(p)^2}{p}$ . From here we derive  $p \mid f(0)$ , contradiction. Hence

$$f(0) = 0.$$

The above then implies that

$$x^2f(-x) = f(x)^2$$

holds for all nonzero  $x$ , but also for  $x = 0$ . Let us now check that  $f$  is an even function. In the above, we may also derive  $f(-x)^2 = x^2f(x)$ . If  $f(x) \neq f(-x)$  (and hence  $x \neq 0$ ), then subtracting the above and factoring implies that  $f(x) + f(-x) = -x^2$ ; we can then obtain by substituting the relation

$$\left[ f(x) + \frac{1}{2}x^2 \right]^2 = -\frac{3}{4}x^4 < 0$$

which is impossible. This means  $f(x)^2 = x^2f(x)$ , thus

$$f(x) \in \{0, x^2\} \quad \forall x.$$

Now suppose there exists a nonzero integer  $t$  with  $f(t) = 0$ . We will prove that  $f(x) \equiv 0$ . Put  $y = t$  in the given to obtain that

$$t^2f(2x) = 0$$

for any integer  $x \neq 0$ , and hence conclude that  $f(2\mathbb{Z}) \equiv 0$ . Then selecting  $x = 2k \neq 0$  in the given implies that

$$y^2f(4k - f(y)) = f(yf(y)).$$

Assume for contradiction that  $f(m) = m^2$  now for some odd  $m \neq 0$ . Evidently

$$m^2f(4k - m^2) = f(m^3).$$

If  $f(m^3) \neq 0$  this forces  $f(4k - m^2) \neq 0$ , and hence  $m^2(4k - m^2)^2 = m^6$  for arbitrary  $k \neq 0$ , which is clearly absurd. That means

$$f(4k - m^2) = f(m^2 - 4k) = f(m^3) = 0$$

for each  $k \neq 0$ . Since  $m$  is odd,  $m^2 \equiv 1 \pmod{4}$ , and so  $f(n) = 0$  for all  $n$  other than  $\pm m^2$  (since we cannot select  $k = 0$ ).

Now  $f(m) = m^2$  means that  $m = \pm 1$ . Hence either  $f(x) \equiv 0$  or

$$f(x) = \begin{cases} 1 & x = \pm 1 \\ 0 & \text{otherwise.} \end{cases}$$

To show that the latter fails, we simply take  $x = 5$  and  $y = 1$  in the given.

Hence, the only solutions are  $f(x) \equiv 0$  and  $f(x) \equiv x^2$ .

## §4 JMO 2014/4, proposed by Palmer Mebane

Let  $b \geq 2$  be a fixed integer, and let  $s_b(n)$  denote the sum of the base- $b$  digits of  $n$ . Show that there are infinitely many positive integers that cannot be represented in the form  $n + s_b(n)$  where  $n$  is a positive integer.

---

For brevity let  $f(n) = n + s_b(n)$ . Select any integer  $M$ . Observe that  $f(x) \geq b^{2M}$  for any  $x \geq b^{2M}$ , but also  $f(b^{2M} - k) \geq b^{2M}$  for  $k = 1, 2, \dots, M$ , since the base- $b$  expansion of  $b^{2M} - k$  will start out with at least  $M$  digits  $b - 1$ .

Thus  $f$  omits at least  $M$  values in  $[1, b^{2M}]$  for any  $M$ .




## §5 JMO 2014/5, proposed by Palmer Mebane

Let  $k$  be a positive integer. Two players  $A$  and  $B$  play a game on an infinite grid of regular hexagons. Initially all the grid cells are empty. Then the players alternately take turns with  $A$  moving first. In her move,  $A$  may choose two adjacent hexagons in the grid which are empty and place a counter in both of them. In his move,  $B$  may choose any counter on the board and remove it. If at any time there are  $k$  consecutive grid cells in a line all of which contain a counter,  $A$  wins. Find the minimum value of  $k$  for which  $A$  cannot win in a finite number of moves, or prove that no such minimum value exists.


---

The answer is  $k = 6$ .

**Proof that  $A$  cannot win if  $k = 6$ .** We give a strategy for  $B$  to prevent  $A$ 's victory. Shade in every third cell, as shown in the figure below. Then  $A$  can never cover two shaded cells simultaneously on her turn. Now suppose  $B$  always removes a counter on a shaded cell (and otherwise does whatever he wants). Then he can prevent  $A$  from ever getting six consecutive counters, because any six consecutive cells contain two shaded cells.




**Example of a strategy for  $A$  when  $k = 5$ .** We describe a winning strategy for  $A$  explicitly. Note that after  $B$ 's first turn there is one counter, so then  $A$  may create an equilateral triangle, and hence after  $B$ 's second turn there are two consecutive counters. Then, on her third turn,  $A$  places a pair of counters two spaces away on the same line. Label the two inner cells  $x$  and  $y$  as shown below.



Now it is  $B$ 's turn to move; in order to avoid losing immediately, he must remove either  $x$  or  $y$ . Then on any subsequent turn,  $A$  can replace  $x$  or  $y$  (whichever was removed) and add one more adjacent counter. This continues until either  $x$  or  $y$  has all its neighbors filled (we ask  $A$  to do so in such a way that she avoids filling in the two central cells between  $x$  and  $y$  as long as possible).

So, let's say without loss of generality (by symmetry) that  $x$  is completely surrounded by tokens. Again,  $B$  must choose to remove  $x$  (or  $A$  wins on her next turn). After  $x$  is removed by  $B$ , consider the following figure.



We let  $A$  play in the two marked green cells. Then, regardless of what move  $B$  plays, one of the two choices of moves marked in red lets  $A$  win. Thus, we have described a winning strategy when  $k = 5$  for  $A$ .

## §6 JMO 2014/6, proposed by Titu Andreescu, Cosmin Pohoata

Let  $ABC$  be a triangle with incenter  $I$ , incircle  $\gamma$  and circumcircle  $\Gamma$ . Let  $M, N, P$  be the midpoints of  $\overline{BC}, \overline{CA}, \overline{AB}$  and let  $E, F$  be the tangency points of  $\gamma$  with  $\overline{CA}$  and  $\overline{AB}$ , respectively. Let  $U, V$  be the intersections of line  $EF$  with line  $MN$  and line  $MP$ , respectively, and let  $X$  be the midpoint of arc  $BAC$  of  $\Gamma$ .

- Prove that  $I$  lies on ray  $CV$ .
- Prove that line  $XI$  bisects  $\overline{UV}$ .

---

The fact that  $I = \overline{BU} \cap \overline{CV}$  is the so-called *Iran incenter lemma*, and is proved as Lemma 1.45 from my textbook.

As for (b), we note:

**Claim** — Line  $IX$  is a symmedian of  $\triangle IBC$ .

*Proof.* Recall that  $(BIC)$  has circumcenter coinciding with the antipode of  $X$  (by “Fact 5”). So this follows from the fact that  $\overline{XB}$  and  $\overline{XC}$  are tangent.  $\square$

Since  $BVUC$  is cyclic with diagonals intersecting at  $I$ , and  $IX$  is symmedian of  $\triangle IBC$ , it is median of  $\triangle IUV$ , as needed.

**Remark** (Alternate solution to (b) by Gunmay Handa). It’s well known that  $X$  is the midpoint of  $\overline{I_b I_c}$  (by considering the nine-point circle of the excentral triangle). However,  $\overline{UV} \parallel \overline{I_b I_c}$  and  $I = \overline{I_b U} \cap \overline{I_c V}$ , implying the result.