

JMO 2020 Solution Notes

COMPILED BY EVAN CHEN

September 30, 2020

This is an compilation of solutions for the 2020 JMO. Some of the solutions are my own work, but many are from the official solutions provided by the organizers (for which they hold any copyrights), and others were found on the Art of Problem Solving forums.

Corrections and comments are welcome!

Contents

0 Problems	2
1 JMO 2020/1, proposed by Milan Haiman	3
2 JMO 2020/2, proposed by Titu Andreescu and Waldemar Pompe	4
3 JMO 2020/3, proposed by Alex Zhai	5
4 JMO 2020/4, proposed by Milan Haiman	7
5 JMO 2020/5, proposed by Ankan Bhattacharya	9
6 JMO 2020/6, proposed by Ankan Bhattacharya	11

§0 Problems

- Let $n \geq 2$ be an integer. Carl has n books arranged on a bookshelf. Each book has a height and a width. No two books have the same height, and no two books have the same width.

Initially, the books are arranged in increasing order of height from left to right. In a *move*, Carl picks any two adjacent books where the left book is wider and shorter than the right book, and swaps their locations. Carl does this repeatedly until no further moves are possible.

Prove that regardless of how Carl makes his moves, he must stop after a finite number of moves, and when he does stop, the books are sorted in increasing order of width from left to right.

- Let ω be the incircle of a fixed equilateral triangle ABC . Let ℓ be a variable line that is tangent to ω and meets the interior of segments BC and CA at points P and Q , respectively. A point R is chosen such that $PR = PA$ and $QR = QB$. Find all possible locations of the point R , over all choices of ℓ .
- An empty $2020 \times 2020 \times 2020$ cube is given, and a 2020×2020 grid of square unit cells is drawn on each of its six faces. A *beam* is a $1 \times 1 \times 2020$ rectangular prism. Several beams are placed inside the cube subject to the following conditions:
 - The two 1×1 faces of each beam coincide with unit cells lying on opposite faces of the cube. (Hence, there are $3 \cdot 2020^2$ possible positions for a beam.)
 - No two beams have intersecting interiors.
 - The interiors of each of the four 1×2020 faces of each beam touch either a face of the cube or the interior of the face of another beam.

What is the smallest positive number of beams that can be placed to satisfy these conditions?

- Let $ABCD$ be a convex quadrilateral inscribed in a circle and satisfying

$$DA < AB = BC < CD.$$

Points E and F are chosen on sides CD and AB such that $\overline{BE} \perp \overline{AC}$ and $\overline{EF} \parallel \overline{BC}$. Prove that $FB = FD$.

- Suppose that $(a_1, b_1), (a_2, b_2), \dots, (a_{100}, b_{100})$ are distinct ordered pairs of non-negative integers. Let N denote the number of pairs of integers (i, j) satisfying $1 \leq i < j \leq 100$ and $|a_i b_j - a_j b_i| = 1$. Determine the largest possible value of N over all possible choices of the 100 ordered pairs.
- Let $n \geq 2$ be an integer. Let $P(x_1, x_2, \dots, x_n)$ be a nonconstant n -variable polynomial with real coefficients. Assuming that P vanishes whenever two of its arguments are equal, prove that P has at least $n!$ terms.

§1 JMO 2020/1, proposed by Milan Haiman

Let $n \geq 2$ be an integer. Carl has n books arranged on a bookshelf. Each book has a height and a width. No two books have the same height, and no two books have the same width.

Initially, the books are arranged in increasing order of height from left to right. In a *move*, Carl picks any two adjacent books where the left book is wider and shorter than the right book, and swaps their locations. Carl does this repeatedly until no further moves are possible.

Prove that regardless of how Carl makes his moves, he must stop after a finite number of moves, and when he does stop, the books are sorted in increasing order of width from left to right.

We say that a pair of books (A, B) is *height-inverted* if A is to the left of B and taller than B . Similarly define *width-inverted* pairs.


Note that every operation decreases the number of width-inverted pairs. This proves the procedure terminates, since the number of width-inverted pairs starts at $\binom{n}{2}$ and cannot increase indefinitely.

Now consider a situation where no more moves are possible. Assume for contradiction two consecutive books (A, B) are still width-inverted. Since the operation isn't possible anymore, they are also height-inverted. In particular, the operation could never have swapped A and B . But this contradicts the assumption there were no height-inverted pairs initially.

§2 JMO 2020/2, proposed by Titu Andreescu and Waldemar Pompe

Let ω be the incircle of a fixed equilateral triangle ABC . Let ℓ be a variable line that is tangent to ω and meets the interior of segments BC and CA at points P and Q , respectively. A point R is chosen such that $PR = PA$ and $QR = QB$. Find all possible locations of the point R , over all choices of ℓ .

Let r be the inradius. Let T be the tangency point of \overline{PQ} on arc \widehat{DE} of the incircle, which we consider varying. We define R_1 and R_2 to be the two intersections of the circle centered at P with radius PA , and the circle centered at Q with radius QB . We choose R_1 to lie on the opposite side of C as line PQ .



Claim — The point R_1 is the unique point on ray TI with $R_1I = 2r$.

Proof. Define S to be the point on ray TI with $SI = 2r$. Note that there is a homothety at I which maps $\triangle DTE$ to $\triangle ASB$, for some point S .

Note that since $TASD$ is an isosceles trapezoid, it follows $PA = PS$. Similarly, $QB = QS$. So it follows that $S = R_1$. \square

Since T can be any point on the open arc \widehat{DE} , it follows that the locus of R_1 is exactly the open 120° arc of \widehat{AB} of the circle centered at I with radius $2r$ (i.e. the circumcircle of ABC).

It remains to characterize R_2 . Since $TI = r$, $IR_1 = 2r$, it follows $TR_2 = 3r$ and $IR_2 = 4r$. Define A' on ray DI such that $A'I = 4r$, and B' on ray IE such that $B'I = 4r$. Then it follows, again by homothety, that the locus of R_2 is the 120° arc $\widehat{A'B'}$ of the circle centered at I with radius $4r$.

In conclusion, the locus of R is the two open 120° arcs we identified.

§3 JMO 2020/3, proposed by Alex Zhai


An empty $2020 \times 2020 \times 2020$ cube is given, and a 2020×2020 grid of square unit cells is drawn on each of its six faces. A *beam* is a $1 \times 1 \times 2020$ rectangular prism. Several beams are placed inside the cube subject to the following conditions:

- The two 1×1 faces of each beam coincide with unit cells lying on opposite faces of the cube. (Hence, there are $3 \cdot 2020^2$ possible positions for a beam.)
- No two beams have intersecting interiors.
- The interiors of each of the four 1×2020 faces of each beam touch either a face of the cube or the interior of the face of another beam.

What is the smallest positive number of beams that can be placed to satisfy these conditions?

Answer: 3030 beams.

Construction: We first give a construction with $3n/2$ beams for any $n \times n \times n$ box, where n is an even integer. Shown below is the construction for $n = 6$, which generalizes. (The left figure shows the cube in 3d; the right figure shows a direct view of the three visible faces.)



To be explicit, impose coordinate axes such that one corner of the cube is the origin. We specify a beam by two opposite corners. The $3n/2$ beams come in three directions, $n/2$ in each direction:

- $(0, 0, 0) \rightarrow (1, 1, n), (2, 2, 0) \rightarrow (3, 3, n), (4, 4, 0) \rightarrow (5, 5, n)$, and so on;
- $(1, 0, 0) \rightarrow (2, n, 1), (3, 0, 2) \rightarrow (4, n, 3), (5, 0, 4) \rightarrow (6, n, 5)$, and so on;
- $(0, 1, 1) \rightarrow (n, 2, 2), (0, 3, 3) \rightarrow (n, 4, 4), (0, 5, 5) \rightarrow (n, 6, 6)$, and so on.

This gives the figure we drew earlier and shows 3030 beams is possible.

Necessity: We now show at least $3n/2$ beams are necessary. Maintain coordinates, and call the beams x -beams, y -beams, z -beams according to which plane their long edges are perpendicular too. Let N_x, N_y, N_z be the number of these.

Claim — If $\min(N_x, N_y, N_z) = 0$, then at least n^2 beams are needed.

Proof. Assume WLOG that $N_z = 0$. Orient the cube so the z -plane touches the ground. Then each of the n layers of the cube (from top to bottom) must be completely filled, and so at least n^2 beams are necessary. \square

We henceforth assume $\min(N_x, N_y, N_z) > 0$.

Claim — If $N_z > 0$, then we have $N_x + N_y \geq n$.

Proof. Again orient the cube so the z -plane touches the ground. We see that for each of the n layers of the cube (from top to bottom), there is at least one x -beam or y -beam. (Pictorially, some of the x and y beams form a “staircase”.) This completes the proof. \square

Proceeding in a similar fashion, we arrive at the three relations

$$N_x + N_y \geq n$$

$$N_y + N_z \geq n$$

$$N_z + N_x \geq n.$$

Summing gives $N_x + N_y + N_z \geq 3n/2$ too.

Remark. The problem condition has the following “physics” interpretation. Imagine the cube is a metal box which is sturdy enough that all beams must remain orthogonal to the faces of the box (i.e. the beams cannot spin). Then the condition of the problem is exactly what is needed so that, if the box is shaken or rotated, the beams will not move.

Remark. Walter Stromquist points out that the number of constructions with 3030 beams is actually enormous: not dividing out by isometries, the number is $(2 \cdot 1010!)^3$.

§4 JMO 2020/4, proposed by Milan Haiman


Let $ABCD$ be a convex quadrilateral inscribed in a circle and satisfying

$$DA < AB = BC < CD.$$

Points E and F are chosen on sides CD and AB such that $\overline{BE} \perp \overline{AC}$ and $\overline{EF} \parallel \overline{BC}$. Prove that $FB = FD$.

We present three approaches. We note that in the second two approaches, the result remains valid even if $AB \neq BC$, as long as E is replaced by the point on \overline{AC} satisfying $EA = EC$. So the result is actually somewhat more general.


First solution by inscribed angle theorem Since $\overline{EF} \parallel \overline{BC}$ we may set $\theta = \angle FEB = \angle CBE = \angle EBF$. This already implies $FE = FB$, so we will in fact prove that F is the circumcenter of $\triangle BED$.



Note that $\angle BDC = \angle BAC = 90^\circ - \theta$. However, $\angle BFE = 180^\circ - 2\theta$. So by the inscribed angle theorem, D lies on the circle centered at F with radius $FE = FB$, as desired.

Remark. Another approach to the given problem is to show that B is the D -excenter of $\triangle DAE$, and F is the arc midpoint of \widehat{DAE} of the circumcircle of $\triangle DAE$. In my opinion, this approach is much clumsier.

Second general solution by angle chasing By Reim's theorem, $AFED$ is cyclic.




Hence

$$\begin{aligned} \angle FDB &= \angle FDC - \angle BDC = \angle FAE - \angle FAC \\ &= \angle CAE = \angle ECA = \angle DCA = \angle DBA = \angle DBF \end{aligned}$$

as desired.

Third general solution by Pascal Extend rays AE and DF to meet the circumcircle again at G and H . By Pascal's theorem on $HDCBAG$, it follows that E, F , and $GH \cap BC$ are collinear, which means that $\overline{EF} \parallel \overline{GH} \parallel \overline{BC}$.



Since $EA = EC$, it follows $DAGC$ is isosceles trapezoid. But also $GHBC$ is an isosceles trapezoid. Thus $m\widehat{DA} = m\widehat{GC} = m\widehat{BH}$, so $DAHB$ is an isosceles trapezoid. Thus $FD = FB$.

Remark. Addicts of projective geometry can use Pascal on $DBCAHG$ to finish rather than noting the equal arcs.

§5 JMO 2020/5, proposed by Ankan Bhattacharya

Suppose that $(a_1, b_1), (a_2, b_2), \dots, (a_{100}, b_{100})$ are distinct ordered pairs of nonnegative integers. Let N denote the number of pairs of integers (i, j) satisfying $1 \leq i < j \leq 100$ and $|a_i b_j - a_j b_i| = 1$. Determine the largest possible value of N over all possible choices of the 100 ordered pairs.


The answer is 197. In general, if 100 is replaced by $n \geq 2$ the answer is $2n - 3$.

The idea is that if we let $P_i = (a_i, b_i)$ be a point in the coordinate plane, and let $O = (0, 0)$ then we wish to maximize the number of triangles $\triangle OP_i P_j$ which have area $1/2$. Call such a triangle *good*.

Construction of 197 points: It suffices to use the points $(1, 0), (1, 1), (2, 1), (3, 1), \dots, (99, 1)$ as shown. Notice that:

- There are 98 good triangles with vertices $(0, 0), (k, 1)$ and $(k+1, 1)$ for $k = 1, \dots, 98$.
- There are 99 good triangles with vertices $(0, 0), (1, 0)$ and $(k, 1)$ for $k = 1, \dots, 99$.

This is a total of $98 + 99 = 197$ triangles.




Proof that 197 points is optimal: We proceed by induction on n to show the bound of $2n - 3$. The base case $n = 2$ is evident.

For the inductive step, suppose (without loss of generality) that the point $P = P_n = (a, b)$ is the farthest away from the point O among all points.

Claim — This farthest point $P = P_n$ is part of at most two good triangles.

Proof. We must have $\gcd(a, b) = 1$ for P to be in any good triangles at all, since otherwise any divisor of $\gcd(a, b)$ also divides $2[OPQ]$. Now, we consider the locus of all points Q for which $[OPQ] = 1/2$. It consists of two parallel lines passing with slope OP , as shown.



Since $\gcd(a, b) = 1$, see that only two lattice points on this locus actually lie inside the quarter-circle centered at O with radius OP . Indeed if one of the points is (u, v) then the others on the line are $(u \pm a, v \pm b)$ where the signs match. This proves the claim. \square

This claim allows us to complete the induction by simply deleting P_n .

§6 JMO 2020/6, proposed by Ankan Bhattacharya

Let $n \geq 2$ be an integer. Let $P(x_1, x_2, \dots, x_n)$ be a nonconstant n -variable polynomial with real coefficients. Assuming that P vanishes whenever two of its arguments are equal, prove that P has at least $n!$ terms.

We present Ankan's original solution. Begin with the following observation:

Claim — Let $1 \leq i < j \leq n$. There is no term of P which omits both x_i and x_j .

Proof. Note that P ought to become identically zero if we set $x_i = x_j = 0$, since it is zero for any choice of the remaining $n - 2$ variables, and the base field \mathbb{R} is infinite. \square

Remark (Technical warning for experts). The fact we used is not true if \mathbb{R} is replaced by a field with finitely many elements, such as \mathbb{F}_p , even with one variable. For example the one-variable polynomial $X^p - X$ vanishes on every element of \mathbb{F}_p , by Fermat's little theorem.

We proceed by induction on $n \geq 2$ with the base case $n = 2$ being clear. Assume WLOG P is not divisible by any of x_1, \dots, x_n , since otherwise we may simply divide out this factor. Now for the inductive step, note that

- The polynomial $P(0, x_2, x_3, \dots, x_n)$ obviously satisfies the inductive hypothesis and is not identically zero since $x_1 \nmid P$, so it has at least $(n - 1)!$ terms.
- Similarly, $P(x_1, 0, x_3, \dots, x_n)$ also has at least $(n - 1)!$ terms.
- Similarly, $P(x_1, x_2, 0, \dots, x_n)$ also has at least $(n - 1)!$ terms.
- ... and so on.

By the claim, all the terms obtained in this way came from different terms of the original polynomial P . Therefore, P itself has at least $n \cdot (n - 1)! = n!$ terms.

Remark. Equality is achieved by the Vandermonde polynomial $P = \prod_{1 \leq i < j \leq n} (x_i - x_j)$.