

USAMO 2001 Solution Notes

COMPILED BY EVAN CHEN

January 22, 2020

This is an compilation of solutions for the 2001 USAMO. Some of the solutions are my own work, but many are from the official solutions provided by the organizers (for which they hold any copyrights), and others were found on the Art of Problem Solving forums.

Corrections and comments are welcome!

Contents

0 Problems	2
1 USAMO 2001/1	3
2 USAMO 2001/2	5
3 USAMO 2001/3	6
4 USAMO 2001/4	8
5 USAMO 2001/5	9
6 USAMO 2001/6	10

§0 Problems

1. Each of eight boxes contains six balls. Each ball has been colored with one of n colors, such that no two balls in the same box are the same color, and no two colors occur together in more than one box. Find with proof the smallest possible n .
2. Let ABC be a triangle and let ω be its incircle. Denote by D_1 and E_1 the points where ω is tangent to sides BC and AC , respectively. Denote by D_2 and E_2 the points on sides BC and AC , respectively, such that $CD_2 = BD_1$ and $CE_2 = AE_1$, and denote by P the point of intersection of segments AD_2 and BE_2 . Circle ω intersects segment AD_2 at two points, the closer of which to the vertex A is denoted by Q . Prove that $AQ = D_2P$.
3. Let a, b, c be nonnegative real numbers such that $a^2 + b^2 + c^2 + abc = 4$. Show that

$$0 \leq ab + bc + ca - abc \leq 2.$$

4. Let ABC be a triangle and P any point such that PA, PB, PC are the sides of an obtuse triangle, with PA the longest side. Prove that $\angle BAC$ is acute.
5. Let $S \subseteq \mathbb{Z}$ be such that:
 - (a) there exist $a, b \in S$ with $\gcd(a, b) = \gcd(a - 2, b - 2) = 1$;
 - (b) if x and y are element of S (possibly equal), then $x^2 - y$ also belongs to S .Prove that $S = \mathbb{Z}$.
6. Each point in the plane is assigned a real number. Suppose that for any nondegenerate triangle, the number at its incenter is the arithmetic mean of the three numbers at its vertices. Prove that all points in the plane are equal to each other.

§1 USAMO 2001/1

Each of eight boxes contains six balls. Each ball has been colored with one of n colors, such that no two balls in the same box are the same color, and no two colors occur together in more than one box. Find with proof the smallest possible n .

The answer is $n = 23$. Shown below is a construction using that many colors, which we call $\{1, 2, \dots, 15, a, \dots, f, X, Y\}$.

$$\begin{bmatrix} X & X & X & 1 & 2 & 3 & 4 & 5 \\ 1 & 6 & 11 & 6 & 7 & 8 & 9 & 10 \\ 2 & 7 & 12 & 11 & 12 & 13 & 14 & 15 \\ 3 & 8 & 13 & Y & Y & Y & a & b \\ 4 & 9 & 14 & a & c & e & c & d \\ 5 & 10 & 15 & b & d & f & e & f \end{bmatrix}$$

We present now two proofs that $n = 23$ is best possible. I think the first is more motivated — it will actually show us how we could come up with the example above.

First solution (hands-on) We say a color x is *overrated* if it is used at least three times. First we make the following smoothing argument.

Claim — Suppose some box contains a ball of overrated color x plus a ball of color y used only once. Then we can change one ball of color x to color y while preserving all the conditions.

Proof. Obvious. (Though the color x could cease to be overrated after this operation.) \square

By applying this operation as many times as possible, we arrive at a situation in which whenever we have a box with an overrated color, the other colors in the box are used twice or more.

Assume now $n \leq 23$ and the assumption; we will show the equality case must of the form we gave. Since there are a total of 48 balls, at least two colors are overrated. Let X be an overrated color and take three boxes where it appears. Then there are 15 more distinct colors, say $\{1, \dots, 15\}$ lying in those boxes. Each of them must appear at least once more, so we arrive at the situation

$$\begin{bmatrix} X & X & X & 1 & 2 & 3 & 4 & 5 \\ 1 & 6 & 11 & 6 & 7 & 8 & 9 & 10 \\ 2 & 7 & 12 & 11 & 12 & 13 & 14 & 15 \\ 3 & 8 & 13 & & & & & \\ 4 & 9 & 14 & & & & & \\ 5 & 10 & 15 & & & & & \end{bmatrix}$$

up to harmless permutation of the color names. Now, note that none of these 15 colors can reappear. So it remains to fill up the last five boxes.

Now, there is at least one more overrated color, distinct from any we have seen; call it Y . In the three boxes Y appears in, there must be six new colors, and this gives the lower bound $n \geq 1 + 15 + 1 + 6 = 23$ which we sought, with equality occurring as we saw above.

Remark (Partial progresses). The fact that $\binom{16}{2} = 120 = 8\binom{6}{2}$ (suggesting the bound $n \geq 16$) is misleading and not that helpful.

There is a simple argument showing that n should be much larger than 16. Imagine opening the boxes in any order. The first box must contain six new colors. The second box must contain five new colors, and so on; thus $n \geq 6 + 5 + 4 + 3 + 2 + 1 = 21$. This is sharp for seven boxes, as the example below shows.

$$\begin{bmatrix} 1 & 1 & 2 & 3 & 4 & 5 & 6 \\ 2 & 7 & 7 & 8 & 9 & 10 & 11 \\ 3 & 8 & 12 & 12 & 13 & 14 & 15 \\ 4 & 9 & 13 & 16 & 16 & 17 & 18 \\ 5 & 10 & 14 & 17 & 19 & 19 & 20 \\ 6 & 11 & 15 & 18 & 20 & 21 & 21 \end{bmatrix}$$

However, one cannot add an eight box, suggesting the answer should be a little larger than 21. One possible eight box is $\{1, 12, 19, a, b, c\}$ which gives $n \leq 24$; but the true answer is a little trickier.

Second solution (slick) Here is a short proof from the official solutions of the bound. Consider the 8×6 grid of colors as before. For each ball b , count the number of times n_b its color is used, and write the fraction $\frac{1}{n_b}$.

On the one hand, we should have

$$n = \sum_{\text{all 48 balls } b} \frac{1}{n_b}.$$

On the other hand, for any given box B , we have $\sum_{b \in B} (n_b - 1) \leq 7$, as among the other seven boxes at most one color from B appears. Therefore, $\sum_{b \in B} n_b \leq 13$, and a smoothing argument this implies

$$\sum_{b \in B} \frac{1}{n_b} \geq \frac{1}{3} \cdot 1 + \frac{1}{2} \cdot 5 = \frac{17}{6}.$$

Thus, $n \geq 8 \cdot \frac{17}{6} = 22.66\dots$, so $n \geq 23$.

§2 USAMO 2001/2

Let ABC be a triangle and let ω be its incircle. Denote by D_1 and E_1 the points where ω is tangent to sides BC and AC , respectively. Denote by D_2 and E_2 the points on sides BC and AC , respectively, such that $CD_2 = BD_1$ and $CE_2 = AE_1$, and denote by P the point of intersection of segments AD_2 and BE_2 . Circle ω intersects segment AD_2 at two points, the closer of which to the vertex A is denoted by Q . Prove that $AQ = D_2P$.

We have that P is the Nagel point

$$P = (s - a : s - b : s - c).$$

Therefore,

$$\frac{PD_2}{AD_2} = \frac{s - a}{(s - a) + (s - b) + (s - c)} = \frac{s - a}{s}.$$

Meanwhile, Q is the antipode of D_1 . The classical homothety at A mapping Q to D_1 (by mapping the incircle to the A -excircle) has ratio $\frac{s-a}{s}$ as well (by considering the length of the tangents from A), so we are done.

§3 USAMO 2001/3

Let a, b, c be nonnegative real numbers such that $a^2 + b^2 + c^2 + abc = 4$. Show that

$$0 \leq ab + bc + ca - abc \leq 2.$$

The left-hand side of the inequality is trivial; just note that $\min\{a, b, c\} \leq 1$. Hence, we focus on the right side. We use Lagrange Multipliers.

Define

$$U = \{(a, b, c) \mid a, b, c > 0 \text{ and } a^2 + b^2 + c^2 < 1000\}.$$

This is an intersection of open sets, so it is open. Its closure is

$$\bar{U} = \{(a, b, c) \mid a, b, c \geq 0 \text{ and } a^2 + b^2 + c^2 \leq 1000\}.$$

Hence the constraint set

$$\bar{S} = \{\mathbf{x} \in \bar{U} : g(\bar{\mathbf{x}}) = 3\}$$

is compact, where $g(a, b, c) = a^2 + b^2 + c^2 + abc$.

Define

$$f(a, b, c) = a^2 + b^2 + c^2 + ab + bc + ca.$$

It's equivalent to show that $f \leq 6$ subject to g . Over \bar{S} , it must achieve a global maximum. Now we consider two cases.

If \mathbf{x} lies on the boundary, that means one of the components is zero (since $a^2 + b^2 + c^2 = 1000$ is clearly impossible). WLOG $c = 0$, then we wish to show $a^2 + b^2 + ab \leq 6$ for $a^2 + b^2 = 4$, which is trivial.

Now for the interior U , we may use the method of Lagrange Multipliers. Consider a local maximum $\mathbf{x} \in U$. Compute

$$\nabla f = \langle 2a + b + c, 2b + c + a, 2c + a + b \rangle$$

and

$$\nabla g = \langle 2a + bc, 2b + ca, 2c + ab \rangle.$$

Of course, $\nabla g \neq \mathbf{0}$ everywhere, so introducing our multiplier yields

$$\langle 2a + b + c, a + 2b + c, a + b + 2c \rangle = \lambda \langle 2a + bc, 2b + ca, 2c + ab \rangle.$$

Note that $\lambda \neq 0$ since $a, b, c > 0$. Subtracting $2a + b + c = \lambda(2a + bc)$ from $a + 2b + c = \lambda(2b + ca)$ implies that

$$(a - b)([2\lambda - 1] - \lambda c) = 0.$$

We can derive similar equations for the others. Hence, we have three cases.

1. If $a = b = c$, then $a = b = c = 1$, and this satisfies $f(1, 1, 1) \leq 6$.
2. If a, b, c are pairwise distinct, then we derive $a = b = c = 2 - \lambda^{-1}$, contradiction.
3. Now suppose that $a = b \neq c$. That means $\lambda = \frac{1}{2-a}$ (of course our conditions force $c < 2$). Now

$$2a + 2c = a + b + 2c = \lambda(2c + ab) = \frac{1}{2-a}(2c + a^2)$$

which implies

$$4a + 4c - 2a^2 - 2ac = 2c + a^2$$

meaning (with the additional note that $a \neq 1$) we have

$$c = \frac{3a^2 - 4a}{2 - 2a}.$$

Note that at this point, $c > 0$ forces $1 < a < \frac{4}{3}$.

The constraint $a^2 + b^2 + c^2 + abc = 4 \iff c^2 + a^2c + (2a^2 - 4) = 0$ now gives

$$(3a^2 - 4a)^2 + a^2(3a^2 - 4a)(2 - 2a) + (2a^2 - 4)(2 - 2a)^2 = 0.$$

Before expanding this, it is prudent to see if it has any rational roots.

A quick inspection finds that $a = 2$ is such a root (precisely, $16 - 32 + 16 = 0$; or even $(a, b, c) = (2, 2, -2)$). Now, we can expand and try to factor:

$$\begin{aligned} 0 &= -6a^5 + 31a^4 - 48a^3 + 8a^2 + 32a - 16 \\ &= (a - 2)(-6a^4 + 19a^3 - 10a^2 - 12a + 8) \\ &= (a - 2)^2(-6a^3 + 7a^2 + 4a - 4) \\ &= (a - 2)^2(2 - 3a)(2a^2 - a - 2). \end{aligned}$$

The only root a in the interval $(1, \frac{4}{3})$ is $a = \frac{1}{4}(1 + \sqrt{17})$. To finish, write

$$c = \frac{a(3a - 4)}{2 - 2a} = \frac{1}{8}(7 - \sqrt{17})$$

and

$$f(a, b, c) = 3a^2 + 2ac + c^2 = \frac{1}{32}(121 + 17\sqrt{17}).$$

This is the last critical point, so we're done once we check this is less than 6. This follows from the inequality $17^3 < (6 \cdot 32 - 121)^2$; in fact, we actually have

$$\frac{1}{32}(121 + 17\sqrt{17}) \approx 5.97165.$$

This completes the solution.

Remark. Equality holds for the upper bound if $(a, b, c) = (1, 1, 1)$ or $(a, b, c) = (\sqrt{2}, \sqrt{2}, 0)$ and permutations. The lower bound is achieved if $(a, b, c) = (2, 0, 0)$ and permutations.

§4 USAMO 2001/4

Let ABC be a triangle and P any point such that PA, PB, PC are the sides of an obtuse triangle, with PA the longest side. Prove that $\angle BAC$ is acute.

Using Ptolemy's inequality and Cauchy-Schwarz,

$$\begin{aligned} PA \cdot BC &\leq PB \cdot AC + PC \cdot AB \\ &\leq \sqrt{(PB^2 + PC^2)(AB^2 + AC^2)} \\ &< \sqrt{PA^2 \cdot (AB^2 + AC^2)^2} = PA \cdot \sqrt{AB^2 + AC^2} \end{aligned}$$

meaning $BC^2 < AB^2 + AC^2$, so $\angle BAC$ is acute.

§5 USAMO 2001/5

Let $S \subseteq \mathbb{Z}$ be such that:

- (a) there exist $a, b \in S$ with $\gcd(a, b) = \gcd(a - 2, b - 2) = 1$;
- (b) if x and y are element of S (possibly equal), then $x^2 - y$ also belongs to S .

Prove that $S = \mathbb{Z}$.

Call an integer $d > 0$ *shifty* if $S = S + d$ (meaning S is invariant under shifting by d). First, note that if $u, v \in S$, then for any $x \in S$,

$$v^2 - (u^2 - x) = (v^2 - u^2) + x \in S.$$

Since we can easily check that $|S| > 1$ and $S \neq \{n, -n\}$ we conclude exists a shifty integer.

We claim 1 is shifty, which implies the problem. Assume for contradiction not that 1 is not shifty. Then for GCD reasons the set of shifty integers must be $d\mathbb{Z}$ for some $d \geq 2$.

Claim — We have $S \subseteq \{x : x^2 \equiv m \pmod{d}\}$ for some fixed m .

Proof. Otherwise if we take any $p, q \in S$ with distinct squares modulo d , then $q^2 - p^2 \not\equiv 0 \pmod{d}$ is shifty, which is impossible. \square

Now take $a, b \in S$ as in (a). In that case we need to have

$$a^2 \equiv b^2 \equiv (a^2 - a)^2 \equiv (b^2 - b)^2 \pmod{d}.$$

Passing to a prime $p \mid d$, we have the following:

- Since $a^2 \equiv (a^2 - a)^2 \pmod{p}$ or equivalently $a^3(a - 2) \equiv 0 \pmod{p}$, either $a \equiv 0 \pmod{p}$ or $a \equiv 2 \pmod{p}$.
- Similarly, either $b \equiv 0 \pmod{p}$ or $b \equiv 2 \pmod{p}$.
- Since $a^2 \equiv b^2 \pmod{p}$, or $a \equiv \pm b \pmod{p}$, we find either $a \equiv b \equiv 0 \pmod{p}$ or $a \equiv b \equiv 2 \pmod{p}$ (even if $p = 2$).

This is a contradiction.

Remark. The condition (a) cannot be dropped, since otherwise we may take $S = \{2 \pmod{p}\}$ or $S = \{0 \pmod{p}\}$, say.

§6 USAMO 2001/6

Each point in the plane is assigned a real number. Suppose that for any nondegenerate triangle, the number at its incenter is the arithmetic mean of the three numbers at its vertices. Prove that all points in the plane are equal to each other.

First, we claim that in an isosceles trapezoid $ABCD$ we have $a + c = b + d$. Indeed, suppose WLOG that rays BA and CD meet at X . Then triangles XAC and XBD share an incircle, proving the claim.

Now, given any two points A and B , construct regular pentagon $ABCDE$. We have $a + c = b + d = c + e = d + a = e + b$, so $a = b = c = d = e$.