

USAMO 2009 Solution Notes

EVAN CHEN 《陳誼廷》

11 December 2023

This is a compilation of solutions for the 2009 USAMO. Some of the solutions are my own work, but many are from the official solutions provided by the organizers (for which they hold any copyrights), and others were found by users on the Art of Problem Solving forums.

These notes will tend to be a bit more advanced and terse than the “official” solutions from the organizers. In particular, if a theorem or technique is not known to beginners but is still considered “standard”, then I often prefer to use this theory anyways, rather than try to work around or conceal it. For example, in geometry problems I typically use directed angles without further comment, rather than awkwardly work around configuration issues. Similarly, sentences like “let \mathbb{R} denote the set of real numbers” are typically omitted entirely.


Corrections and comments are welcome!

Contents

0 Problems	2
1 Solutions to Day 1	3
1.1 USAMO 2009/1, proposed by Ian Le	3
1.2 USAMO 2009/2, proposed by Kiran Kedlaya, Tewodos Amdeberhan . . .	4
1.3 USAMO 2009/3, proposed by Sam Vandervelde	5
2 Solutions to Day 2	7
2.1 USAMO 2009/4, proposed by Titu Andreescu	7
2.2 USAMO 2009/5, proposed by Zuming Feng	8
2.3 USAMO 2009/6, proposed by Gabriel Carroll	9

§0 Problems

- Given circles ω_1 and ω_2 intersecting at points X and Y , let ℓ_1 be a line through the center of ω_1 intersecting ω_2 at points P and Q and let ℓ_2 be a line through the center of ω_2 intersecting ω_1 at points R and S . Prove that if P , Q , R , and S lie on a circle then the center of this circle lies on line XY .
- Let n be a positive integer. Determine the size of the largest subset of $\{-n, -n+1, \dots, n-1, n\}$ which does not contain three elements a , b , c (not necessarily distinct) satisfying $a + b + c = 0$.
- We define a *chessboard polygon* to be a simple polygon whose sides are situated along lines of the form $x = a$ or $y = b$, where a and b are integers. These lines divide the interior into unit squares, which are shaded alternately grey and white so that adjacent squares have different colors. To tile a chessboard polygon by dominoes is to exactly cover the polygon by non-overlapping 1×2 rectangles. Finally, a *tasteful tiling* is one which avoids the two configurations of dominoes and colors shown on the left below. Two tilings of a 3×4 rectangle are shown; the first one is tasteful, while the second is not, due to the vertical dominoes in the upper right corner.



Prove that (a) if a chessboard polygon can be tiled by dominoes, then it can be done so tastefully, and (b) such a tasteful tiling is unique.

- For $n \geq 2$, let a_1, a_2, \dots, a_n be positive real numbers such that

$$(a_1 + a_2 + \dots + a_n) \left(\frac{1}{a_1} + \frac{1}{a_2} + \dots + \frac{1}{a_n} \right) \leq \left(n + \frac{1}{2} \right)^2.$$

Prove that $\max(a_1, \dots, a_n) \leq 4 \min(a_1, \dots, a_n)$.

- Trapezoid $ABCD$, with $\overline{AB} \parallel \overline{CD}$, is inscribed in circle ω and point G lies inside triangle BCD . Rays AG and BG meet ω again at points P and Q , respectively. Let the line through G parallel to \overline{AB} intersect \overline{BD} and \overline{BC} at points R and S , respectively. Prove that quadrilateral $PQRS$ is cyclic if and only if \overline{BG} bisects $\angle CBD$.
- Let s_1, s_2, s_3, \dots be an infinite, nonconstant sequence of rational numbers, meaning it is not the case that $s_1 = s_2 = s_3 = \dots$. Suppose that t_1, t_2, t_3, \dots is also an infinite, nonconstant sequence of rational numbers with the property that $(s_i - s_j)(t_i - t_j)$ is an integer for all i and j . Prove that there exists a rational number r such that $(s_i - s_j)r$ and $(t_i - t_j)/r$ are integers for all i and j .

§1 Solutions to Day 1


§1.1 USAMO 2009/1, proposed by Ian Le

Available online at <https://aops.com/community/p1485133>.

Problem statement

Given circles ω_1 and ω_2 intersecting at points X and Y , let ℓ_1 be a line through the center of ω_1 intersecting ω_2 at points P and Q and let ℓ_2 be a line through the center of ω_2 intersecting ω_1 at points R and S . Prove that if P , Q , R , and S lie on a circle then the center of this circle lies on line XY .

Let r_1, r_2, r_3 denote the circumradii of ω_1, ω_2 , and ω_3 , respectively.



We wish to show that O_3 lies on the radical axis of ω_1 and ω_2 . Let us encode the conditions using power of a point. Because O_1 is on the radical axis of ω_2 and ω_3 ,

$$\begin{aligned} \text{Pow}_{\omega_2}(O_1) &= \text{Pow}_{\omega_3}(O_1) \\ \implies O_1O_2^2 - r_2^2 &= O_1O_3^2 - r_3^2. \end{aligned}$$

Similarly, because O_2 is on the radical axis of ω_1 and ω_3 , we have

$$\begin{aligned} \text{Pow}_{\omega_1}(O_2) &= \text{Pow}_{\omega_3}(O_2) \\ \implies O_1O_2^2 - r_1^2 &= O_2O_3^2 - r_3^2. \end{aligned}$$

Subtracting the two gives

$$\begin{aligned} (O_1O_2^2 - r_2^2) - (O_1O_2^2 - r_1^2) &= (O_1O_3^2 - r_3^2) - (O_2O_3^2 - r_3^2) \\ \implies r_1^2 - r_2^2 &= O_1O_3^2 - O_2O_3^2 \\ \implies O_2O_3^2 - r_2^2 &= O_1O_3^2 - r_1^2 \\ \implies \text{Pow}_{\omega_2}(O_3) &= \text{Pow}_{\omega_1}(O_3) \end{aligned}$$

as desired.

§1.2 USAMO 2009/2, proposed by Kiran Kedlaya, Tewodos Amdeberhan

Available online at <https://aops.com/community/p1485139>.

Problem statement

Let n be a positive integer. Determine the size of the largest subset of $\{-n, -n+1, \dots, n-1, n\}$ which does not contain three elements a, b, c (not necessarily distinct) satisfying $a + b + c = 0$.

The answer is n with n even and $n+1$ with n odd; the construction is to take all odd numbers.

To prove this is maximal, it suffices to show it for n even; we do so by induction on even $n \geq 2$ with the base case being trivial. Letting A be the subset, we consider three cases:

- (i) If $|A \cap \{-n, -n+1, n-1, n\}| \leq 2$, then by the hypothesis for $n-2$ we are done.
- (ii) If both $n \in A$ and $-n \in A$, then there can be at most $n-2$ elements in $A \setminus \{\pm n\}$, one from each of the pairs $(1, n-1), (2, n-2), \dots$ and their negations.
- (iii) If $n, n-1, -n+1 \in A$ and $-n \notin A$, and at most $n-3$ more can be added, one from each of $(1, n-2), (2, n-3), \dots$ and $(-2, -n+2), (-3, -n+3), \dots$ (In particular $-1 \notin A$. Analogous case for $-A$ if $n \notin A$.)

Thus in all cases, $|A| \leq n$ as needed.

Remark. A few examples of equality cases:

- All odd numbers
- All numbers with absolute value at least $n/2$
- For n even, the set $\{1, 2, \dots, n\}$


This list isn't exhaustive e.g. for $n=4$, the set $\{-3, 2, 3, 4\}$ achieves the optimum.

§1.3 USAMO 2009/3, proposed by Sam Vandervelde

Available online at <https://aops.com/community/p1485321>.

Problem statement

We define a *chessboard polygon* to be a simple polygon whose sides are situated along lines of the form $x = a$ or $y = b$, where a and b are integers. These lines divide the interior into unit squares, which are shaded alternately grey and white so that adjacent squares have different colors. To tile a chessboard polygon by dominoes is to exactly cover the polygon by non-overlapping 1×2 rectangles. Finally, a *tasteful tiling* is one which avoids the two configurations of dominoes and colors shown on the left below. Two tilings of a 3×4 rectangle are shown; the first one is tasteful, while the second is not, due to the vertical dominoes in the upper right corner.




Prove that (a) if a chessboard polygon can be tiled by dominoes, then it can be done so tastefully, and (b) such a tasteful tiling is unique.

¶ **Proof of (a).** This is easier, and by induction. Let \mathcal{P} denote the chessboard polygon which can be tiled by dominoes.

Consider a *lower-left* square s of the polygon, and WLOG is it white (other case similar). Then we have two cases:

- If there exists a domino tiling of \mathcal{P} where s is covered by a vertical domino, then delete this domino and apply induction on the rest of \mathcal{P} . This additional domino will not cause any distasteful tilings.
- Otherwise, assume s is covered by a horizontal domino in *every* tiling. Again delete this domino and apply induction on the rest of \mathcal{P} . The resulting tasteful tiling should not have another horizontal domino adjacent to the one covering s , because otherwise we could have replaced that 2×2 square with two vertical dominoes to arrive in the first case. So this additional domino will not cause any distasteful tilings.

Remark. The second case can actually arise, for example in the following picture.



Thus one cannot just try to cover s with a vertical domino and claim the rest of \mathcal{P} is tile-able. So the induction is not as easy as one might hope.

One can phrase the solution algorithmically too, in the following way: any time we see a


distasteful tiling, we rotate it to avoid the bad pattern. The bottom-left corner eventually becomes stable, and an induction shows the termination of the algorithm.

¶ Proof of (b). We now turn to proving uniqueness. Suppose for contradiction there are two distinct tasteful tilings. Overlaying the two tilings on top of each other induces several *cycles* of overlapping dominoes at positions where the tilings differ.

Henceforth, it will be convenient to work with the lattice \mathbb{Z}^2 , treating the squares as black/white points, and we do so. Let γ be any such cycle and let s denote a lower left point, and again WLOG it is black. Orient γ counterclockwise henceforth. Restrict attention to the lattice polygon \mathcal{Q} enclosed by γ (we consider points of γ as part of \mathcal{Q}).

In one of the two tilings of (lattice points of) \mathcal{Q} , the point s will be covered by a horizontal domino; in the other tiling s will be covered by a vertical domino. From now on we will focus only on the latter one. Observe that we now have a set of dominoes along γ , such that γ points from the white point to the black point within each domino.

Now impose coordinates so that $s = (0, 0)$. Consider the stair-case sequence of points $p_0 = s = (0, 0)$, $p_1 = (1, 0)$, $p_2 = (1, 1)$, $p_3 = (2, 1)$, and so on. By hypothesis, p_0 is covered by a vertical domino. Then p_1 must be covered by a horizontal domino, to avoid a distasteful tiling. Then if p_2 is in \mathcal{Q} , then it must be covered by a vertical domino to avoid a distasteful tiling, and so on. We may repeat this argument as long the points p_i lie inside \mathcal{Q} . (See figure below; the staircase sequence is highlighted by red halos.)



The curve γ by definition should cross $y = x - 1$ at the point $b = (1, 0)$. Let a denote the first point of this sequence after p_1 for which γ crosses $y = x - 1$ again.

Now a is tiled by a vertical domino whose black point is to the right of ℓ . But the line segment ℓ cuts \mathcal{Q} into two parts, and the orientation of γ has this path also entering from the right. This contradicts the fact that the orientation of γ points only from white to black within dominoes. This contradiction completes the proof.

Remark. Note the problem is false if you allow holes (consider a 3×3 with the middle square deleted).

§2 Solutions to Day 2

§2.1 USAMO 2009/4, proposed by Titu Andreescu

Available online at <https://aops.com/community/p1485147>.

Problem statement

For $n \geq 2$, let a_1, a_2, \dots, a_n be positive real numbers such that

$$(a_1 + a_2 + \dots + a_n) \left(\frac{1}{a_1} + \frac{1}{a_2} + \dots + \frac{1}{a_n} \right) \leq \left(n + \frac{1}{2} \right)^2.$$

Prove that $\max(a_1, \dots, a_n) \leq 4 \min(a_1, \dots, a_n)$.

Assume a_1 is the largest and a_2 is the smallest. Let $M = a_1/a_2$. We wish to show $M \leq 4$.

In left-hand side of given, write as $a_2 + a_1 + \dots + a_n$. By Cauchy Schwarz, one obtains

$$\begin{aligned} \left(n + \frac{1}{2} \right)^2 &\geq (a_2 + a_1 + a_3 + \dots + a_n) \left(\frac{1}{a_1} + \frac{1}{a_2} + \frac{1}{a_3} + \dots + \frac{1}{a_n} \right) \\ &\geq \left(\sqrt{\frac{a_2}{a_1}} + \sqrt{\frac{a_1}{a_2}} + 1 + \dots + 1 \right)^2 \\ &\geq \left(\sqrt{1/M} + \sqrt{M} + (n-2) \right)^2. \end{aligned}$$

Expanding and solving for M gives $1/4 \leq M \leq 4$ as needed.


§2.2 USAMO 2009/5, proposed by Zuming Feng

Available online at <https://aops.com/community/p1485200>.

Problem statement

Trapezoid $ABCD$, with $\overline{AB} \parallel \overline{CD}$, is inscribed in circle ω and point G lies inside triangle BCD . Rays AG and BG meet ω again at points P and Q , respectively. Let the line through G parallel to \overline{AB} intersect \overline{BD} and \overline{BC} at points R and S , respectively. Prove that quadrilateral $PQRS$ is cyclic if and only if \overline{BG} bisects $\angle CBD$.

Perform an inversion around B with arbitrary radius, and denote the inverse of a point Z with Z^* .



After inversion, we obtain a cyclic quadrilateral $BS^*G^*R^*$ and points C^*, D^* on $\overline{BS^*}$, $\overline{BR^*}$, such that (BC^*D^*) is tangent to $(BS^*G^*R^*)$ — in other words, so that $\overline{C^*D^*}$ is parallel to $\overline{S^*R^*}$. Point A^* lies on line $\overline{C^*D^*}$ so that $\overline{A^*B}$ is tangent to $(BS^*G^*R^*)$. Points P^* and Q^* are the intersections of (A^*BG^*) and $\overline{BG^*}$ with line $\overline{C^*D^*}$.

Observe that $P^*Q^*R^*S^*$ is a trapezoid, so it is cyclic if and only if it is isosceles.

Let X be the second intersection of line $\overline{G^*P^*}$ with (BS^*R^*) . Because

$$\angle Q^*P^*G^* = \angle A^*BG^* = \angle BXG^*$$

we find that BXS^*R^* is an isosceles trapezoid.

If G^* is indeed the midpoint of the arc then everything is clear by symmetry now. Conversely, if $P^*R^* = Q^*S^*$ then that means $P^*Q^*R^*S^*$ is a cyclic trapezoid, and hence that the perpendicular bisectors of $\overline{P^*Q^*}$ and $\overline{R^*S^*}$ are the same. Hence B, X, P^*, Q^* are symmetric around this line. This forces G^* to be the midpoint of arc R^*S^* as desired.

§2.3 USAMO 2009/6, proposed by Gabriel Carroll

Available online at <https://aops.com/community/p1485204>.

Problem statement

Let s_1, s_2, s_3, \dots be an infinite, nonconstant sequence of rational numbers, meaning it is not the case that $s_1 = s_2 = s_3 = \dots$. Suppose that t_1, t_2, t_3, \dots is also an infinite, nonconstant sequence of rational numbers with the property that $(s_i - s_j)(t_i - t_j)$ is an integer for all i and j . Prove that there exists a rational number r such that $(s_i - s_j)r$ and $(t_i - t_j)/r$ are integers for all i and j .

First we eliminate the silly edge case:

Claim — For some i and j , we have $(s_i - s_j)(t_i - t_j) \neq 0$.

Proof. Assume not. WLOG $s_1 \neq s_2$, then $t_1 = t_2$. Now select i such that $t_i \neq t_1 = t_2$. Then either $t_i - s_1 \neq 0$ or $t_i - s_2 \neq 0$, contradiction. \square

So, WLOG (by permutation) that $n = (s_1 - s_2)(t_1 - t_2) \neq 0$. By shifting and scaling appropriately, we may assume

$$s_1 = t_1 = 0, \quad s_2 = 1, \quad t_2 = n.$$

Thus we deduce

$$s_i t_i \in \mathbb{Z}, \quad s_i t_j + s_j t_i \in \mathbb{Z} \quad \forall i, j.$$

Claim — For any index i , $t_i \in \mathbb{Z}$, $s_i \in \frac{1}{n}\mathbb{Z}$.

Proof. We have $s_i t_i \in \mathbb{Z}$ and $t_i + n s_i \in \mathbb{Z}$ by problem condition. By looking at ν_p of this, we conclude $n s_i, t_i \in \mathbb{Z}$ (since if either has negative p -adic value, so does the other, and then $s_i t_i \notin \mathbb{Z}$). \square

Last claim:

Claim — If $d = \gcd t_\bullet$, then $d s_i \in \mathbb{Z}$ for all i .

Proof. Consider a prime $p \mid n$, and let $e = \nu_p(t_j)$. We will show $\nu_p(s_i) \geq -e$ for any i .

This is already true for $i = j$, so assume $i \neq j$. Assume for contradiction $\nu_p(s_i) < -e$. Then $\nu_p(t_i) > e = \nu_p(t_k)$. Since $\nu_p(s_k) \geq -e$ we deduce $\nu_p(s_i t_k) < \nu_p(s_k t_i)$; so $\nu_p(s_i t_k) \geq 0$ and $\nu_p(s_i) \geq -e$ as desired. \square