

# USAMO 2020 Solution Notes

COMPILED BY EVAN CHEN

August 29, 2020

This is an compilation of solutions for the 2020 USAMO. Some of the solutions are my own work, but many are from the official solutions provided by the organizers (for which they hold any copyrights), and others were found on the Art of Problem Solving forums.

Corrections and comments are welcome!

## Contents

<a href="#">0 Problems</a>	2
<a href="#">1 USAMO 2020/1, proposed by Zuming Feng</a>	3
<a href="#">2 USAMO 2020/2, proposed by Alex Zhai</a>	5
<a href="#">3 USAMO 2020/3, proposed by Richard Stong and Toni Blucher</a>	7
<a href="#">4 USAMO 2020/4, proposed by Ankan Bhattacharya</a>	8
<a href="#">5 USAMO 2020/5, proposed by Carl Schildkraut</a>	10
<a href="#">6 USAMO 2020/6, proposed by David Speyer and Kiran Kedlaya</a>	12

## §0 Problems

- Let  $ABC$  be a fixed acute triangle inscribed in a circle  $\omega$  with center  $O$ . A variable point  $X$  is chosen on minor arc  $AB$  of  $\omega$ , and segments  $CX$  and  $AB$  meet at  $D$ . Denote by  $O_1$  and  $O_2$  the circumcenters of triangles  $ADX$  and  $BDX$ , respectively. Determine all points  $X$  for which the area of triangle  $OO_1O_2$  is minimized.
- An empty  $2020 \times 2020 \times 2020$  cube is given, and a  $2020 \times 2020$  grid of square unit cells is drawn on each of its six faces. A *beam* is a  $1 \times 1 \times 2020$  rectangular prism. Several beams are placed inside the cube subject to the following conditions:
  - The two  $1 \times 1$  faces of each beam coincide with unit cells lying on opposite faces of the cube. (Hence, there are  $3 \cdot 2020^2$  possible positions for a beam.)
  - No two beams have intersecting interiors.
  - The interiors of each of the four  $1 \times 2020$  faces of each beam touch either a face of the cube or the interior of the face of another beam.

What is the smallest positive number of beams that can be placed to satisfy these conditions?

- Let  $p$  be an odd prime. An integer  $x$  is called a *quadratic non-residue* if  $p$  does not divide  $x - t^2$  for any integer  $t$ .

Denote by  $A$  the set of all integers  $a$  such that  $1 \leq a < p$ , and both  $a$  and  $4 - a$  are quadratic non-residues. Calculate the remainder when the product of the elements of  $A$  is divided by  $p$ .

- Suppose that  $(a_1, b_1), (a_2, b_2), \dots, (a_{100}, b_{100})$  are distinct ordered pairs of non-negative integers. Let  $N$  denote the number of pairs of integers  $(i, j)$  satisfying  $1 \leq i < j \leq 100$  and  $|a_i b_j - a_j b_i| = 1$ . Determine the largest possible value of  $N$  over all possible choices of the 100 ordered pairs.
- A finite set  $S$  of points in the coordinate plane is called *overdetermined* if  $|S| \geq 2$  and there exists a nonzero polynomial  $P(t)$ , with real coefficients and of degree at most  $|S| - 2$ , satisfying  $P(x) = y$  for every point  $(x, y) \in S$ .

For each integer  $n \geq 2$ , find the largest integer  $k$  (in terms of  $n$ ) such that there exists a set of  $n$  distinct points that is *not* overdetermined, but has  $k$  overdetermined subsets.

- Let  $n \geq 2$  be an integer. Let  $x_1 \geq x_2 \geq \dots \geq x_n$  and  $y_1 \geq y_2 \geq \dots \geq y_n$  be  $2n$  real numbers such that

$$0 = x_1 + x_2 + \dots + x_n = y_1 + y_2 + \dots + y_n,$$

$$\text{and } 1 = x_1^2 + x_2^2 + \dots + x_n^2 = y_1^2 + y_2^2 + \dots + y_n^2.$$

Prove that

$$\sum_{i=1}^n (x_i y_i - x_i y_{n+1-i}) \geq \frac{2}{\sqrt{n-1}}.$$

## §1 USAMO 2020/1, proposed by Zuming Feng

Let  $ABC$  be a fixed acute triangle inscribed in a circle  $\omega$  with center  $O$ . A variable point  $X$  is chosen on minor arc  $AB$  of  $\omega$ , and segments  $CX$  and  $AB$  meet at  $D$ . Denote by  $O_1$  and  $O_2$  the circumcenters of triangles  $ADX$  and  $BDX$ , respectively. Determine all points  $X$  for which the area of triangle  $OO_1O_2$  is minimized.

---

We prove  $[OO_1O_2] \geq \frac{1}{4}[ABC]$ , with equality if and only if  $\overline{CX} \perp \overline{AB}$ .

**First approach (Bobby Shen)** We use two simultaneous inequalities:

- Let  $M$  and  $N$  be the midpoints of  $CX$  and  $DX$ . Then  $MN$  equals the length of the  $O$ -altitude of  $\triangle OO_1O_2$ , since  $\overline{O_1O_2}$  and  $\overline{DX}$  meet at  $N$  at a right angle. Moreover, we have

$$MN = \frac{1}{2}CD \geq \frac{1}{2}h_a$$

where  $h_a$  denotes the  $A$ -altitude.

- The projection of  $O_1O_2$  onto line  $AB$  has length exactly  $AB/2$ . Thus

$$O_1O_2 \geq \frac{1}{2}AB.$$

So, we find

$$[OO_1O_2] = \frac{1}{2} \cdot MN \cdot O_1O_2 \geq \frac{1}{8}h_a \cdot AB = \frac{1}{4}[ABC].$$


Note that equality occurs in both cases if and only if  $\overline{CX} \perp \overline{AB}$ . So the area is minimized exactly when this occurs.

**Second approach (Evan's solution)** We need two claims.

**Claim** — We have  $\triangle OO_1O_2 \sim \triangle CBA$ , with opposite orientation.

*Proof.* Notice that  $\overline{OO_1} \perp \overline{AX}$  and  $\overline{O_1O_2} \perp \overline{CX}$ , so  $\angle OO_1O_2 = \angle AXC = \angle ABC$ . Similarly  $\angle OO_2O_1 = \angle BAC$ .  $\square$

Therefore, the problem is equivalent to minimizing  $O_1O_2$ .



**Claim (Salmon theorem)** — We have  $\triangle XO_1O_2 \sim \triangle XAB$ .

*Proof.* It follows from the fact that  $\triangle AO_1X \sim \triangle BO_2X$  (since  $\angle ADX = \angle XDB \implies \angle XO_1A = \angle XO_2B$ ) and that spiral similarities come in pairs.  $\square$

Let  $\theta = \angle ADX$ . The ratio of similarity in the previous claim is equal to  $\frac{XO_1}{XA} = \frac{1}{2 \sin \theta}$ . In other words,

$$O_1O_2 = \frac{AB}{2 \sin \theta}.$$

This is minimized when  $\theta = 90^\circ$ , in which case  $O_1O_2 = AB/2$  and  $[OO_1O_2] = \frac{1}{4}[ABC]$ . This completes the solution.

## §2 USAMO 2020/2, proposed by Alex Zhai


An empty  $2020 \times 2020 \times 2020$  cube is given, and a  $2020 \times 2020$  grid of square unit cells is drawn on each of its six faces. A *beam* is a  $1 \times 1 \times 2020$  rectangular prism. Several beams are placed inside the cube subject to the following conditions:

- The two  $1 \times 1$  faces of each beam coincide with unit cells lying on opposite faces of the cube. (Hence, there are  $3 \cdot 2020^2$  possible positions for a beam.)
- No two beams have intersecting interiors.
- The interiors of each of the four  $1 \times 2020$  faces of each beam touch either a face of the cube or the interior of the face of another beam.

What is the smallest positive number of beams that can be placed to satisfy these conditions?

Answer: 3030 beams.

**Construction:** We first give a construction with  $3n/2$  beams for any  $n \times n \times n$  box, where  $n$  is an even integer. Shown below is the construction for  $n = 6$ , which generalizes. (The left figure shows the cube in 3d; the right figure shows a direct view of the three visible faces.)



To be explicit, impose coordinate axes such that one corner of the cube is the origin. We specify a beam by two opposite corners. The  $3n/2$  beams come in three directions,  $n/2$  in each direction:

- $(0, 0, 0) \rightarrow (1, 1, n), (2, 2, 0) \rightarrow (3, 3, n), (4, 4, 0) \rightarrow (5, 5, n)$ , and so on;
- $(1, 0, 0) \rightarrow (2, n, 1), (3, 0, 2) \rightarrow (4, n, 3), (5, 0, 4) \rightarrow (6, n, 5)$ , and so on;
- $(0, 1, 1) \rightarrow (n, 2, 2), (0, 3, 3) \rightarrow (n, 4, 4), (0, 5, 5) \rightarrow (n, 6, 6)$ , and so on.

This gives the figure we drew earlier and shows 3030 beams is possible.

**Necessity:** We now show at least  $3n/2$  beams are necessary. Maintain coordinates, and call the beams  $x$ -beams,  $y$ -beams,  $z$ -beams according to which plane their long edges are perpendicular too. Let  $N_x, N_y, N_z$  be the number of these.

**Claim** — If  $\min(N_x, N_y, N_z) = 0$ , then at least  $n^2$  beams are needed.

*Proof.* Assume WLOG that  $N_z = 0$ . Orient the cube so the  $z$ -plane touches the ground. Then each of the  $n$  layers of the cube (from top to bottom) must be completely filled, and so at least  $n^2$  beams are necessary.  $\square$

We henceforth assume  $\min(N_x, N_y, N_z) > 0$ .

**Claim** — If  $N_z > 0$ , then we have  $N_x + N_y \geq n$ .

*Proof.* Again orient the cube so the  $z$ -plane touches the ground. We see that for each of the  $n$  layers of the cube (from top to bottom), there is at least one  $x$ -beam or  $y$ -beam. (Pictorially, some of the  $x$  and  $y$  beams form a “staircase”.) This completes the proof.  $\square$

Proceeding in a similar fashion, we arrive at the three relations

$$N_x + N_y \geq n$$

$$N_y + N_z \geq n$$

$$N_z + N_x \geq n.$$

Summing gives  $N_x + N_y + N_z \geq 3n/2$  too.

**Remark.** The problem condition has the following “physics” interpretation. Imagine the cube is a metal box which is sturdy enough that all beams must remain orthogonal to the faces of the box (i.e. the beams cannot spin). Then the condition of the problem is exactly what is needed so that, if the box is shaken or rotated, the beams will not move.

**Remark.** Walter Stromquist points out that the number of constructions with 3030 beams is actually enormous: not dividing out by isometries, the number is  $(2 \cdot 1010!)^3$ .

### §3 USAMO 2020/3, proposed by Richard Stong and Toni Blucher

Let  $p$  be an odd prime. An integer  $x$  is called a *quadratic non-residue* if  $p$  does not divide  $x - t^2$  for any integer  $t$ .

Denote by  $A$  the set of all integers  $a$  such that  $1 \leq a < p$ , and both  $a$  and  $4 - a$  are quadratic non-residues. Calculate the remainder when the product of the elements of  $A$  is divided by  $p$ .

The answer is that  $\prod_{a \in A} a \equiv 2 \pmod{p}$  regardless of the value of  $p$ . Here is the official solution, where we always work in  $\mathbb{F}_p$ .

We define

$$A = \{a \in \mathbb{F}_p \mid a, 4 - a \text{ not qr}\}$$

$$B = \{b \in \mathbb{F}_p \mid b, 4 - b \text{ qr}, b \neq 0, b \neq 4\}.$$

Note that  $a \in A \iff 4 - a \in A$  and  $b \in B \iff 4 - b \in B$ .

We now do the following magical calculation in  $\mathbb{F}_p$ :

$$\begin{aligned} \prod_{b \in B} b &= \prod_{b \in B} (4 - b) = \prod_{\substack{1 \leq y \leq (p-1)/2 \\ y \neq 2 \\ 4 - y^2 \text{ is qr}}} (4 - y^2) \\ &= \prod_{\substack{1 \leq y \leq (p-1)/2 \\ y \neq 2 \\ 4 - y^2 \text{ is qr}}} (2 + y) \prod_{\substack{1 \leq y \leq (p-1)/2 \\ y \neq 2 \\ 4 - y^2 \text{ is qr}}} (2 - y) \\ &= \prod_{\substack{1 \leq y \leq (p-1)/2 \\ y \neq 2 \\ 4 - y^2 \text{ is qr}}} (2 + y) \prod_{\substack{(p+1)/2 \leq y \leq p-1 \\ y \neq p-2 \\ 4 - y^2 \text{ is qr}}} (2 + y) \\ &= \prod_{\substack{1 \leq y \leq p-1 \\ y \neq 2, p-2 \\ 4 - y^2 \text{ is qr}}} (2 + y) \\ &= \prod_{\substack{3 \leq z \leq p+1 \\ z \neq 4, p \\ z(4-z) \text{ is qr}}} z \\ &= \prod_{\substack{0 \leq z \leq p-1 \\ z \neq 0, 4, 2 \\ z(4-z) \text{ is qr}}} z. \end{aligned}$$

Note  $z(4 - z)$  is a nonzero quadratic residue if and only if  $z \in A \cup B$ . So the right-hand side is almost the product over  $z \in A \cup B$ , except it is missing the  $z = 2$  term. Adding it in gives

$$\prod_{b \in B} b = \frac{1}{2} \prod_{\substack{0 \leq z \leq p-1 \\ z \neq 0, 4 \\ z(4-z) \text{ is qr}}} z = \frac{1}{2} \prod_{a \in A} a \prod_{b \in B} b.$$

This gives  $\prod_{a \in A} a = 2$  as desired.

### §4 USAMO 2020/4, proposed by Ankan Bhattacharya

Suppose that  $(a_1, b_1), (a_2, b_2), \dots, (a_{100}, b_{100})$  are distinct ordered pairs of nonnegative integers. Let  $N$  denote the number of pairs of integers  $(i, j)$  satisfying  $1 \leq i < j \leq 100$  and  $|a_i b_j - a_j b_i| = 1$ . Determine the largest possible value of  $N$  over all possible choices of the 100 ordered pairs.


The answer is 197. In general, if 100 is replaced by  $n \geq 2$  the answer is  $2n - 3$ .

The idea is that if we let  $P_i = (a_i, b_i)$  be a point in the coordinate plane, and let  $O = (0, 0)$  then we wish to maximize the number of triangles  $\triangle OP_i P_j$  which have area  $1/2$ . Call such a triangle *good*.

**Construction of 197 points:** It suffices to use the points  $(1, 0), (1, 1), (2, 1), (3, 1), \dots, (99, 1)$  as shown. Notice that:

- There are 98 good triangles with vertices  $(0, 0), (k, 1)$  and  $(k + 1, 1)$  for  $k = 1, \dots, 98$ .
- There are 99 good triangles with vertices  $(0, 0), (1, 0)$  and  $(k, 1)$  for  $k = 1, \dots, 99$ .

This is a total of  $98 + 99 = 197$  triangles.




**Proof that 197 points is optimal:** We proceed by induction on  $n$  to show the bound of  $2n - 3$ . The base case  $n = 2$  is evident.

For the inductive step, suppose (without loss of generality) that the point  $P = P_n = (a, b)$  is the farthest away from the point  $O$  among all points.

**Claim —** This farthest point  $P = P_n$  is part of at most two good triangles.

*Proof.* We must have  $\gcd(a, b) = 1$  for  $P$  to be in any good triangles at all, since otherwise any divisor of  $\gcd(a, b)$  also divides  $2[OPQ]$ . Now, we consider the locus of all points  $Q$  for which  $[OPQ] = 1/2$ . It consists of two parallel lines passing with slope  $OP$ , as shown.





Since  $\gcd(a, b) = 1$ , see that only two lattice points on this locus actually lie inside the quarter-circle centered at  $O$  with radius  $OP$ . Indeed if one of the points is  $(u, v)$  then the others on the line are  $(u \pm a, v \pm b)$  where the signs match. This proves the claim.  $\square$

This claim allows us to complete the induction by simply deleting  $P_n$ .

## §5 USAMO 2020/5, proposed by Carl Schildkraut

A finite set  $S$  of points in the coordinate plane is called *overdetermined* if  $|S| \geq 2$  and there exists a nonzero polynomial  $P(t)$ , with real coefficients and of degree at most  $|S| - 2$ , satisfying  $P(x) = y$  for every point  $(x, y) \in S$ .

For each integer  $n \geq 2$ , find the largest integer  $k$  (in terms of  $n$ ) such that there exists a set of  $n$  distinct points that is *not* overdetermined, but has  $k$  overdetermined subsets.

We claim the answer is  $k = 2^{n-1} - n$ . We denote the  $n$  points by  $A$ .

Throughout the solution we will repeatedly use the following fact:

### Lemma

If  $S$  is a finite set of points in the plane there is at most one polynomial with real coefficients and of degree at most  $|S| - 1$  whose graph passes through all points of  $S$ .

*Proof.* If any two of the points have the same  $x$ -coordinate then obviously no such polynomial may exist at all.

Otherwise, suppose  $f$  and  $g$  are two such polynomials. Then  $f - g$  has degree at most  $|S| - 1$ , but it has  $|S|$  roots, so is the zero polynomial.  $\square$

**Remark.** Actually Lagrange interpolation implies that such a polynomial exists as long as all the  $x$ -coordinates are different!

**Construction:** Consider the set  $A = \{(1, a), (2, b), (3, b), (4, b), \dots, (n, b)\}$ , where  $a$  and  $b$  are two distinct nonzero real numbers. Any subset of the latter  $n - 1$  points with at least one element is overdetermined, and there are  $2^{n-1} - n$  such sets.

**Bound:** Say a subset  $S$  of  $A$  is *flooded* if it is not overdetermined. For brevity, an  $m$ -set refers simply to a subset of  $A$  with  $m$  elements.

**Claim** — If  $S$  is an flooded  $m$ -set for  $m \geq 3$ , then at most one  $(m - 1)$ -subset of  $S$  is not flooded.

*Proof.* Let  $S = \{p_1, \dots, p_m\}$  be flooded. Assume for contradiction that  $S - \{p_i\}$  and  $S - \{p_j\}$  are both overdetermined. Then we can find polynomials  $f$  and  $g$  of degree at most  $m - 3$  passing through  $S - \{p_i\}$  and  $S - \{p_j\}$ , respectively.

Since  $f$  and  $g$  agree on  $S - \{p_i, p_j\}$ , which has  $m - 2$  elements, by the lemma we have  $f = g$ . Thus this common polynomial (actually of degree at most  $m - 3$ ) witnesses that  $S$  is overdetermined, which is a contradiction.  $\square$

**Claim** — For all  $m = 2, 3, \dots, n$  there are at least  $\binom{n-1}{m-1}$  flooded  $m$ -sets of  $A$ .

*Proof.* The proof is by downwards induction on  $m$ . The base case  $m = n$  is by assumption.

For the inductive step, suppose it's true for  $m$ . Each of the  $\binom{n-1}{m-1}$  flooded  $m$ -sets has at least  $m - 1$  flooded  $(m - 1)$ -subsets. Meanwhile, each  $(m - 1)$ -set has exactly  $n - (m - 1)$  parent  $m$ -sets. We conclude the number of flooded sets of size  $m - 1$  is at least

$$\frac{m-1}{n-(m-1)} \binom{n-1}{m-1} = \binom{n-1}{m-2}$$

as desired. □

This final claim completes the proof, since it shows there are at most

$$\sum_{m=2}^n \left( \binom{n}{m} - \binom{n-1}{m-1} \right) = 2^{n-1} - n$$

overdetermined sets, as desired.

**Remark** (On repeated  $x$ -coordinates). Suppose  $A$  has two points  $p$  and  $q$  with repeated  $x$ -coordinates. We can argue directly that  $A$  satisfies the bound. Indeed, any overdetermined set contains at most one of  $p$  and  $q$ . Moreover, given  $S \subseteq A - \{p, q\}$ , if  $S \cup \{p\}$  is overdetermined then  $S \cup \{q\}$  is not, and vice-versa. (Recall that overdetermined sets always have distinct  $x$ -coordinates.) This gives a bound  $[2^{n-2} - (n-2) - 1] + [2^{n-2} - 1] = 2^{n-1} - n$  already.

**Remark** (Alex Zhai). An alternative approach to the double-counting argument is to show that any overdetermined  $m$ -set has an flooded  $m$ -superset. Together with the first claim, this lets us pair overdetermined sets in a way that implies the bound.

## §6 USAMO 2020/6, proposed by David Speyer and Kiran Kedlaya

Let  $n \geq 2$  be an integer. Let  $x_1 \geq x_2 \geq \dots \geq x_n$  and  $y_1 \geq y_2 \geq \dots \geq y_n$  be  $2n$  real numbers such that

$$0 = x_1 + x_2 + \dots + x_n = y_1 + y_2 + \dots + y_n,$$

$$\text{and } 1 = x_1^2 + x_2^2 + \dots + x_n^2 = y_1^2 + y_2^2 + \dots + y_n^2.$$

Prove that

$$\sum_{i=1}^n (x_i y_i - x_i y_{n+1-i}) \geq \frac{2}{\sqrt{n-1}}.$$

We present two approaches. In both approaches, it's helpful motivation that for even  $n$ , equality occurs at

$$(x_i) = \left( \underbrace{\frac{1}{\sqrt{n}}, \dots, \frac{1}{\sqrt{n}}}_{n/2}, \underbrace{-\frac{1}{\sqrt{n}}, \dots, -\frac{1}{\sqrt{n}}}_{n/2} \right)$$

$$(y_i) = \left( \frac{n-1}{\sqrt{n(n-1)}}, \underbrace{-\frac{1}{\sqrt{n(n-1)}}, \dots, -\frac{1}{\sqrt{n(n-1)}}}_{n-1} \right)$$

**First approach (expected value)** For a permutation  $\sigma$  on  $\{1, 2, \dots, n\}$  we define

$$S_\sigma = \sum_{i=1}^n x_i y_{\sigma(i)}.$$

**Claim** — For random permutations  $\sigma$ ,  $\mathbb{E}[S_\sigma] = 0$  and  $\mathbb{E}[S_\sigma^2] = \frac{1}{n-1}$ .

*Proof.* The first one is clear.

Since  $\sum_{i < j} 2x_i x_j = -1$ , it follows that (for fixed  $i$  and  $j$ ),  $\mathbb{E}[y_{\sigma(i)} y_{\sigma(j)}] = -\frac{1}{n(n-1)}$ . Thus

$$\sum_i x_i^2 \cdot \mathbb{E}[y_{\sigma(i)}^2] = \frac{1}{n}$$

$$2 \sum_{i < j} x_i x_j \cdot \mathbb{E}[y_{\sigma(i)} y_{\sigma(j)}] = (-1) \cdot \frac{1}{n(n-1)}$$

the conclusion follows. □

**Claim** (A random variable in  $[0, 1]$  has variance at most  $1/4$ ) — If  $A$  is a random variable with mean  $\mu$  taking values in the closed interval  $[m, M]$  then

$$\mathbb{E}[(A - \mu)^2] \leq \frac{1}{4}(M - m)^2.$$

*Proof.* By shifting and scaling, we may assume  $m = 0$  and  $M = 1$ , so  $A \in [0, 1]$  and hence  $A^2 \leq A$ . Then

$$\mathbb{E}[(A - \mu)^2] = \mathbb{E}[A^2] - \mu^2 \leq \mathbb{E}[A] - \mu^2 = \mu - \mu^2 \leq \frac{1}{4}.$$

This concludes the proof. □

Thus the previous two claims together give

$$\max_{\sigma} S_{\sigma} - \min_{\sigma} S_{\sigma} \geq \sqrt{\frac{4}{n-1}} = \frac{2}{\sqrt{n-1}}.$$

But  $\sum_i x_i y_i = \max_{\sigma} S_{\sigma}$  and  $\sum_i x_i y_{n+1-i} = \min_{\sigma} S_{\sigma}$  by rearrangement inequality and therefore we are done.

**Outline of second approach (by convexity, due to Alex Zhai)** We will instead prove a converse result: given the hypotheses

- $x_1 \geq \dots \geq x_n$
- $y_1 \geq \dots \geq y_n$
- $\sum_i x_i = \sum_i y_i = 0$
- $\sum_i x_i y_i - \sum_i x_i y_{n+1-i} = \frac{2}{\sqrt{n-1}}$

we will prove that  $\sum x_i^2 \sum y_i^2 \leq 1$ .

Fix the choice of  $y$ 's. We see that we are trying to maximize a convex function in  $n$  variables  $(x_1, \dots, x_n)$  over a convex domain (actually the intersection of two planes with several half planes). So a maximum can only happen at the boundaries: when at most two of the  $x$ 's are different.

An analogous argument applies to  $y$ . In this way we find that it suffices to consider situations where  $x_{\bullet}$  takes on at most two different values. The same argument applies to  $y_{\bullet}$ .

At this point the problem can be checked directly.