

USAMO 2021 Solution Notes

COMPILED BY EVAN CHEN

April 17, 2021

This is an compilation of solutions for the 2021 USAMO. Some of the solutions are my own work, but many are from the official solutions provided by the organizers (for which they hold any copyrights), and others were found on the Art of Problem Solving forums.

Corrections and comments are welcome!

Contents

0	Problems	2
1	USAMO 2021/1, proposed by Ankan Bhattacharya	4
2	USAMO 2021/2, proposed by Zoran Sunic	5
3	USAMO 2021/3, proposed by Alex Zhai and Shaunak Kishore	7
4	USAMO 2021/4, proposed by Carl Schildkraut	8
5	USAMO 2021/5, proposed by Mohsen Jamaali	9
6	USAMO 2021/6, proposed by Ankan Bhattacharya	11

§0 Problems


1. Rectangles BCC_1B_2 , CAA_1C_2 , and ABB_1A_2 are erected outside an acute triangle ABC . Suppose that

$$\angle BC_1C + \angle CA_1A + \angle AB_1B = 180^\circ.$$

Prove that lines B_1C_2 , C_1A_2 , and A_1B_2 are concurrent.

2. The Planar National Park is a undirected 3-regular planar graph (i.e. all vertices have degree 3). A visitor walks through the park as follows: she begins at a vertex and starts walking along an edge. When she reaches the other endpoint, she turns left. On the next vertex she turns right, and so on, alternating left and right turns at each vertex. She does this until she gets back to the vertex where she started. What is the largest possible number of times she could have entered any vertex during her walk, over all possible layouts of the park?
3. Let $n \geq 2$ be an integer. An $n \times n$ board is initially empty. Each minute, you may perform one of three moves:

- If there is an L-shaped tromino region of three cells without stones on the board (see figure; rotations not allowed), you may place a stone in each of those cells.



- If all cells in a column have a stone, you may remove all stones from that column.
- If all cells in a row have a stone, you may remove all stones from that row.

For which n is it possible that, after some non-zero number of moves, the board has no stones?

4. A finite set S of positive integers has the property that, for each $s \in S$, and each positive integer divisor d of s , there exists a unique element $t \in S$ satisfying $\gcd(s, t) = d$. (The elements s and t could be equal.)

Given this information, find all possible values for the number of elements of S .

5. Let $n \geq 4$ be an integer. Find all positive real solutions to the following system of $2n$ equations:

$$\begin{aligned} a_1 &= \frac{1}{a_{2n}} + \frac{1}{a_2}, & a_2 &= a_1 + a_3, \\ a_3 &= \frac{1}{a_2} + \frac{1}{a_4}, & a_4 &= a_3 + a_5, \\ a_5 &= \frac{1}{a_4} + \frac{1}{a_6}, & a_6 &= a_5 + a_7, \\ &\vdots & &\vdots \\ a_{2n-1} &= \frac{1}{a_{2n-2}} + \frac{1}{a_{2n}}, & a_{2n} &= a_{2n-1} + a_1. \end{aligned}$$

6. Let $ABCDEF$ be a convex hexagon satisfying $\overline{AB} \parallel \overline{DE}$, $\overline{BC} \parallel \overline{EF}$, $\overline{CD} \parallel \overline{FA}$, and

$$AB \cdot DE = BC \cdot EF = CD \cdot FA.$$

Let X , Y , and Z be the midpoints of \overline{AD} , \overline{BE} , and \overline{CF} . Prove that the circumcenter of $\triangle ACE$, the circumcenter of $\triangle BDF$, and the orthocenter of $\triangle XYZ$ are collinear.

§1 USAMO 2021/1, proposed by Ankan Bhattacharya

Rectangles BCC_1B_2 , CAA_1C_2 , and ABB_1A_2 are erected outside an acute triangle ABC . Suppose that

$$\angle BC_1C + \angle CA_1A + \angle AB_1B = 180^\circ.$$

Prove that lines B_1C_2 , C_1A_2 , and A_1B_2 are concurrent.

The angle condition implies the circumcircles of the three rectangles concur at a single point P . Then $\angle CPB_2 = \angle CPA_1 = 90^\circ$, hence P lies on A_1B_2 etc., so we're done.

Remark. As one might guess from the two-sentence solution, the entire difficulty of the problem is getting the characterization of the concurrence point.

§2 USAMO 2021/2, proposed by Zoran Sunic

The Planar National Park is a undirected 3-regular planar graph (i.e. all vertices have degree 3). A visitor walks through the park as follows: she begins at a vertex and starts walking along an edge. When she reaches the other endpoint, she turns left. On the next vertex she turns right, and so on, alternating left and right turns at each vertex. She does this until she gets back to the vertex where she started. What is the largest possible number of times she could have entered any vertex during her walk, over all possible layouts of the park?

The answer is 3.

We consider the trajectory of the visitor as an ordered sequence of *turns*. A turn is defined by specifying a vertex, the incoming edge, and the outgoing edge. Hence there are six possible turns for each vertex.

Claim — Given one turn in the sequence, one can reconstruct the entire sequence of turns.


Proof. This is clear from the process's definition: given a turn t , one can compute the turn after it and the turn before it. \square

This implies already that the trajectory of the visitor, when extended to an infinite sequence, is totally periodic (not just eventually periodic), because there are finitely many possible turns, so some turn must be repeated. So, any turn appears at most once in the period of the sequence, giving a naïve bound of 6 for the original problem.

However, the following claim improves the bound to 3.


Claim — It is impossible for both of the turns $a \rightarrow b \rightarrow c$ and $c \rightarrow b \rightarrow a$ to occur in the same trajectory.

Proof. If so, then extending the path, we get $a \rightarrow b \rightarrow c \rightarrow d \rightarrow e \rightarrow \dots$ and $\dots \rightarrow e \rightarrow d \rightarrow c \rightarrow b \rightarrow a$, as illustrated below in red and blue respectively.



However, we assumed for contradiction the red and blue paths were part of the same trajectory, yet they clearly never meet. \square

It remains to give a construction showing 3 can be achieved. There are many, many valid constructions. One construction due to Danielle Wang is given here, who provided the following motivation: “I was lying in bed and drew the first thing I could think of”. The path is $CAHIFGDBAHEFGJBAC$ which visits A three times.




Remark. As the above example shows it is possible to transverse an edge more than once even in the same direction, as in edge AH above.

§3 USAMO 2021/3, proposed by Alex Zhai and Shaunak Kishore

Let $n \geq 2$ be an integer. An $n \times n$ board is initially empty. Each minute, you may perform one of three moves:

- If there is an L-shaped tromino region of three cells without stones on the board (see figure; rotations not allowed), you may place a stone in each of those cells.




- If all cells in a column have a stone, you may remove all stones from that column.
- If all cells in a row have a stone, you may remove all stones from that row.

For which n is it possible that, after some non-zero number of moves, the board has no stones?

The answer is $3 \mid n$.

Construction: For $n = 3$, the construction is fairly straightforward, shown below.



This can be extended to any $3 \mid n$.

Converse: Assume for contradiction $3 \nmid n$. We will show the task is impossible even if we allow stones to have real weights in our process. A valid elimination corresponds to a polynomial $P \in \mathbb{R}[x, y]$ such that

$$\begin{aligned} \deg_x P &\leq n - 2 \\ \deg_y P &\leq n - 2 \\ (1 + x + y)P(x, y) &\in \langle 1 + x + \dots + x^{n-1}, 1 + y + \dots + y^{n-1} \rangle. \end{aligned}$$

(Here $\langle \dots \rangle$ is an ideal of $\mathbb{R}[x, y]$.) In particular, if S is the set of n th roots of unity other than 1, we should have

$$(1 + z_1 + z_2)P(z_1, z_2) = 0$$

for any $z_1, z_2 \in S$. Since $3 \nmid n$, it follows that $1 + z_1 + z_2 \neq 0$ always.

So P vanishes on $S \times S$, a contradiction to the bounds on $\deg P$ (by, say, combinatorial nullstellensatz on any nonzero term).

§4 USAMO 2021/4, proposed by Carl Schildkraut

A finite set S of positive integers has the property that, for each $s \in S$, and each positive integer divisor d of s , there exists a unique element $t \in S$ satisfying $\gcd(s, t) = d$. (The elements s and t could be equal.)

Given this information, find all possible values for the number of elements of S .

The answer is that $|S|$ must be a power of 2 (including 1), or $|S| = 0$ (a trivial case we do not discuss further).

Construction: For any nonnegative integer k , a construction for $|S| = 2^k$ is given by

$$S = \{(p_1 \text{ or } q_1) \times (p_2 \text{ or } q_2) \times \cdots \times (p_k \text{ or } q_k)\}$$

for $2k$ distinct primes $p_1, \dots, p_k, q_1, \dots, q_k$.

Converse: the main claim is as follows.

Claim — In any valid set S , for any prime p and $x \in S$, $\nu_p(x) \leq 1$.

Proof. Assume for contradiction $e = \nu_p(x) \geq 2$.

- On the one hand, by taking x in the statement, we see $\frac{e}{e+1}$ of the elements of S are divisible by p .
- On the other hand, consider a $y \in S$ such that $\nu_p(y) = 1$ which must exist (say if $\gcd(x, y) = p$). Taking y in the statement, we see $\frac{1}{2}$ of the elements of S are divisible by p .

So $e = 1$, contradiction. □

Now since $|S|$ equals the number of divisors of any element of S , we are done.

§5 USAMO 2021/5, proposed by Mohsen Jamaali

Let $n \geq 4$ be an integer. Find all positive real solutions to the following system of $2n$ equations:

$$\begin{aligned} a_1 &= \frac{1}{a_{2n}} + \frac{1}{a_2}, & a_2 &= a_1 + a_3, \\ a_3 &= \frac{1}{a_2} + \frac{1}{a_4}, & a_4 &= a_3 + a_5, \\ a_5 &= \frac{1}{a_4} + \frac{1}{a_6}, & a_6 &= a_5 + a_7, \\ & \vdots & & \vdots \\ a_{2n-1} &= \frac{1}{a_{2n-2}} + \frac{1}{a_{2n}}, & a_{2n} &= a_{2n-1} + a_1. \end{aligned}$$

The answer is that the only solution is $(1, 2, 1, 2, \dots, 1, 2)$ which works.

We will prove a_{2k} is a constant sequence, at which point the result is obvious.

First approach (Andrew Gu) Apparently, with indices modulo $2n$, we should have

$$a_{2k} = \frac{1}{a_{2k-2}} + \frac{2}{a_{2k}} + \frac{1}{a_{2k+2}}$$

for every index k (this eliminates all a_{odd} 's). Define

$$m = \min_k a_{2k} \quad \text{and} \quad M = \max_k a_{2k}.$$

Look at the indices i and j achieving m and M to respectively get

$$\begin{aligned} m &= \frac{2}{m} + \frac{1}{a_{2i-2}} + \frac{1}{a_{2i+2}} \geq \frac{2}{m} + \frac{1}{M} + \frac{1}{M} = \frac{2}{m} + \frac{2}{M} \\ M &= \frac{2}{M} + \frac{1}{a_{2j-2}} + \frac{1}{a_{2j+2}} \leq \frac{2}{M} + \frac{1}{m} + \frac{1}{m} = \frac{2}{M} + \frac{2}{m}. \end{aligned}$$

Together this gives $m \geq M$, so $m = M$. That means a_{2i} is constant as i varies, solving the problem.

Second approach (author's solution) As before, we have

$$a_{2k} = \frac{1}{a_{2k-2}} + \frac{2}{a_{2k}} + \frac{1}{a_{2k+2}}$$

The proof proceeds in three steps.

- Define

$$S = \sum_k a_{2k}, \quad \text{and} \quad T = \sum_k \frac{1}{a_{2k}}.$$

Summing gives $S = 4T$. On the other hand, Cauchy-Schwarz says $S \cdot T \geq n^2$, so $T \geq \frac{1}{2}n$.

- On the other hand,

$$1 = \frac{1}{a_{2k-2}a_{2k}} + \frac{2}{a_{2k}^2} + \frac{1}{a_{2k}a_{2k+2}}$$

Sum this modified statement to obtain

$$n = \sum_k \left(\frac{1}{a_{2k}} + \frac{1}{a_{2k+2}} \right)^2 \stackrel{\text{QM-AM}}{\geq} \frac{1}{n} \left(\sum_k \frac{1}{a_{2k}} + \frac{1}{a_{2k+2}} \right)^2 = \frac{1}{n} (2T)^2$$

So $T \leq \frac{1}{2}n$.

- Since $T \leq \frac{1}{2}n$ and $T \geq \frac{1}{2}n$, we must have equality everywhere above. This means a_{2k} is a constant sequence.

Remark. The problem is likely intractable over \mathbb{C} , in the sense that one gets a high-degree polynomial which almost certainly has many complex roots. So it seems likely that most solutions must involve some sort of inequality, using the fact we are over $\mathbb{R}_{>0}$ instead.

§6 USAMO 2021/6, proposed by Ankan Bhattacharya


Let $ABCDEF$ be a convex hexagon satisfying $\overline{AB} \parallel \overline{DE}$, $\overline{BC} \parallel \overline{EF}$, $\overline{CD} \parallel \overline{FA}$, and

$$AB \cdot DE = BC \cdot EF = CD \cdot FA.$$

Let X , Y , and Z be the midpoints of \overline{AD} , \overline{BE} , and \overline{CF} . Prove that the circumcenter of $\triangle ACE$, the circumcenter of $\triangle BDF$, and the orthocenter of $\triangle XYZ$ are collinear.

We present two solutions.

Parallelogram solution found by contestants Note that the following figure is intentionally *not* drawn to scale, to aid legibility. We construct parallelograms $ABCE'$, etc as shown. Note that this gives two congruent triangles $A'C'E'$ and $B'D'F'$. (Assuming that triangle XYZ is non-degenerate, the triangles $A'C'E'$ and $B'D'F'$ will also be non-degenerate.)



Claim — If $AB \cdot DE = BC \cdot EF = CD \cdot FA = k$, then the circumcenters of ACE and $A'C'E'$ coincide.


Proof. The power of A to $(A'C'E')$ is $AE' \cdot AC' = BC \cdot EF = k$; same for C and E . \square

Claim — Triangle XYZ is the vector average of the (congruent) medial triangles of triangles $A'C'E'$ and $B'D'F'$.

Proof. If M and N are the midpoints of $\overline{C'E'}$ and $\overline{B'F'}$, then X is the midpoint of \overline{MN} by vector calculation. \square

Hence the orthocenter of XYZ is the midpoint of the orthocenters of the medial triangles of $A'C'E'$ and $B'D'F'$ — that is, their circumcenters.

Author's solution Call MNP and UVW the medial triangles of ACE and BDF .



Claim — In trapezoid $ABDE$, the perpendicular bisector of \overline{XY} is the same as the perpendicular bisector of the midline \overline{WN} .

Proof. This is true for any trapezoid: because $WX = \frac{1}{2}AB = YN$. □

Claim — The points V, W, M, N are cyclic.

Proof. By power of a point from Y , since

$$WY \cdot YN = \frac{1}{2}DE \cdot \frac{1}{2}AB = \frac{1}{2}EF \cdot \frac{1}{2}BC = VY \cdot YM. \quad \square$$

Applying all the cyclic variations of the above two claims, it follows that all six points U, V, W, M, N, P are concyclic, and the center of that circle coincides with the circumcenter of $\triangle XYZ$.

Remark. It is also possible to implement ideas from the first solution here, by showing all six midpoints have equal power to (XYZ) .

Claim — The orthocenter of XYZ is the midpoint of the circumcenters of $\triangle ACE$ and $\triangle BDF$.

Proof. Apply complex numbers with the unit circle coinciding with the circumcircle of

NVPWMU. Then

$$\text{orthocenter}(XYZ) = x + y + z = \frac{a + b + c + d + e + f}{2}$$

$$\text{circumcenter}(ACE) = \text{orthocenter}(MNP)$$

$$= m + n + p = \frac{c + e}{2} + \frac{e + a}{2} + \frac{a + c}{2} = a + c + e$$

$$\text{circumcenter}(BDF) = \text{orthocenter}(UVW)$$

$$= u + v + w = \frac{d + f}{2} + \frac{f + b}{2} + \frac{b + d}{2} = b + d + f. \quad \square$$