

# USA IMO TST 2014 Solutions

United States of America — IMO Team Selection Tests

EVAN CHEN 《陳誼廷》

55<sup>th</sup> IMO 2014 South Africa

## Contents

<b>1</b>	<b>Solutions to Day 1</b>	<b>2</b>
1.1	Solution to TST 1 . . . . .	2
1.2	Solution to TST 2, by Victor Wang . . . . .	3
1.3	Solution to TST 3 . . . . .	5
<b>2</b>	<b>Solutions to Day 2</b>	<b>6</b>
2.1	Solution to TST 4 . . . . .	6
2.2	Solution to TST 5 . . . . .	7
2.3	Solution to TST 6 . . . . .	8

## §1 Solutions to Day 1

### §1.1 Solution to TST 1

Let  $ABC$  be an acute triangle, and let  $X$  be a variable interior point on the minor arc  $BC$  of its circumcircle. Let  $P$  and  $Q$  be the feet of the perpendiculars from  $X$  to lines  $CA$  and  $CB$ , respectively. Let  $R$  be the intersection of line  $PQ$  and the perpendicular from  $B$  to  $AC$ . Let  $\ell$  be the line through  $P$  parallel to  $XR$ . Prove that as  $X$  varies along minor arc  $BC$ , the line  $\ell$  always passes through a fixed point.

---

The fixed point is the orthocenter, since  $\ell$  is a Simson line. See Lemma 4.4 of *Euclidean Geometry in Math Olympiads*.

## §1.2 Solution to TST 2, by Victor Wang

Let  $a_1, a_2, a_3, \dots$  be a sequence of integers, with the property that every consecutive group of  $a_i$ 's averages to a perfect square. More precisely, for all positive integers  $n$  and  $k$ , the quantity

$$\frac{a_n + a_{n+1} + \dots + a_{n+k-1}}{k}$$

is always the square of an integer. Prove that the sequence must be constant (all  $a_i$  are equal to the same perfect square).

Let  $\nu_p(n)$  denote the largest exponent of  $p$  dividing  $n$ . The problem follows from the following proposition.

### Proposition

Let  $(a_n)$  be a sequence of integers and let  $p$  be a prime. Suppose that every consecutive group of  $a_i$ 's with length at most  $p$  averages to a perfect square. Then  $\nu_p(a_i)$  is independent of  $i$ .

We proceed by induction on the smallest value of  $\nu_p(a_i)$  as  $i$  ranges (which must be even, as each of the  $a_i$  are themselves a square). First we prove two claims.

**Claim —** If  $j \equiv k \pmod{p}$  then  $a_j \equiv a_k \pmod{p}$ .

*Proof.* Taking groups of length  $p$  in our given, we find that  $p \mid a_j + \dots + a_{j+p-1}$  and  $p \mid a_{j+1} + \dots + a_{j+p}$  for any  $j$ . So  $a_j \equiv a_{j+p} \pmod{p}$  and the conclusion follows.  $\square$

**Claim —** If some  $a_i$  is divisible by  $p$  then all of them are.

*Proof.* The case  $p = 2$  is trivial so assume  $p \geq 3$ . Without loss of generality (via shifting indices) assume that  $a_1 \equiv 0 \pmod{p}$ , and define

$$S_n = a_1 + a_2 + \dots + a_n \equiv a_2 + \dots + a_n \pmod{p}.$$

Call an integer  $k$  with  $2 \leq k < p$  a **pivot** if  $1 - k^{-1}$  is a quadratic nonresidue modulo  $p$ .

We claim that for any pivot  $k$ ,  $S_k \equiv 0 \pmod{p}$ . If not, then

$$\frac{a_1 + a_2 + \dots + a_k}{k} \text{ and } \frac{a_2 + \dots + a_k}{k-1}$$

are both quadratic residues. Division implies that  $\frac{k-1}{k} = 1 - k^{-1}$  is a quadratic residue, contradiction.

Next we claim that there is an integer  $m$  with  $S_m \equiv S_{m+1} \equiv 0 \pmod{p}$ , which implies  $p \mid a_{m+1}$ . If 2 is a pivot, then we simply take  $m = 1$ . Otherwise, there are  $\frac{1}{2}(p-1)$  pivots, one for each nonresidue (which includes neither 0 nor 1), and all pivots lie in  $[3, p-1]$ , so we can find an  $m$  such that  $m$  and  $m+1$  are both pivots.

Repeating this procedure starting with  $a_{m+1}$  shows that  $a_{2m+1}, a_{3m+1}, \dots$  must all be divisible by  $p$ . Combined with the first claim and the fact that  $m < p$ , we find that all the  $a_i$  are divisible by  $p$ .  $\square$

The second claim establishes the base case of our induction. Now assume all  $a_i$  are divisible by  $p$  and hence  $p^2$ . Then all the averages in our proposition (with length at most  $p$ ) are divisible by  $p$  and hence  $p^2$ . Thus the map  $a_i \mapsto \frac{1}{p^2}a_i$  gives a new sequence satisfying the proposition, and our inductive hypothesis completes the proof.

**Remark.** There is a subtle bug that arises if one omits the condition that  $k \leq p$  in the proposition. When  $k = p^2$  the average  $\frac{a_1 + \dots + a_{p^2}}{p^2}$  is not necessarily divisible by  $p$  even if all the  $a_i$  are. Hence it is not valid to divide through by  $p$ . This is why the condition  $k \leq p$  was added.

### §1.3 Solution to TST 3

Let  $n$  be an even positive integer, and let  $G$  be an  $n$ -vertex (simple) graph with exactly  $\frac{n^2}{4}$  edges. An unordered pair of distinct vertices  $\{x, y\}$  is said to be *amicable* if they have a common neighbor (there is a vertex  $z$  such that  $xz$  and  $yz$  are both edges). Prove that  $G$  has at least  $2\binom{n/2}{2}$  pairs of vertices which are amicable.

First, we prove the following lemma. ([https://en.wikipedia.org/wiki/Friendship\\_paradox](https://en.wikipedia.org/wiki/Friendship_paradox)).

**Lemma** (On average, your friends are more popular than you)

For a vertex  $v$ , let  $a(v)$  denote the average degree of the neighbors of  $v$  (setting  $a(v) = 0$  if  $\deg v = 0$ ). Then

$$\sum_v a(v) \geq \sum_v \deg v = 2\#E.$$

*Proof.* Ignoring isolated vertices, we can write

$$\begin{aligned} \sum_v a(v) &= \sum_v \frac{\sum_{w \sim v} \deg w}{\deg v} \\ &= \sum_v \sum_{w \sim v} \frac{\deg w}{\deg v} \\ &= \sum_{\text{edges } vw} \left( \frac{\deg w}{\deg v} + \frac{\deg v}{\deg w} \right) \\ &\stackrel{\text{AM-GM}}{\geq} \sum_{\text{edges } vw} 2 = 2\#E = \sum_v \deg v \end{aligned}$$

as desired. □

**Corollary** (On average, your most popular friend is more popular than you)

For a vertex  $v$ , let  $m(v)$  denote the maximum degree of the neighbors of  $v$  (setting  $m(v) = 0$  if  $\deg v = 0$ ). Then

$$\sum_v m(v) \geq \sum_v \deg v = 2\#E.$$

We can use this to count amicable pairs by noting that any particular vertex  $v$  is in at least  $m(v) - 1$  amicable pairs. So, the number of amicable pairs is at least

$$\frac{1}{2} \sum_v (m(v) - 1) \geq \#E - \frac{1}{2}\#V.$$

Note that up until now we haven't used any information about  $G$ . But now if we plug in  $\#E = n^2/4$ ,  $\#V = n$ , then we get exactly the desired answer. (Equality holds for  $G = K_{n/2, n/2}$ .)

## §2 Solutions to Day 2

### §2.1 Solution to TST 4

Let  $n$  be a positive even integer, and let  $c_1, c_2, \dots, c_{n-1}$  be real numbers satisfying

$$\sum_{i=1}^{n-1} |c_i - 1| < 1.$$

Prove that

$$2x^n - c_{n-1}x^{n-1} + c_{n-2}x^{n-2} - \dots - c_1x^1 + 2$$

has no real roots.

We will prove the polynomial is positive for all  $x \in \mathbb{R}$ . As  $c_i > 0$ , the result is vacuous for  $x \leq 0$ , so we restrict attention to  $x > 0$ .

Then letting  $c_i = 1 - d_i$  for each  $i$ , the inequality we want to prove becomes

$$x^n + 1 + \frac{x^{n+1} + 1}{x + 1} > \sum_1^{n-1} d_i x^i \quad \text{given } \sum |d_i| < 1.$$

But obviously  $x^n + 1 > x^i$  for any  $1 \leq i \leq n-1$  and  $x > 0$ . So in fact  $x^n + 1 > \sum_1^{n-1} |d_i| x^i$  holds for  $x > 0$ , as needed.

## §2.2 Solution to TST 5

Let  $ABCD$  be a cyclic quadrilateral, and let  $E, F, G,$  and  $H$  be the midpoints of  $AB, BC, CD,$  and  $DA$  respectively. Let  $W, X, Y$  and  $Z$  be the orthocenters of triangles  $AHE, BEF, CFG$  and  $DGH,$  respectively. Prove that the quadrilaterals  $ABCD$  and  $WXYZ$  have the same area.

The following solution is due to Grace Wang.

We begin with:

**Claim** — Point  $W$  has coordinates  $\frac{1}{2}(2a + b + d)$ .

*Proof.* The orthocenter of  $\triangle DAB$  is  $d + a + b$ , and  $\triangle AHE$  is homothetic to  $\triangle DAB$  through  $A$  with ratio  $1/2$ . Hence  $w = \frac{1}{2}(a + (d + a + b))$  as needed.  $\square$

By symmetry, we have

$$\begin{aligned} w &= \frac{1}{2}(2a + b + d) \\ x &= \frac{1}{2}(2b + c + a) \\ y &= \frac{1}{2}(2c + d + b) \\ z &= \frac{1}{2}(2d + a + c). \end{aligned}$$

We see that  $w - y = a - c$ ,  $x - z = b - d$ . So the diagonals of  $WXYZ$  have the same length as those of  $ABCD$  as well as the same directed angle between them. This implies the areas are equal, too.

### §2.3 Solution to TST 6

For a prime  $p$ , a subset  $S$  of residues modulo  $p$  is called a *sum-free multiplicative subgroup* of  $\mathbb{F}_p$  if

- there is a nonzero residue  $\alpha$  modulo  $p$  such that  $S = \{1, \alpha, \alpha^2, \dots\}$  (all considered mod  $p$ ), and
- there are no  $a, b, c \in S$  (not necessarily distinct) such that  $a + b \equiv c \pmod{p}$ .

Prove that for every integer  $N$ , there is a prime  $p$  and a sum-free multiplicative subgroup  $S$  of  $\mathbb{F}_p$  such that  $|S| \geq N$ .

We first prove the following general lemma.

#### Lemma

If  $f, g \in \mathbb{Z}[X]$  are relatively prime nonconstant polynomials, then for sufficiently large primes  $p$ , they have no common root modulo  $p$ .

*Proof.* By Bézout Lemma, there exist polynomials  $a(X)$  and  $b(X)$  in  $\mathbb{Z}[X]$  and a nonzero constant  $c \in \mathbb{Z}$  satisfying the identity

$$a(X)f(X) + b(X)g(X) \equiv c.$$

So, plugging in  $X = r$  we get  $p \mid c$ , so the set of permissible primes  $p$  is finite.  $\square$

With this we can give the construction.

**Claim** — Suppose that

- $n$  is a positive integer with  $n \not\equiv 0 \pmod{3}$ ;
- $p$  is a prime which is  $1 \pmod{n}$ ; and
- $\alpha$  is a primitive  $n$ 'th root of unity modulo  $p$ .

Then  $|S| = n$  and, if  $p$  is sufficiently large in  $n$ , is also sum-free.

*Proof.* The assertion  $|S| = n$  is immediate from the choice of  $\alpha$ . As for sum-free, assume for contradiction that

$$1 + \alpha^k \equiv \alpha^m \pmod{p}$$

for some integers  $k, m \in \mathbb{Z}$ . This means  $(X + 1)^n - 1$  and  $X^n - 1$  have common root  $X = \alpha^k$ .

But

$$\gcd_{\mathbb{Z}[x]} \left( (X + 1)^n - 1, X^n - 1 \right) = 1 \quad \forall n \not\equiv 0 \pmod{3}$$

because when  $3 \nmid n$  the two polynomials have no common complex roots. (Indeed, if  $|\omega| = |1 + \omega| = 1$  then  $\omega = -\frac{1}{2} \pm \frac{\sqrt{3}}{2}i$ .)

Thus  $p$  is bounded by the lemma, as desired.  $\square$