



Saturday, June 11, 2016

Problem 1. Let ABC be an acute triangle with orthocenter H . Let G be the point such that the quadrilateral $ABGH$ is a parallelogram. Let I be the point on the line GH such that AC bisects HI . Suppose that the line AC intersects the circumcircle of the triangle GCI at C and J . Prove that $IJ = AH$.

Problem 2. Suppose that a sequence of a_1, a_2, \dots of positive real numbers satisfies

$$a_{k+1} \geq \frac{ka_k}{a_k^2 + (k-1)}$$

for every positive integer k . Prove that $a_1 + a_2 + \dots + a_n \geq n$ for every $n \geq 2$.

Problem 3. Let S be a nonempty set of positive integers. We say that a positive integer n is *clean* if it has a unique representation as a sum of an odd number of distinct elements from S . Prove that there exist infinitely many positive integers that are not clean.

*Time limit: 4 hours and 30 minutes.
Each problem is worth seven points.*



Saturday, June 18, 2016

Problem 4. In Lineland there are $n \geq 1$ towns, arranged along a road running from left to right. Each town has a *left bulldozer* (put to the left of the town and facing left) and a *right bulldozer* (put to the right of the town and facing right). The sizes of the $2n$ bulldozers are distinct. Every time when a right and a left bulldozer confront each other, the larger bulldozer pushes the smaller one off the road. On the other hand, the bulldozers are quite unprotected at their rears; so, if a bulldozer reaches the rear-end of another one, the first one pushes the second one off the road, regardless of their sizes.

Let A and B be two towns, with B to the right of A . We say that town A can *sweep* town B *away* if the right bulldozer of A can move over to B pushing off all bulldozers it meets. Similarly, B can sweep A away if the left bulldozer of B can move to A pushing off all bulldozers of all towns on its way.

Prove that there is exactly one town which cannot be swept away by any other one.

Problem 5. Let m and n be positive integers such that $m > n$. Define $x_k = (m+k)/(n+k)$ for $k = 1, 2, \dots, n+1$. Prove that if all the numbers x_1, x_2, \dots, x_{n+1} are integers, then $x_1 x_2 \dots x_{n+1} - 1$ is divisible by an odd prime.

Problem 6. Let ABC be a triangle with $CA \neq CB$. Let D , F , and G be the midpoints of the sides AB , AC , and BC , respectively. A circle Γ passing through C and tangent to AB at D meets the segments AF and BG at H and I , respectively. The points H' and I' are symmetric to H and I about F and G , respectively. The line $H'I'$ meets CD and FG at Q and M , respectively. The line CM meets Γ again at P . Prove that $CQ = QP$.

*Time limit: 4 hours and 30 minutes.
Each problem is worth seven points.*