

Shortlist 2007 N5

Evan Chen

TWITCH SOLVES ISL

Episode 13

Problem

Find all functions $f: \mathbb{N} \rightarrow \mathbb{N}$ such that $f(1) = 1$ and for all positive integers m and n and primes p , the number $f(m + n)$ is divisible by p if and only if $f(m) + f(n)$ is divisible by p .

Video

<https://youtu.be/0ExdK27KHx8>

External Link

<https://aops.com/community/p1187222>

Solution

The answer is that f must be the identity (which works). We let $\text{rad } n$ denote the product of distinct prime factors of n . We also let $P(m, n)$ denote the given statement.

Remark. The original problem gave the hypothesis f was surjective in place of $f(1) = 1$. Our version is better: to see that f surjective implies $f(1) = 1$ anyways, consider $P(1, N - 1)$ if $f(N) = 1$ held for any $N \geq 2$.

Claim. We have $\text{rad } f(2^e) = 2$ for every positive integer e ; that is, $f(2), f(4), \dots$ are all powers of 2.

Proof. Follows by taking $P(2^{e-1}, 2^{e-1})$ and induction. □

Claim. We have $f(2) = 2, f(3) = 3, f(4) = 4$.

Proof. First, $P(3, 1)$ gives that $f(3) = 2^y - 1$ for some y . Now if $f(2) = 2^x$ then $P(2, 1)$ gives

$$\text{rad}(2^x + 1) = \text{rad}(2^y - 1).$$

We now consider two cases.

- If $x = 3$ then this forces $y = 2$. So $f(2) = 8$ and $f(3) = 3$. But $\text{rad}(1 + f(4)) = \text{rad}(f(2) + f(3)) = 11$, so there is no possible value of $f(4)$, contradiction.
- Otherwise, let p be a primitive prime divisor of $2^{2x} - 1$ which exists by Zsigmondy. It also divides the right-hand side, so $2x \mid y$, and $2^{2x} - 1 \mid 2^y - 1$. This could only happen if $2^x - 1 = 1$, so $x = 1$. This gives $y = 2$ (again by Zsigmondy) and we're done.

Hence $f(2) = 2$ and $f(3) = 3$. Now $\text{rad}(1 + f(4)) = \text{rad}(f(2) + f(3)) = 5$, and since $f(4)$ is a power of 2, Zsigmondy implies $f(4) = 4$. □

Moving forward, we apply the following simple consequence of Zsigmondy:

Lemma. If $\text{rad}(2^x - 1) = \text{rad}(2^y - 1)$ then $x = y$. Similarly if $\text{rad}(2^x + 1) = \text{rad}(2^y + 1)$ then $x = y$.

We now prove by induction on $e \geq 1$ the statement that

$$f(n) = n \quad \text{for } 2^e \leq n \leq 2^{e+1}.$$

The base case is already set for us. For the inductive step, suppose f is the identity for $n \leq 2^e$. We proceed in three steps:

- First, we have

$$\begin{aligned} \text{rad}(1 + f(2^{e+1} - 1)) &= 2 \\ \text{rad}(f(2^{e+1} - 1)) &= \text{rad}(f(2^e - 1) + f(2^e)) = \text{rad}(2^{e+1} - 1). \end{aligned}$$

The first equation just says $f(2^{e+1} - 1) = 2^t - 1$ for some t . So the latter equation, together with the lemma, gives $f(2^{e+1} - 1) = 2^{e+1} - 1$.

- Next, we have

$$\begin{aligned} \text{rad}(f(2^{e+1}) + 1) &= \text{rad } f(2^{e+1} + 1) = \text{rad}((2^{e+1} - 1) + 2) \\ &= \text{rad}(f(2^{e+1} - 1) + f(2)) = \text{rad}(2^{e+1} + 1) \end{aligned}$$

which gives $f(2^{e+1}) = 2^{e+1}$.

- Now assume $2^e + 1 \leq n \leq 2^{e+1} - 2$. Let $1 \leq k \leq 2^e - 2$ be such that $n + k = 2^{e+1} - 1$. Then

$$\begin{aligned}\operatorname{rad}(f(n) + k) &= \operatorname{rad}(f(n) + f(k)) = \operatorname{rad}(f(n + k)) = \operatorname{rad}(2^{e+1} - 1) \\ \operatorname{rad}(f(n) + k + 1) &= \operatorname{rad}(f(n) + f(k + 1)) = \operatorname{rad}(f(n + k + 1)) = 2.\end{aligned}$$

So we have an integer x (depending on k) such that

$$f(n) + k + 1 = 2^x \implies \operatorname{rad}(2^x - 1) = \operatorname{rad}(f(n) + k) = \operatorname{rad}(2^{e+1} - 1).$$

Hence $x = e + 1$. And so $f(n) = n$.

This completes the induction and hence the problem is solved.