

JMO 2020/2

Evan Chen

TWITCH SOLVES ISL

Episode 16

Problem

Let ω be the incircle of a fixed equilateral triangle ABC . Let ℓ be a variable line that is tangent to ω and meets the interior of segments BC and CA at points P and Q , respectively. A point R is chosen such that $PR = PA$ and $QR = QB$. Find all possible locations of the point R , over all choices of ℓ .

Video


<https://youtu.be/r7j0oRtpErA>

External Link

<https://aops.com/community/p15952801>

Solution

Let r be the inradius. Let T be the tangency point of \overline{PQ} on arc \widehat{DE} of the incircle, which we consider varying. We define R_1 and R_2 to be the two intersections of the circle centered at P with radius PA , and the circle centered at Q with radius QB . We choose R_1 to lie on the opposite side of C as line PQ .



Claim. The point R_1 is the unique point on ray TI with $R_1I = 2r$.

Proof. Define S to be the point on ray TI with $SI = 2r$. Note that there is a homothety at I which maps $\triangle DTE$ to $\triangle ASB$, for some point S .

Note that since $TASD$ is an isosceles trapezoid, it follows $PA = PS$. Similarly, $QB = QS$. So it follows that $S = R_1$. \square

Since T can be any point on the open arc \widehat{DE} , it follows that the locus of R_1 is exactly the open 120° arc of \widehat{AB} of the circle centered at I with radius $2r$ (i.e. the circumcircle of ABC).

It remains to characterize R_2 . Since $TI = r$, $IR_1 = 2r$, it follows $TR_2 = 3r$ and $IR_2 = 4r$. Define A' on ray DI such that $A'I = 4r$, and B' on ray IE such that $B'I = 4r$. Then it follows, again by homothety, that the locus of R_2 is the 120° arc $\widehat{A'B'}$ of the circle centered at I with radius $4r$.

In conclusion, the locus of R is the two open 120° arcs we identified.