

# USAMO 2020/4

Evan Chen

TWITCH SOLVES ISL

Episode 16

## Problem

Suppose that  $(a_1, b_1), (a_2, b_2), \dots, (a_{100}, b_{100})$  are distinct ordered pairs of nonnegative integers. Let  $N$  denote the number of pairs of integers  $(i, j)$  satisfying  $1 \leq i < j \leq 100$  and  $|a_i b_j - a_j b_i| = 1$ . Determine the largest possible value of  $N$  over all possible choices of the 100 ordered pairs.

## Video

<https://youtu.be/r7j0oRtpErA?>

## External Link

<https://aops.com/community/p15952792>

## Solution


The answer is 197. In general, if 100 is replaced by  $n \geq 2$  the answer is  $2n - 3$ .

The idea is that if we let  $P_i = (a_i, b_i)$  be a point in the coordinate plane, and let  $O = (0, 0)$  then we wish to maximize the number of triangles  $\triangle OP_i P_j$  which have area  $1/2$ . Call such a triangle *good*.

**Construction of 197 points.** It suffices to use the points  $(1, 0), (1, 1), (2, 1), (3, 1), \dots, (99, 1)$  as shown. Notice that:

- There are 98 good triangles with vertices  $(0, 0), (k, 1)$  and  $(k+1, 1)$  for  $k = 1, \dots, 98$ .
- There are 99 good triangles with vertices  $(0, 0), (1, 0)$  and  $(k, 1)$  for  $k = 1, \dots, 99$ .

This is a total of  $98 + 99 = 197$  triangles.




**Proof that 197 points is optimal.** We proceed by induction on  $n$  to show the bound of  $2n - 3$ . The base case  $n = 2$  is evident.

For the inductive step, suppose (without loss of generality) that the point  $P = P_n = (a, b)$  is the farthest away from the point  $O$  among all points.

**Claim.** This farthest point  $P = P_n$  is part of at most two good triangles.

*Proof.* We must have  $\gcd(a, b) = 1$  for  $P$  to be in any good triangles at all, since otherwise any divisor of  $\gcd(a, b)$  also divides  $2[OPQ]$ . Now, we consider the locus of all points  $Q$  for which  $[OPQ] = 1/2$ . It consists of two parallel lines passing with slope  $OP$ , as shown.



Since  $\gcd(a, b) = 1$ , see that only two lattice points on this locus actually lie inside the quarter-circle centered at  $O$  with radius  $OP$ . Indeed if one of the points is  $(u, v)$  then the others on the line are  $(u \pm a, v \pm b)$  where the signs match. This proves the claim.  $\square$

This claim allows us to complete the induction by simply deleting  $P_n$ .