USAMO 2020/5 Evan Chen

TWITCH SOLVES ISL

Episode 16

Problem

A finite set S of points in the coordinate plane is called *overdetermined* if $|S| \ge 2$ and there exists a nonzero polynomial P(t), with real coefficients and of degree at most |S|-2, satisfying P(x) = y for every point $(x, y) \in S$.

For each integer $n \ge 2$, find the largest integer k (in terms of n) such that there exists a set of n distinct points that is not overdetermined, but has k overdetermined subsets.

Video

https://youtu.be/r7j0oRtpErA

External Link

https://aops.com/community/p15952824

Solution

We claim the answer is $k = 2^{n-1} - n$. We denote the *n* points by *A*. Throughout the solution we will repeatedly use the following fact:

Lemma. If S is a finite set of points in the plane there is at most one polynomial with real coefficients and of degree at most |S| - 1 whose graph passes through all points of S.

Proof. If any two of the points have the same x-coordinate then obviously no such polynomial may exist at all.

Otherwise, suppose f and g are two such polynomials. Then f - g has degree at most |S| - 1, but it has |S| roots, so is the zero polynomial.

Remark. Actually Lagrange interpolation implies that such a polynomial exists as long as all the *x*-coordinates are different!

Construction: Consider the set $A = \{(1, a), (2, b), (3, b), (4, b), \dots, (n, b)\}$, where a and b are two distinct nonzero real numbers. Any subset of the latter n - 1 points with at least one element is overdetermined, and there are $2^{n-1} - n$ such sets.

Bound: Say a subset S of A is *flooded* if it is not overdetermined. For brevity, an m-set refers simply to a subset of A with m elements.

Claim. If S is an flooded m-set for $m \ge 3$, then at most one (m-1)-subset of S is not flooded.

Proof. Let $S = \{p_1, \ldots, p_m\}$ be flooded. Assume for contradiction that $S - \{p_i\}$ and $S - \{p_j\}$ are both overdetermined. Then we can find polynomials f and g of degree at most m - 3 passing through $S - \{p_i\}$ and $S - \{p_j\}$, respectively.

Since f and g agree on $S - \{p_i, p_j\}$, which has m - 2 elements, by the lemma we have f = g. Thus this common polynomial (actually of degree at most m - 3) witnesses that S is overdetermined, which is a contradiction.

Claim. For all m = 2, 3, ..., n there are at least $\binom{n-1}{m-1}$ flooded *m*-sets of *A*.

Proof. The proof is by downwards induction on m. The base case m = n is by assumption. For the inductive step, suppose it's true for m. Each of the $\binom{n-1}{m-1}$ flooded m-sets has at

least m-1 flooded (m-1)-subsets. Meanwhile, each (m-1)-set has exactly n-(m-1)parent *m*-sets. We conclude the number of flooded sets of size m-1 is at least

$$\frac{m-1}{n-(m-1)}\binom{n-1}{m-1} = \binom{n-1}{m-2}$$

as desired.

This final claim completes the proof, since it shows there are at most

$$\sum_{m=2}^{n} \left(\binom{n}{m} - \binom{n-1}{m-1} \right) = 2^{n-1} - n$$

overdetermined sets, as desired.

Remark (On repeated x-coordinates). Suppose A has two points p and q with repeated x-coordinates. We can argue directly that A satisfies the bound. Indeed, any overdetermined set contains at most one of p and q. Moreover, given $S \subseteq A - \{p,q\}$, if $S \cup \{p\}$ is overdetermined then $S \cup \{q\}$ is not, and vice-versa. (Recall that overdetermined sets always have distinct x-coordinates.) This gives a bound $[2^{n-2} - (n-2) - 1] + [2^{n-2} - 1] = 2^{n-1} - n$ already.

Remark (Alex Zhai). An alternative approach to the double-counting argument is to show that any overdetermined m-set has an flooded m-superset. Together with the first claim, this lets us pair overdetermined sets in a way that implies the bound.