

TSTST 2020/7

Evan Chen

TWITCH SOLVES ISL

Episode 50

Problem

Find all nonconstant polynomials $P(z)$ with complex coefficients for which all complex roots of the polynomials $P(z)$ and $P(z) - 1$ have absolute value 1.

Video

<https://youtu.be/jsw3c3yAn7o>

Solution

The answer is $P(x)$ should be a polynomial of the form $P(x) = \lambda x^n - \mu$ where $|\lambda| = |\mu|$ and $\operatorname{Re} \mu = -\frac{1}{2}$. One may check these all work; let's prove they are the only solutions.

First approach (Evan Chen) We introduce the following notations:

$$\begin{aligned} P(x) &= c_n x^n + c_{n-1} x^{n-1} + \cdots + c_1 x + c_0 \\ &= c_n (x + \alpha_1) \cdots (x + \alpha_n) \\ P(x) - 1 &= c_n (x + \beta_1) \cdots (x + \beta_n) \end{aligned}$$

By taking conjugates,

$$\begin{aligned} (x + \alpha_1) \cdots (x + \alpha_n) &= (x + \beta_1) \cdots (x + \beta_n) + c_n^{-1} \\ \implies \left(x + \frac{1}{\alpha_1}\right) \cdots \left(x + \frac{1}{\alpha_n}\right) &= \left(x + \frac{1}{\beta_1}\right) \cdots \left(x + \frac{1}{\beta_n}\right) + (\overline{c_n})^{-1} \quad (\spadesuit) \end{aligned}$$

The equation (\spadesuit) is the main player:

Claim. We have $c_k = 0$ for all $k = 1, \dots, n-1$.

Proof. By comparing coefficients of x^k in (\spadesuit) we obtain

$$\frac{c_{n-k}}{\prod_i \alpha_i} = \frac{c_{n-k}}{\prod_i \beta_i}$$

but $\prod_i \alpha_i - \prod_i \beta_i = \frac{1}{c_n} \neq 0$. Hence $c_k = 0$. \square

It follows that $P(x)$ must be of the form $P(x) = \lambda x^n - \mu$, so that $P(x) = \lambda x^n - (\mu + 1)$. This requires $|\mu| = |\mu + 1| = |\lambda|$ which is equivalent to the stated part.

Second approach (from the author) We let $A = P$ and $B = P - 1$ to make the notation more symmetric. We will as before show that A and B have all coefficients equal to zero other than the leading and constant coefficient; the finish is the same.

First, we rule out double roots.

Claim. Neither A nor B have double roots.

Proof. Suppose that b is a double root of B . By differentiating, we obtain $A' = B'$, so $A'(b) = 0$. However, by Gauss-Lucas, this forces $A(b) = 0$, contradiction. \square

Let $\omega = e^{2\pi i/n}$, let a_1, \dots, a_n be the roots of A , and let b_1, \dots, b_n be the roots of B . For each k , let A_k and B_k be the points in the complex plane corresponding to a_k and b_k .

Claim (Main claim). For any i and j , $\frac{a_i}{a_j}$ is a power of ω .

Proof. Note that

$$\frac{a_i - b_1}{a_j - b_1} \cdots \frac{a_i - b_n}{a_j - b_n} = \frac{B(a_i)}{B(a_j)} = \frac{A(a_i) - 1}{A(a_j) - 1} = \frac{0 - 1}{0 - 1} = 1.$$

Since the points A_i, A_j, B_k all lie on the unit circle, interpreting the left-hand side geometrically gives

$$\angle A_i B_1 A_j + \cdots + \angle A_i B_n A_j = 0 \implies n \widehat{A_i A_j} = 0,$$

where angles are directed modulo 180° and arcs are directed modulo 360° . This implies that $\frac{a_i}{a_j}$ is a power of ω . \square

Now the finish is easy: since a_1, \dots, a_n are all different, they must be $a_1\omega^0, \dots, a_1\omega^{n-1}$ in some order; this shows that A is a multiple of $x^n - a_1^n$, as needed.