

# ELMO 2010/2

Evan Chen

TWITCH SOLVES ISL

Episode 59

## Problem

Let  $r$  and  $s$  be positive integers. Define  $a_0 = 0$ ,  $a_1 = 1$ , and  $a_n = ra_{n-1} + sa_{n-2}$  for  $n \geq 2$ . Let  $f_n = a_1 a_2 \dots a_n$ . Prove that  $\frac{f_n}{f_k f_{n-k}}$  is an integer for all integers  $n$  and  $k$  such that  $0 < k < n$ .

## Video

<https://youtu.be/-tfVetEaVdo>

## External Link

<https://aops.com/community/p2731491>

## Solution

It's equivalent to show that for integers  $k, m$  we have

$$a_1 \dots a_k \mid a_m \dots a_{m+k-1}. \quad (\spadesuit)$$

From the theory of linear recurrences, we know there are algebraic integers  $\alpha$  and  $\beta$  such that  $a_n$  has a closed form

$$a_n = \frac{\alpha^n - \beta^n}{\alpha - \beta}.$$

(Namely,  $\alpha$  and  $\beta$  are the roots of the polynomial  $T^2 - rT - s = 0$ . Note that since  $s > 0$  it follows  $\alpha \neq \beta$ .)

Then,  $(\spadesuit)$  rewrites as

$$\frac{(\alpha^m - \beta^m)(\alpha^{m+1} - \beta^{m+1}) \dots (\alpha^{m+k-1} - \beta^{m+k-1})}{(\alpha - \beta)(\alpha^2 - \beta^2) \dots (\alpha^k - \beta^k)}.$$

**Claim.** We get a divisibility as polynomials in  $\alpha$  and  $\beta$ .

*Proof.* Since this is homogeneous, it suffices to show for  $\beta = 1, \alpha = X$ :

$$\frac{(X^m - 1^m)(X^{m+1} - 1^{m+1}) \dots (X^{m+k-1} - 1^{m+k-1})}{(X - 1)(X^2 - 1^2) \dots (X^k - 1^k)}$$

If  $\zeta$  is a primitive  $r$ th root of unity, it appears at least as many times on top as on bottom. □

Hence, the number in  $(\spadesuit)$  is a rational algebraic integer, so it is an integer.