

# USAMO 2021/5

Evan Chen

TWITCH SOLVES ISL

Episode 65

## Problem

Let  $n \geq 4$  be an integer. Find all positive real solutions to the following system of  $2n$  equations:

$$\begin{array}{ll} a_1 = \frac{1}{a_{2n}} + \frac{1}{a_2}, & a_2 = a_1 + a_3, \\ a_3 = \frac{1}{a_2} + \frac{1}{a_4}, & a_4 = a_3 + a_5, \\ a_5 = \frac{1}{a_4} + \frac{1}{a_6}, & a_6 = a_5 + a_7, \\ \vdots & \vdots \\ a_{2n-1} = \frac{1}{a_{2n-2}} + \frac{1}{a_{2n}}, & a_{2n} = a_{2n-1} + a_1. \end{array}$$

## Video

<https://youtu.be/9WNgDETH0II>

## External Link

<https://aops.com/community/p21498967>

## Solution

The answer is that the only solution is  $(1, 2, 1, 2, \dots, 1, 2)$  which works.

We will prove  $a_{2k}$  is a constant sequence, at which point the result is obvious.

**First approach (Andrew Gu).** Apparently, with indices modulo  $2n$ , we should have

$$a_{2k} = \frac{1}{a_{2k-2}} + \frac{2}{a_{2k}} + \frac{1}{a_{2k+2}}$$

for every index  $k$  (this eliminates all  $a_{\text{odd}}$ 's). Define

$$m = \min_k a_{2k} \quad \text{and} \quad M = \max_k a_{2k}.$$

Look at the indices  $i$  and  $j$  achieving  $m$  and  $M$  to respectively get

$$\begin{aligned} m &= \frac{2}{m} + \frac{1}{a_{2i-2}} + \frac{1}{a_{2i+2}} \geq \frac{2}{m} + \frac{1}{M} + \frac{1}{M} = \frac{2}{m} + \frac{2}{M} \\ M &= \frac{2}{M} + \frac{1}{a_{2j-2}} + \frac{1}{a_{2j+2}} \leq \frac{2}{M} + \frac{1}{m} + \frac{1}{m} = \frac{2}{M} + \frac{2}{m}. \end{aligned}$$

Together this gives  $m \geq M$ , so  $m = M$ . That means  $a_{2i}$  is constant as  $i$  varies, solving the problem.

**Second approach (author's solution).** As before, we have

$$a_{2k} = \frac{1}{a_{2k-2}} + \frac{2}{a_{2k}} + \frac{1}{a_{2k+2}}$$

The proof proceeds in three steps.

- Define

$$S = \sum_k a_{2k}, \quad \text{and} \quad T = \sum_k \frac{1}{a_{2k}}.$$

Summing gives  $S = 4T$ . On the other hand, Cauchy-Schwarz says  $S \cdot T \geq n^2$ , so  $T \geq \frac{1}{2}n$ .

- On the other hand,

$$1 = \frac{1}{a_{2k-2}a_{2k}} + \frac{2}{a_{2k}^2} + \frac{1}{a_{2k}a_{2k+2}}$$

Sum this modified statement to obtain

$$n = \sum_k \left( \frac{1}{a_{2k}} + \frac{1}{a_{2k+2}} \right)^2 \stackrel{\text{QM-AM}}{\geq} \frac{1}{n} \left( \sum_k \frac{1}{a_{2k}} + \frac{1}{a_{2k+2}} \right)^2 = \frac{1}{n} (2T)^2$$

So  $T \leq \frac{1}{2}n$ .

- Since  $T \leq \frac{1}{2}n$  and  $T \geq \frac{1}{2}n$ , we must have equality everywhere above. This means  $a_{2k}$  is a constant sequence.

**Remark.** The problem is likely intractable over  $\mathbb{C}$ , in the sense that one gets a high-degree polynomial which almost certainly has many complex roots. So it seems likely that most solutions must involve some sort of inequality, using the fact we are over  $\mathbb{R}_{>0}$  instead.