

# Shortlist 2012 N3

Evan Chen

TWITCH SOLVES ISL

Episode 70

## Problem

Determine all integers  $m \geq 2$  such that every  $n$  with  $\frac{m}{3} \leq n \leq \frac{m}{2}$  divides the binomial coefficient  $\binom{n}{m-2n}$ .

## Video

<https://youtu.be/zVmGnDNw6LE>

## External Link

<https://aops.com/community/p3156840>

## Solution

The answer is  $m$  prime. The canonical way to think of this is:

**Claim.** The number  $m$  works if and only if  $\gcd(n, m) = 1$  for all  $n \in [m/3, m/2]$ .

*Proof.* Fix  $m$ , and choose any prime  $p$ . The divisibility is rewritten as

$$\nu_p((n-1)!) \geq \nu_p((m-2n)!) + (\nu_p(2n)!).$$

Define

$$f(n, q) = \left\lfloor \frac{n-1}{q} \right\rfloor - \left\lfloor \frac{m-2n}{q} \right\rfloor - \left\lfloor \frac{3n-m-1}{q} \right\rfloor.$$

Then we have

$$\nu_p\left(\binom{n}{m-2n}\right) = \sum_{e \geq 1} f(n, p^e).$$

We want this sum to be nonnegative for any prime  $p$ .

The following properties are true:

- $f(n, q) \geq 0$  if  $q \nmid n$ , since then  $\left\lfloor \frac{n-1}{q} \right\rfloor = \left\lfloor \frac{n}{q} \right\rfloor$ . So there are no issues when  $p \nmid n$ .
- Now suppose  $q$  is a prime power of a prime  $p$  dividing  $n$ . We may explicitly compute

$$\begin{aligned} f(qk, q) &= (k-1) - \left( \left\lfloor \frac{m-2(qk)}{q} \right\rfloor + \left\lfloor \frac{3(qk)-m-1}{q} \right\rfloor \right) \\ &= -1 - \left\lfloor \frac{m}{q} \right\rfloor - \left\lfloor \frac{-(m+1)}{q} \right\rfloor \\ &= \begin{cases} -1 & q \mid m \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

In that case, it follows that  $\nu_p\left(\binom{n}{m-2n}\right)$  is the sum of some non-positive terms, at least one of which is  $-1$  (namely the first term  $f(n, p^1) = -1$ .) So  $m$  fails.

In conclusion,  $m$  works if and only if  $n$  never has a common prime factor with  $m$ , as desired.  $\square$