

# IMO 1999/6

Evan Chen

TWITCH SOLVES ISL

Episode 91

## Problem

Find all the functions  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that

$$f(x - f(y)) = f(f(y)) + xf(y) + f(x) - 1$$

for all  $x, y \in \mathbb{R}$ .

## Video

[https://youtu.be/r1MW65i\\_VtI](https://youtu.be/r1MW65i_VtI)

## External Link

<https://aops.com/community/p131856>

## Solution

The answer is  $f(x) = -\frac{1}{2}x^2 + 1$  which obviously works.

For the other direction, first note that

$$P(f(y), y) \implies 2f(f(y)) + f(y)^2 - 1 = f(0).$$

We introduce the notation  $c = \frac{f(0)-1}{2}$ , and  $S = \text{img } f$ . Then the above assertion says

$$f(s) = -\frac{1}{2}s^2 + (c+1).$$

Thus, the given functional equation can be rewritten as

$$Q(x, s) : f(x-s) = -\frac{1}{2}s^2 + sx + f(x) - c.$$

**Claim** (Main claim). We can find a function  $g: \mathbb{R} \rightarrow \mathbb{R}$  such that

$$f(x-z) = zx + f(x) + g(z). \quad (\spadesuit).$$

*Proof.* If  $z \neq 0$ , the idea is to fix a nonzero value  $s_0 \in S$  (it exists) and then choose  $x_0$  such that  $-\frac{1}{2}s_0^2 + s_0x_0 - c = z$ . Then,  $Q(x_0, s)$  gives an pair  $(u, v)$  with  $u - v = z$ .

But now for any  $x$ , using  $Q(x+v, u)$  and  $Q(x, -v)$  gives

$$\begin{aligned} f(x-z) - f(x) &= f(x-u+v) - f(x) = f(x+v) - f(x) + u(x+v) - \frac{1}{2}u^2 + c \\ &= -vx - \frac{1}{2}s^2 - c + u(x+v) - \frac{1}{2}u^2 + c \\ &= -vx - \frac{1}{2}v^2 + u(x+v) - \frac{1}{2}u^2 = zx + g(z) \end{aligned}$$

where  $g(z) = -\frac{1}{2}(u^2 + v^2)$  depends only on  $z$ . □

Now, let

$$h(x) := \frac{1}{2}x^2 + f(x) - (2c+1),$$

so  $h(0) = 0$ .

**Claim.** The function  $h$  is additive.

*Proof.* We just need to rewrite  $(\spadesuit)$ . Letting  $x = z$  in  $(\spadesuit)$ , we find that actually  $g(x) = f(0) - x^2 - f(x)$ . Using the definition of  $h$  now gives

$$h(x-z) = h(x) + h(z). \quad \square$$

To finish, we need to remember that  $f$ , hence  $h$ , is known on the image

$$S = \{f(x) \mid x \in \mathbb{R}\} = \left\{ h(x) - \frac{1}{2}x^2 + (2c+1) \mid x \in \mathbb{R} \right\}.$$

Thus, we derive

$$h\left(h(x) - \frac{1}{2}x^2 + (2c+1)\right) = -c \quad \forall x \in \mathbb{R}. \quad (\heartsuit)$$

We can take the following two instances of  $\heartsuit$ :

$$\begin{aligned}h(h(2x) - 2x^2 + (2c + 1)) &= -c \\h(2h(x) - x^2 + 2(2c + 1)) &= -2c.\end{aligned}$$

Now subtracting these and using  $2h(x) = h(2x)$  gives

$$c = h(-x^2 - (2c + 1)).$$

Together with  $h$  additive, this implies readily  $h$  is constant. That means  $c = 0$  and the problem is solved.