# IMO 1998/3 Evan Chen

TWITCH SOLVES ISL

Episode 94

#### Problem

For any positive integer n, let  $\tau(n)$  denote the number of its positive divisors (including 1 and itself). Determine all positive integers m for which there exists a positive integer n such that

$$\frac{\tau(n^2)}{\tau(n)} = m.$$

### Video

https://youtu.be/mKfgxyfdn2w

## **External Link**

https://aops.com/community/p124439

#### Solution

The answer is odd integers m only. If we write  $n = p_1^{e_1} \dots p_k^{e_k}$  we get

$$\prod \frac{2e_i+1}{e_i+1} = m.$$

It's clear now that m must be odd, since every fraction has odd numerator.

We now endeavor to construct odd numbers. The proof is by induction, in which we are curating sets of fractions of the form  $\frac{2e+1}{e+1}$  that multiply to a given target. The base cases are easy to verify by hand. Generally, assume  $p = 2^t k - 1$  is odd, where

k is odd. Then we can write

$$\frac{2^{2t}k - 2^{t}(k+1) + 1}{2^{2t-1}k - 2^{t-1}(k+1) + 1} \cdot \frac{2^{2t-1}k - 2^{t-1}(k+1) + 1}{2^{2t-2}k - 2^{t-2}(k+1) + 1} \cdots \frac{2^{t+1}k - 2(k+1) + 1}{2^{t}k - 2^{0}(k+1) + 1}.$$

Note that  $2^{2t}k - 2^t(k+1) + 1 = (2^tk - 1)(2^t - 1)$ , and  $2^tk - k = k(2^t - 1)$ , so the above fraction simplifies to

$$\frac{2^t k - 1}{k}$$

meaning we just need to multiply by k, which we can do using induction hypothesis.