

# gcd det

Evan Chen

TWITCH SOLVES ISL

Episode 97

## Problem

Let  $n$  be a given positive integer. Find the determinant of the  $n \times n$  matrix whose  $(i, j)$ th entry is  $\gcd(i, j)$ .

## Video

<https://youtu.be/SnEj4CljU18>

## Solution

The answer is the product  $\varphi(1)\varphi(2)\dots\varphi(n)$ , where  $\varphi$  is the Euler phi function.

The proof is by induction on  $n$ , with base cases immediate. The main claim is that we can eliminate nearly all of the entries in the last column by doing column operations. To be precise:

**Claim.** Let  $n = p_1^{e_1} \dots p_k^{e_k}$ . Then for any integer  $1 \leq m \leq n$ , we have

$$\sum_{S \subseteq \{1,2,\dots,k\}} (-1)^{|S|} \gcd\left(m, \frac{n}{\prod_{i \in S} p_i}\right) = \begin{cases} 0 & m < n \\ \varphi(n) & m = n. \end{cases}$$

Hence, if the claim is true, then we can take the  $\frac{n}{\prod_{i \in S} p_i}$ 'th columns for each nonempty  $S$  and add/subtract them from the  $n$ th column (subtraction if  $|S|$  is odd, addition if  $|S|$  is even). This would leave all zeros except  $\varphi(n)$  in the bottom right. Since we didn't touch any of the other rows or columns, the original problem is solved by induction.

*Proof of claim.* Let  $m = p_1^{a_1} p_2^{a_2} \dots p_k^{a_k}$ . The sum can be factored in the following way:

$$\prod_{i=1}^k \left( p_i^{\min(a_i, e_i)} - p_i^{\min(a_i, e_i - 1)} \right).$$

If  $m = n$ , then  $a_i = e_i$  and the above product becomes the usual factorization of the Euler phi function.

On the other hand, if  $m < n$ , then there exists some particular index  $i$  such that  $a_i < e_i$ . That makes the corresponding term in the product zero, and so the whole product vanishes.  $\square$