

# bissue 2022/15

Evan Chen

TWITCH SOLVES ISL

Episode 120

## Problem

Quadrilateral  $ABCD$  is such that  $2\angle A = \angle B$  and  $2\angle D = \angle C$ . If  $AB = 3$ ,  $BC = 4$ , and  $CD = 5$ , find the area of quadrilateral  $ABCD$ .

## Video

<https://youtu.be/PBS0cmLosrw>

## External Link

<https://aops.com/community/p24130067>

## Solution


This problem and solution were contributed by Benny Wang.

Let  $E$  be the intersection of the angle bisector of  $\angle ABC$  with  $\overline{AD}$ ; let  $F$  be the intersection of the angle bisector of  $\angle BCD$  with  $\overline{AD}$ .

**Claim.** We have  $E = F$ .

*Proof.* Let the perpendiculars from  $E$  to  $\overline{AB}$  and  $\overline{BC}$  be  $E_1$  and  $E_2$ . Note that due to  $\triangle ABE$  isosceles with  $AE = BE$  (using the angle condition), it follows that  $E_1B = \frac{AB}{2} = 1.5$ . Then, since we defined  $\overline{BE}$  as the bisector of  $\angle ABC$ , we have that  $E_2B = E_1B = 1.5$ .


Similarly, if we define  $F_1$  and  $F_2$  as the perpendiculars from  $F$  to  $\overline{CD}$  and  $\overline{BC}$ , we have that  $F_2C = F_1C = \frac{CD}{2} = 2.5$ . Note that  $BE_2 + F_2C = 4 = BC$ , implying that  $E_2 = F_2$ , implying that  $E = F$ , as desired.  $\square$



Now we compute a bunch. Let  $G$  be the point  $E_2 = F_2$ , and let  $X$  be the point  $E = F$ .

Consider  $\triangle BXC$ . Then  $G$  is defined as the foot of the altitude from  $X$  to  $\overline{BC}$ ; we know  $BG = 1.5$  and  $GC = 2.5$ . We also know that  $\angle BXC = 60^\circ$  by some straightforward angle chasing. This is enough information to solve for  $\overline{BXC}$ ; in particular, we desire the length of  $XG$ ; then, since  $X$  is equidistant from  $\overline{AB}$ ,  $\overline{BC}$  and  $\overline{CD}$ , we could compute the area of  $ABCD$  to be  $\frac{(AB+BC+CD) \times XG}{2}$ .

To find  $XG$  is a straightforward geometry problem now. There are several ways we could go about this. Here is one: let  $M$  be the midpoint of  $\overline{BC}$ , and let  $O$  be the circumcenter of  $\triangle BXC$ .



Note that  $\angle BOC = 2\angle BXC = 120^\circ$ , so  $\triangle BMO$  and  $\triangle CMO$  are 30-60-90 triangles. Therefore,  $MO = \frac{2}{\sqrt{3}}$  and  $BO = OC = OX = \frac{4}{\sqrt{3}}$ . Now, note that

$$\begin{aligned} GX &= \sqrt{XO^2 - GM^2} + MO \\ &= \sqrt{\left(\frac{4}{\sqrt{3}}\right)^2 - 0.5^2} + \frac{2}{\sqrt{3}} \\ &= \sqrt{\frac{61}{12}} + \frac{4}{\sqrt{12}} \\ &= \frac{4 + \sqrt{61}}{\sqrt{12}}. \end{aligned}$$

Therefore,

$$[ABCD] = \frac{XG \cdot (AB + BC + CD)}{2} = \frac{4 + \sqrt{61}}{2\sqrt{12}} \cdot 12 = \boxed{4\sqrt{3} + \sqrt{183}}.$$