

# Longlist 1974/2

Evan Chen

TWITCH SOLVES ISL

Episode 121

## Problem

Let  $u_n$  be the Fibonacci sequence, i.e.,  $u_0 = 0$ ,  $u_1 = 1$ ,  $u_n = u_{n-1} + u_{n-2}$  for  $n > 1$ . Prove that there exist infinitely many prime numbers  $p$  that divide  $u_{p-1}$ .

## Video

<https://youtu.be/QS7jBXIVf8s>

## External Link

<https://aops.com/community/p2136111>

## Solution

Let  $\alpha = \frac{1}{2}(1 + \sqrt{5})$  and  $\beta = \frac{1}{2}(1 - \sqrt{5})$ . We have Binet's formula:

$$u_n = \frac{\alpha^n - \beta^n}{\alpha - \beta}.$$

In fact, if  $p \equiv 1 \pmod{5}$ , then by quadratic reciprocity, 5 is a quadratic residue, so  $\alpha$  and  $\beta$  can be viewed as elements of  $\mathbb{F}_p$ , and Fermat's theorem applies to give  $\alpha^{p-1} \equiv \beta^{p-1} \equiv 1 \pmod{p}$ , as desired.

**Remark.** In fact, if  $p \equiv \pm 2 \pmod{5}$  and  $p > 2$  then  $\alpha^p = \beta$  and  $\beta^p = \alpha$ , so in this case

$$u_{p-1} \equiv \frac{\frac{\beta}{\alpha} - \frac{\alpha}{\beta}}{\alpha - \beta} = -\frac{\alpha + \beta}{\alpha\beta} = -\frac{-1}{1} = 1 \pmod{p}.$$