

Marcin E Kuczma

Evan Chen

TWITCH SOLVES ISL

Episode 122

Problem

Find all functions f taking integer-lattice vectors in \mathbb{Z}^2 to \mathbb{R} such that $f(\mathbf{a} + \mathbf{b}) = f(\mathbf{a}) + f(\mathbf{b})$ whenever $\mathbf{a} \perp \mathbf{b}$.

Video

<https://youtu.be/2mn3De0yc8A>

External Link

<https://aops.com/community/p27461531>

Solution

Abbreviate $f(\langle x, y \rangle)$ to just $f(x, y)$. Note that the set of solutions is obviously an \mathbb{R} -vector space. Consider the four solutions:

$$\begin{aligned} f_1(x, y) &:= x \\ f_2(x, y) &:= y \\ f_3(x, y) &:= x^2 + y^2 \\ f_4(x, y) &:= (x\%2) - (y\%2). \end{aligned}$$

(Verification left as exercise.) They are linearly independent, so any linear combination of this works, and we have found a 4-dimensional space of solutions.

We now show conversely that the dimension of the vector space of solutions is at most 4. To begin, first note that

$$f(x, y) = f(x, 0) + f(0, y).$$

In particular, $f(0, 0) = 0$. Next, note that for all $n \geq 1$

$$\begin{aligned} f(n, 0) + f(0, -1) + f(1, 0) + f(0, n) &= f(n, -1) + f(1, n) \\ &= f(n + 1, n - 1) \\ &= f(n + 1, 0) + f(0, n - 1) \\ \implies f(n + 1, 0) &= f(n, 0) + f(0, -1) + f(1, 0) + f(0, n) - f(0, n - 1) \end{aligned}$$

By writing the analogous equations for $f(-(n + 1), 0)$, $f(0, n + 1)$ and $f(0, -(n + 1))$, it follows that all the values of f are determined by these recursions as soon as the “initial” values of $f(1, 0)$, $f(-1, 0)$, $f(0, 1)$, $f(0, -1)$ are chosen. Ergo, the space of valid f is at most four-dimensional, as needed.