

Twitch 125.2

Evan Chen

TWITCH SOLVES ISL

Episode 125

Problem

Regular hexagon $ABCDEF$ is drawn, and a point P inside this hexagon satisfies $[PAB] = 6$, $[PCD] = 10$, and $[PEF] = 11$. Compute $[PBC][PDE][PFA]$.

Video

<https://youtu.be/3DdyaBnRYPN30>

Solution

Let x denote the side length of the hexagon, so the heights from P to AB , CD , EF are $\frac{12}{x}$, $\frac{20}{x}$, $\frac{22}{x}$ respectively. Let h_1 , h_2 , h_3 be the heights from P to BC , DE , FA ; then

$$\begin{aligned}\sqrt{3}x &= \frac{12}{x} + h_1 \\ &= \frac{20}{x} + h_2 \\ &= \frac{22}{x} + h_3.\end{aligned}$$

On the other hand, by considering the equilateral triangles formed by extending the sides of the hexagon (or quoting Viviani theorem),

$$h_1 + h_2 + h_3 = \frac{12}{x} + \frac{20}{x} + \frac{22}{x}.$$

Hence, it follows that

$$3 \cdot \sqrt{3}x = 2 \left(\frac{12}{x} + \frac{20}{x} + \frac{22}{x} \right)$$

so $x^2 = 12\sqrt{3}$.

To finish up, compute

$$\begin{aligned}[PDE] &= \frac{1}{2}x \cdot h_1 &&= \frac{\sqrt{3}}{2}x^2 - 6 = 12 \\ [PFA] &= \frac{1}{2}x \cdot h_2 &&= \frac{\sqrt{3}}{2}x^2 - 10 = 8 \\ [PBC] &= \frac{1}{2}x \cdot h_3 &&= \frac{\sqrt{3}}{2}x^2 - 11 = 7\end{aligned}$$

giving answer $12 \cdot 8 \cdot 7 = 672$.