

# JMO 2024/4

Evan Chen

TWITCH SOLVES ISL

Episode 142

## Problem

Let  $n \geq 3$  be an integer. Rowan and Colin play a game on an  $n \times n$  grid of squares, where each square is colored either red or blue. Rowan is allowed to permute the rows of the grid and Colin is allowed to permute the columns. A grid coloring is orderly if:

- no matter how Rowan permutes the rows of the coloring, Colin can then permute the columns to restore the original grid coloring; and
- no matter how Colin permutes the columns of the coloring, Rowan can then permute the rows to restore the original grid coloring.

In terms of  $n$ , how many orderly colorings are there?

## Video

[https://youtu.be/s\\_bUukq7HiA](https://youtu.be/s_bUukq7HiA)

## External Link

<https://aops.com/community/p30227193>

## Solution


The answer is  $2n! + 2$ . In fact, we can describe all the orderly colorings as follows:

- The all-blue coloring.
- The all-red coloring.
- Each of the  $n!$  colorings where every row/column has exactly one red cell.
- Each of the  $n!$  colorings where every row/column has exactly one blue cell.

These obviously work; we turn our attention to proving these are the only ones.

For the other direction, fix a orderly coloring  $\mathcal{A}$ .

Consider any particular column  $C$  in  $\mathcal{A}$  and let  $m$  denote the number of red cells that  $C$  has. Any row permutation (say  $\sigma$ ) that Rowan chooses will transform  $C$  into some column  $\sigma(C)$ , and our assumption requires  $\sigma(C)$  has to appear somewhere in the original assignment  $\mathcal{A}$ . An example for  $n = 7$ ,  $m = 2$ , and a random  $\sigma$  is shown below.



On the other hand, the number of possible patterns of  $\sigma(C)$  is easily seen to be exactly  $\binom{n}{m}$  — and they must all appear. In particular, if  $m \notin \{0, 1, n-1, n\}$ , then we immediately get a contradiction because  $\mathcal{A}$  would need too many columns (there are only  $n$  columns in  $\mathcal{A}$ , which is fewer than  $\binom{n}{m}$ ). Moreover, if either  $m = 1$  or  $m = n-1$ , these columns are all the columns of  $\mathcal{A}$ ; these lead to the  $2n!$  main cases we found before.

The only remaining case is when  $m \in \{0, n\}$  for every column, i.e. every column is monochromatic. It's easy to see in that case the columns must all be the same color.