

# Notes for 18.02 Recitation 13

## 18.02 Recitation MW9

EVAN CHEN

23 October 2024

*The world is quiet here.*

— Lemony Snicket, in *A Series of Unfortunate Events*

This handout (and any other DLC's I write) are posted at <https://web.evanchen.cc/1802.html>. (This was originally titled Recitation 14, but the numbering changed.)

### §1 Reading

You should read section 21 of LAMV for the full details, which don't fit on the page. The following are just excerpts for quick reference.

### §2 Recipes

#### ☰ Recipe for integrating over a rectangle

To integrate something of the form  $\int(\int dy) dx$ :

1. Evaluate the inner integral as in 18.01, treating  $x$  as constant.
2. You should get something only depending on  $x$ . Integrate it as in 18.01.

#### ☰ Recipe for converting to $xy$ -integration

1. Draw a picture of the region as best you can.
2. Write the region as a list of inequalities.<sup>1</sup>
3. Pick *one* of  $x$  and  $y$ , and use your picture to describe all the values it could take.
4. Solve for the *other* variable in all the inequalities.

### §3 Pictures for the example from Poonen's notes




#### 🔗 Sample Question

Show both ways of setting up an integral of a function  $f(x, y)$  over the region bounded by  $y - x = 2$  and  $y = x^2$ .

Here the region would be described as  $y \geq x^2$  and  $y - x \leq 2$ .

---

<sup>1</sup>I don't think other sources always write the inequalities the way I do. But I think this will help you a lot with making sure bounds go the right way.



### §4 Recitation questions from official course

1. Calculate the double integral of the function  $f(x, y) = 6x^2 + 2y$  over the rectangle  $R = [0, 2] \times [-1, 1]$ . Use both vertical and horizontal slicings and check you get the same answer.
2. Let  $R$  be the first-quadrant region bounded by the two curves  $y = \sqrt{x}$  and  $y = x^3$ . Compute in two different ways the double integral

$$\iint_R xy^2 \, dA.$$

3. Let  $R$  be the (bounded) region between the parabola  $y^2 = x$  and the line through  $(2, 0)$  having slope 1. Find the points where the curves intersect and describe the region  $R$  in terms of horizontal slices and vertical slices. Express the double integral  $\iint_R f(x, y) \, dA$  as an iterated integral in both ways, using both horizontal and vertical slicings. In the second case, you will have to write the integral in two pieces.